



# 86<sup>TH</sup> CONGRESS OF THE EUROPEAN ORTHODONTIC SOCIETY

**JUNE 15 - 19, 2010  
PORTOROŽ, SLOVENIA**



## ABSTRACT BOOK



<http://www.eos2010.si>

I FEEL  
SLOVENIA



**86<sup>TH</sup> CONGRESS OF THE  
EUROPEAN  
ORTHODONTIC  
SOCIETY**

**JUNE 15 - 19, 2010  
PORTOROŽ, SLOVENIA**

**ABSTRACT  
BOOK**



# CONTENTS

03	<b>Sheldon Friel Memorial Lecture</b>
04	<b>Houston Award</b>
04	Oral Presentations
06	Posters
09	<b>Keynote Lectures</b>
13	<b>Oral Presentations</b>
57	<b>Scientific posters</b>
57	Biotechnology and the Future in Orthodontics
69	Contemporary Diagnostics and Treatment Approaches
85	Class III Malocclusion
97	Free Topics
147	<b>Clinical posters</b>
147	Biotechnology and the Future in Orthodontics
149	Contemporary Diagnostics and Treatment Approaches
155	Class III Malocclusion
165	Free Topics
175	<b>Index of Authors</b>

## Legend

Colour	Topics
	Sheldon Friel Memorial Lecture WJB Houston Research Awards - Oral Presentations & Posters Keynote Lectures
	Biotechnology and the Future in Orthodontics
	Contemporary Diagnostics and Treatment Approaches
	Class III Malocclusion
	Free Topics

NB: The authors of abstracts marked \*\*\* have either declared a financial interest or have not indicated whether or not they have a financial interest



## SHELDON FRIEL MEMORIAL LECTURE

Wednesday, June 16 - at 8:30

### THE FACIAL PROFILE – CHANGES WITH AND WITHOUT TREATMENT

**Urban Hägg**, Hong Kong

Orthodontists pay attention to the patient's face in profile when they plan and evaluate treatment. Such treatment planning and evaluation were in the past – and still are today – commonly based on estimates of changes in the patient's hard and soft tissue profiles after certain angular and linear measurements have been made on a series of lateral cephalograms. Is this approach good enough?

An individual's appearance and face in profile are affected by the position of the mandible. But the final position of the mandible depends on bone remodelling, the magnitude and direction of growth of the condyle, maxilla, and other parts of the facial complex, as well as muscles and ligaments. Can the position of the mandible be influenced by orthodontic devices?

For patients who have a skeletal Class II malocclusion with mandibular retrognathism, treatment usually aims to 'position' the mandible forward. Orthodontics, dentofacial orthopaedics, and orthognathic surgery are the three main treatment modalities currently used to 'improve' the profile of such patients. How effective are these modalities?

Orthodontic treatment can focus on dental displacement only, or can deal with the control and modification of facial growth...defined as 'dentofacial orthopaedics' (Wikipedia, 2008). In recent randomized controlled clinical trials studying the treatment of skeletal Class II malocclusions, conventional cephalometric analyses showed that the growth modification effect, if any, of using dentofacial orthopaedic devices did not exceed the effect of using comprehensive orthodontics only. Is 'control and modification of facial growth' a clinical reality?

Recent prospective long-term studies on sagittal-split osteotomies have demonstrated that mandibular advancement can be followed by marked relapse. Does this indicate that there might be a 'natural limit' of forward mandibular positioning?

Randomized controlled clinical trials and prospective studies on orthodontics can be riddled with flaws, such as inappropriate sampling, operators' attitudes and level of experience, choice of treatment devices and retention protocol, timing and length of treatment, and choice of methods of analysis. Are there any methods that could allow more valid estimates of facial changes? Will there be any methods that really control and regulate jaw growth, in the foreseeable future? In this lecture, an attempt to will be made to address these questions and provide clinical guidance based on science, a dose of common sense, and clinical preference.



## Oral Presentations

### HAO 01

Wednesday, June 16 - at 9:20

#### RAPID VERSUS SLOW MAXILLARY EXPANSION: A THREE-DIMENSIONAL RANDOMIZED CLINICAL TRIAL

**I Cioffi**<sup>1</sup>, G Cordasco<sup>2</sup>, P Manzo<sup>1</sup>, R Nucera<sup>2</sup>, R Martina<sup>1</sup>, <sup>1</sup>Department of Oral Sciences, Section of Orthodontics and Temporomandibular Disorders, University of Naples Federico II and <sup>2</sup>School of Dentistry, University of Messina, Italy

**AIM:** A posterior crossbite is a common malocclusion in children, and is generally treated to alleviate dental crowding, to prevent the developing of mandibular skeletal asymmetries in adulthood and to reduce the risk of developing temporomandibular disorders. A PXB is mostly the result of skeletal deficiency of the upper jaw and is often corrected by means of by orthopaedic maxillary expansion. The purpose of this study was to evaluate the skeletal and dentoalveolar effects of rapid maxillary expansion (RME) as compared with to slow maxillary expansion (SME) by means of low dose computed tomography. It was hypothesized that RME was more effective in treating a constricted maxilla because of major orthopaedic effects as compared with SME.

**SUBJECTS AND METHOD:** The subjects were treated by means of a banded palatal expander and were randomly allocated to two treatment groups: 11 subjects (8 males, mean age  $\pm$  SD=10.8  $\pm$  2.9; 3 females, mean age  $\pm$  SD = 9.3  $\pm$  0.4) were allocated to SME group and 15 subjects to the RME group (6 males mean age  $\pm$  SD = 9.6  $\pm$  1.6; 9 females, mean age  $\pm$  SD = 10.3  $\pm$  1.2). A low dose three-dimensional computed tomography protocol was used to identify skeletal and dental landmarks. Data were also extracted from dental casts.

**RESULTS:** Maxillary expansion resulted in an increase of dental and skeletal transverse diameters in both the RME and SME groups. No statistical difference in expansion at different tooth levels was found between groups, with the exception of molar expansion measured at the molar apices, which was significantly lower in the RME group ( $P = 0.02$ ) than in the SME group. Skeletal expansion measured at different bony locations did not differ significantly between the groups.

**CONCLUSION:** RME was not more effective than SME in orthopaedic expansion of the upper jaw.

### HAO 02

Wednesday, June 16 - at 9:30

#### THE EFFECTS OF PLAYING A WIND INSTRUMENT ON THE OCCLUSION

**E Grammatopoulos**<sup>1</sup>, A White<sup>2</sup>, A Dhoptkar<sup>3</sup>, <sup>1</sup>Birmingham Dental Hospital, <sup>2</sup>Department of Statistics and <sup>3</sup>School of Dental Sciences, University of Birmingham, England

**AIM:** Dentists and orthodontists are often asked by patients and their parents whether playing a wind instrument may affect the position of their teeth. Indeed, amongst wind instrument players and teachers it has been a popular belief that playing a wind instrument leads to malocclusion. The aim of this study was to investigate the effects of playing a wind instrument on the occlusion

**SUBJECTS AND METHOD:** This was a cross-sectional observational study. One hundred and seventy professional musicians were selected from 21 classical orchestras and organisations. The subjects were subdivided according to the type of instrument mouthpiece and included: 32 large cup-shaped mouthpiece brass players (group AL), 42 small cup-shaped mouthpiece brass players (group AS), 37 single reed mouthpiece woodwind players (group B) and 59 string and percussion

instrument players (control group). Impressions were taken for each subject and various parameters were assessed from the study casts. Statistical analysis was undertaken for interval variables with one-way analysis of variance and for categorical variables with chi-square tests

**RESULTS:** No statistically significant differences were found in overjet, overbite, crowding, Little's Irregularity Index or prevalence of incisor classification between the wind instrument players and the control group ( $P > 0.05$ ). However, group AL had a significantly higher prevalence of buccal crossbites than all other groups ( $P < 0.05$ ).

**CONCLUSION:** Playing a wind instrument does not significantly influence the position of the anterior teeth and is not a major aetiological factor in the development of malocclusion. However, playing a brass instrument with a large cup-shaped mouthpiece may predispose to buccal crossbite development.

### HAO 03

Wednesday, June 16 - at 9:40

#### ALVEOLAR CREST HEIGHT AFTER DISTRACTION OF THE MANDIBULAR ANTERIOR ALVEOLAR SEGMENT

**C Joss**<sup>1</sup>, M Antonini<sup>2</sup>, A Triaca<sup>2</sup>, A M Kuijpers-Jagtman<sup>1</sup>, S Kiliaridis<sup>3</sup>,  
<sup>1</sup>Department of Orthodontics and Oral Biology, Radboud University Nijmegen, Netherlands, <sup>2</sup>Pyramide Klinik, Zürich, and <sup>3</sup>Department of Orthodontics, University of Geneva, Switzerland

**AIM:** To compare anterior alveolar crest height in consecutive patients receiving (i) anterior mandibular alveolar distraction osteogenesis combined with orthodontic treatment (DOG group), (ii) non-extraction orthodontic treatment (non-ex group), and (iii) lower premolar extraction orthodontic treatment (ex group) to resolve mandibular space deficiency.

**SUBJECTS AND METHOD:** The ex group contained 18 patients, the non-ex group 18 patients and the DOG group 19 subjects. Study casts and lateral cephalograms were retrospectively analysed before (T1) and clinically re-examined at a mean of 3.9 (DOG group), 5.0 (ex group) and 4.6 years (non-ex group) after orthodontic treatment (T4). Cone beam computed tomograms were available only for T4 and were used to measure lower incisor buccal and lingual alveolar crest height.

**RESULTS:** The DOG group showed a significant lower buccal (ex-group  $P = 0.038$ ) and lingual alveolar crest height (non-ex group  $P = 0.013$ ; ex group  $P = 0.005$ ). No difference regarding alveolar crest height was found between the non-ex and ex groups even though the lower incisors were significantly more proclined in the non-ex group at T4. Age and alveolar process depth were the only variables significantly correlated with lower alveolar crest level in the DOG group while the inclination of lower incisors or the amount of surgical advancement at point B or incision inferior showed no correlation. Age and decrease in alveolar process depth (T4-T1) in the ex group and proclination of the lower incisors at T4 in the non-ex group were significantly correlated with lower buccal alveolar crest level.

**CONCLUSION:** Lingual and buccal alveolar bone loss as a long-term consequence of DOG treatment cannot be excluded, but increased age in combination with pre-existing periodontal problems should be considered as the main source for alveolar bone loss. A smaller depth of the alveolar process resulted in more alveolar bone dehiscence on the lingual and buccal side in the DOG group.

### HAO 04

Wednesday, June 16 - at 9:50

#### GINGIVAL CREVICULAR FLUID ALKALINE PHOSPHATE ACTIVITY AS A MARKER OF SKELETAL MATURATION

**G Perinetti**<sup>1</sup>, T Baccetti<sup>2</sup>, <sup>1</sup>Department of Biomedicine, Unit of Orthodontics, University of Trieste and <sup>2</sup>Department of Orthodontics, University of Florence, Italy

**AIM:** Treatment timing plays a significant role in the outcome of orthopaedic treatment of dentoskeletal disharmonies in growing patients. Therefore, correct identification of the different phases in skeletal maturation represents a crucial issue in orthodontic diagnosis and treatment planning. The present study aimed to evaluate gingival crevicular fluid (GCF) alkaline phosphatase activity, a key enzyme in bone mineralization, in growing subjects in relation to chronological age, phase of the dentition, and stage in individual skeletal growth.

**SUBJECTS AND METHOD:** Forty-nine healthy subjects (30 females, 19 males,

7.8-17.0 years of age). After appraisal of the dentition phase and skeletal maturation using the cervical vertebral method. GFC was collected at both the maxilla and the mandible at periodontally healthy sites around the central incisors. Enzymatic activity was determined spectrophotometrically. **RESULTS:** GCF alkaline phosphatase activity when collected from the mandible was correlated with the cervical vertebral stages. In particular, a peak in enzyme activity was seen at the pubertal stage CS 3. On the contrary, phases of the dentition and chronological age showed either no significant or very low correlations, respectively, with enzyme activity. The pubertal increase in the enzyme activity from the mandible may have clinical value when planning orthodontic treatment in growing subjects with dentoskeletal disharmonies. **CONCLUSION:** GCF alkaline phosphatase activity may be a valid candidate as a new, biochemical and non-invasive, marker of skeletal maturation, i.e. mandibular growth spurt, in periodontally healthy subjects scheduled for orthodontic treatment during adolescence



## HOUSTON AWARD

### Posters

#### HAP 01

*Allways on display*

#### **PREDICTIVE VALUE OF MOLAR BITE FORCE ON CLASS II FUNCTIONAL APPLIANCE TREATMENT OUTCOMES**

**G Antonarakis<sup>1</sup>**, H Kjellberg<sup>2</sup>, S Kiliaridis<sup>1</sup>, Departments of Orthodontics, Dental Schools, <sup>1</sup>University of Geneva, Switzerland and <sup>2</sup>Göteborg University, Sweden

**AIMS:** Sagittal intermaxillary changes using functional appliances are unpredictable, showing large interindividual variation. One factor for these differences may be the masticatory musculature and its functional capacity, which varies widely between growing individuals. The aims of this research were to investigate changes in maximal molar bite force during functional appliance treatment, and to assess the influence of pre-treatment maximal molar bite force on treatment outcomes with functional appliances used in Class II malocclusion children.

**SUBJECTS AND METHOD:** Twenty-five children, aged 9-13 years, with a skeletal and dental Class II relationship and increased overjet were treated with functional appliances for 1-2 years. Dental casts, lateral cephalograms, maximal molar bite force, and finger force measurements were performed before and after treatment. These same measurements were also performed 1-2 years before treatment; the intermediate period before starting treatment serving as a control. Multiple regression analysis was used to determine possible correlations between initial maximal molar bite force and dental or cephalometric changes during treatment.

**RESULTS:** Maximal molar bite force increased pre-treatment, but decreased during functional appliance treatment. Children treated with functional appliances who had a weaker pre-treatment maximal molar bite force showed a larger overjet reduction, greater improvement in molar relationship, a greater reduction in the ANB angle, and a greater augmentation in SNB angle during treatment.

**CONCLUSION:** Treatment of a dental Class II relationship with functional appliances seems to lead to more favourable treatment outcomes in children with a weaker maximal molar bite force. This is observed both as regards improvements in the dental sagittal relationships, namely overjet and molar Class, as well as skeletal changes detected by a decrease in ANB and an increase in SNB angle.

### COMPARISON BETWEEN THE INDEX OF COMPLEXITY OUTCOME AND NEED AND THE INDEX OF ORTHODONTIC TREATMENT NEED

**A Borzabadi-Farahani**<sup>1</sup>, A Borzabadi-Farahani<sup>2</sup>, F Eslamipour<sup>3</sup>,

<sup>1</sup>Department of Orthodontics, Birmingham Dental Hospital, England,

<sup>2</sup>Private Practice, Tehran and <sup>3</sup>Department of Community Dentistry, Isfahan University of Medical Sciences, Iran

**AIM:** A cross-sectional study to investigate the agreement between assessments of orthodontic treatment need using the Index of Complexity, Outcome and Need (ICON) and the Index of Orthodontic Treatment Need (IOTN).

**SUBJECTS AND METHOD:** Five hundred and two subjects (253 female, 249 males, aged 11-14 years), including those who were wearing an orthodontic appliance (1 female and 5 males). ICON and IOTN [Dental Health Component (DHC) and Aesthetic Component (AC)] scores were recorded in those not undergoing treatment. The percentage of subjects needing treatment (ICON >43) and different complexity compartments of the ICON were compared in different genders. Observed percentage agreement and kappa statistics were used to analyse the agreement between different components of the IOTN (AC and DHC) and also the ICON and DC and AC of the IOTN dichotomized into yes or no categories of orthodontic treatment need.

**RESULTS:** No gender differences were found for treatment need (ICON >43,  $P > 0.05$ ) or treatment complexity ( $P > 0.05$ ). Kappa statistics for diagnostic agreement between the DHC and AC of the IOTN was 0.55 (95% CI 0.48-0.63) and for diagnostic agreement between the ICON and AC and DHC of the IOTN 0.40 (95% CI 0.33-0.46) and 0.78 (95% CI 0.73-0.83), respectively. The observed agreement between the DHC and AC of the IOTN was 81.8 per cent. The observed agreement between the ICON and AC and DHC of the IOTN were 71.3 and 89.5 per cent, respectively.

**CONCLUSION:** In terms of orthodontic treatment need, there was moderate agreement between ICON and AC (IOTN) and between DHC and AC, respectively. There was good agreement between the ICON and the DHC of the IOTN. ICON is a good substitute for the DHC of ION.

### TREATMENT OF PALATALLY DISPLACED CANINES WITH RAPID MAXILLARY EXPANSION AND/OR A TRANSPALATAL ARCH

**C Masucci**, T Baccetti, Università degli Studi di Firenze, Italy

**AIM:** To investigate the effect of rapid maxillary expansion (RME) and/or transpalatal arch (TPA) therapy in combination with primary canine extraction on the eruption rate of palatally displaced canines by means of a randomized clinical study.

**SUBJECTS AND METHOD:** One hundred and twenty subjects were enrolled based on palatally displaced canines (PDC) diagnosed on panoramic radiographs. They were randomly assigned to one of four groups: three treatment groups (RME followed by TPA therapy plus extraction of primary canines, RME/TPA/EC group, 40 subjects; TPA therapy plus extraction of primary canines, TPA/EC group, 25 subjects; extraction of primary canines, EC group, 25 subjects). A control group (CG; 30 subjects) received no orthodontic treatment. All subjects were evaluated before treatment at the initial observation (T1) and then re-evaluated in the early permanent dentition (T2) to assess the rate of successful eruption of PDCs. Panoramic radiographs and dental casts at T1 and the prevalence rates of successful cases in the three treatment groups compared with the CG at T2 were statistically evaluated. Predictive features at T1 for successful canine eruption were determined in the three treatment groups.

**RESULTS:** No significant difference was found for any measurement at T1. Three dropouts were recorded. The prevalence of canine eruption was 80 per cent for the RME/TPA/EC group, 79 per cent for the TPA/EC group, 62.5 per cent for the EC group, versus 28 per cent in the CG, with statistically significant differences between all groups, with the exception of the comparison between RME/TPA/EC and TPA/EC. Predictive pre-treatment variables for the success of treatment were: less severe sectors of canine displacement, prepubertal stages of skeletal maturity, and an open root apex of the PDC.

**CONCLUSION:** The use of a TPA without RME can be equally effective as RME/TPA in PDC cases not requiring maxillary expansion, thus reducing the burden of treatment for the patient.

### EFFECTS OF VASCULAR ENDOTHELIAL GROWTH FACTOR ON MC3T3-E1 CELL LINE

**Y Y Tan**, Y-Q Yang, L Chai, R W K Wong, A B M Rabie, Orthodontic Department, University of Hong Kong, SAR China

**AIM:** Bone support and blood supply are necessary for effective orthodontic tooth movement. Osteogenesis and angiogenesis are closely correlated. Vascular endothelial growth factor (VEGF) is believed to play a critical role in skeletal development including mandibular condylar growth, by enhancement of angiogenesis. This aim of this study was to investigate whether VEGF has any direct effect on increasing bone cell activity, in an attempt to better understand how VEGF promotes bone remodelling.

**MATERIALS AND METHOD:** A preosteoblastic cell line, MC3T3-E1, was cultured with and without VEGF in vitro. The cells in both the control and test groups were collected at different culture time points of 24, 48 and 72 hours ( $n = 3$  for each time point). Real-time polymerase chain reaction (PCR) was carried out to quantify the expressions of two markers related to bone formation [alkaline phosphatase (ALP) and osteocalcin (OCN)] and two factors related to osteoclastogenesis [osteoprotegerin (OPG) and receptor activator of nuclear factor kappa  $\beta$  ligand (RANKL)] at the mRNA level.

**RESULTS:** The expression of OPG significantly decreased by 7 per cent compared with the control at 24 hours ( $P < 0.001$ ), while it increased by 133 per cent at 72 hours ( $P < 0.001$ ). RANKL remained unchanged at all three time points ( $P > 0.05$ ). ALP was upregulated by 73 per cent at 24 hours ( $P < 0.001$ ), but decreased by 14 and 41 per cent at 48 and 72 hours, respectively ( $P < 0.05$ ). OCN showed the same trend of expression change as that of OPG, which was down-regulated by 41 per cent at 24 hours but upregulated by 149 per cent at 72 hours ( $P < 0.001$ ).

**CONCLUSION:** VEGF promotes bone remodelling by direct effects on osteoblastic cells via regulating gene expressions of ALP, OCN and OPG.

### IN VITRO DIFFERENTIATION OF DENTAL PULP STEM CELLS INTO THE CHONDROGENIC ROUTE

**P Yugaraj**, P Zuo, A B M Rabie, R W K Wong, Department of Paediatric Dentistry and Paediatric Dentistry, Faculty of Dentistry, University of Hong Kong, SAR China

**AIM:** Cartilage cells have a limited ability to regenerate and self-repair. A large number of people are affected by cartilage deformity due to trauma, degenerative disease, congenital craniofacial defect, idiopathic condylar resorption, ageing, etc. The limited availability of donor sites for cartilage reconstruction makes it an ideal candidate for tissue engineering. Currently bone marrow aspirate is the most popular source of mesenchymal stem cells. Human dental pulp tissue has been found to be an alternative rich source of stem cells. The aim of this study was to examine the in vitro potential of dental pulp stem cells (DPSC) to differentiate into chondrogenic lineage when added to human transforming growth factor beta 3 (TGF- $\beta$ 3).

**MATERIALS AND METHOD:** A strain of stem cells were isolated from human dental pulp, expanded and further characterized. TGF- $\beta$ 3 was chosen to trigger its differentiation to cartilage. The TGF- $\beta$ 3 gene was subcloned as a recombinant pcDNA3.1/V5-His-TOPO-TGF- $\beta$ 3 vector. The vector was transfected into the DPSC. Three different negative controls were used; blank cells, empty vector and non-related green fluorescent protein (GFP) vector. After 48 hours, the mRNA level of collagen type II and aggrecan was measured by real-time polymerase chain reaction.

**RESULTS:** The relative transcription level of collagen type II mRNA in the TGF- $\beta$ 3 group was significantly higher than all three negative controls. The relative levels of collagen type II mRNA were 2.746 for blank cell, 154.3 for empty vector, 49.41 for non-related GFP vector and 233.4 for TGF- $\beta$ 3 vector.

**CONCLUSION:** TGF- $\beta$ 3 was successful in triggering DPSC differentiation into chondrogenic progenitor cells. It is concluded that DPSC have the capability to form cartilage cells.



## KEYNOTE LECTURES

### KL 01

Wednesday, June 16 - at 10:00

#### ETHICAL ISSUES IN 'ENHANCEMENT' MEDICINE

**Jože Trontelj**, Slovenia

Orthodontics is a medical field where most of the difficult ethical dilemmas of medicine seem to occur rather infrequently. Commonly, its goal is a dual effect: improvement of function and better aesthetic appearance. From the ethical point of view, this puts orthodontics close to the framework of enhancement medicine.

The scope of human enhancement with the aid of medicine, however, is much wider, and may, with advancement of technologies, present not only new challenges but also increasingly disturbing ethical doubts. Among these is the use of high-tech medicine for non-medical purposes, the gradual increase in standards of individual appearance and physical abilities, as well as discrimination against those who cannot afford the costly procedures. There is also the controversial possibility of genetic manipulation. It is important to recognize the ethical limitations of human enhancement in order to avoid far reaching undesired consequences.

### KL 02

Wednesday, June 16 - at 11:00

#### BIOTECHNOLOGY AND THE FUTURE IN ORTHODONTICS

**Fraser McDonald**, UK

There has been an exponential increase in the development of technology, especially with regard to the development of information technology. In particular both hard and software developments have meant the easy handling of large numbers of data points in such things as images. The most single advance that has occurred is the development of clinically applicable three-dimensional radiography. This can assist with our diagnosis but leaves us open to new clinical questions: Is a maxillary canine that is seen to be contacting adjacent teeth permanently damaging these teeth? Are we able to identify precise locations of ankylosis? Are large volume views able to provide more data which assist in treatment planning orthognathic surgery? All these questions will require objective evaluation in order to establish the clinical and therapeutic value of such imaging technology. It also raises questions about where our current imaging techniques will be placed in terms of supportive clinical information. Whilst techniques exist to superimpose radiographs, only limited data is available to identify what superimpositions can be carried out with confidence (Treil J, Faure J, Braga J, Casteigt J, Borianne P 2002 Three-dimensional imaging and cephalometry of cranio-facial asymmetry. L'Orthodontie française 73: 179-97). Alternative use of the ability to handle such large data sets include both study model image (storage and manipulation; derived from impressions negating the requirement to cast up the models) and facial images which in turn can then be integrated with the hard tissues. The data on soft tissue analysis whilst essentially of significant potential still retains the underlying inconsistencies of any imaging of soft tissue; that is the lack of reproducibility.

### KL 03

Wednesday, June 16 - at 14:30

#### BIOTECHNOLOGY ASPECTS IN ORTHODONTICS

**Andrea Wichelhaus**, Germany

Customised nickel-titanium (NiTi) arches allow customised treatment. Experimental research shows the effects of this programming and allows

application through thermal treatment and reshaping using forceps. This permits management of the biomechanical side effects of self-ligating systems.

Mechanics for treatment of spacing in the form of closing and controlled tipping are biomechanically favourable. Applications for treating adult patients are varied. An alloy arch is used as a segmental arch for this group of patients. Scientific research and case-studies show the biomechanical effects and associated clinical outcomes.

#### **KL 04**

*Wednesday, June 16 - at 16:30*

##### **THE USE OF BIOMECHANICAL MODELLING TO PREDICT SURGICAL CHANGE**

**Stephen Richmond, UK**

Predicting the outcome of orthognathic surgery is challenging in that the hard and soft tissue changes in the face require simulation and these changes are dependent on the surgeon being able to carry out the surgical plan within an overall mean surface error of less than 2 mm. The period of treatment is often in excess of two years during which the individual's face is continually maturing. The facial contours continue to change post-surgery as the facial swelling reduces. Therefore simulation of facial changes should ideally reflect a combination of maturation and surgical change.

However, in spite of efforts to provide a detailed predictive plan the outcome often falls short of reality. These errors may be minimized by: i) Full characterization of the facial features. Each face contains subtle differences that make appearance unique and it is important to identify these features. ii) Detailing the relationship of the various soft tissues to the underlying skeletal structures. iii) Carrying out a full assessment of horizontal and vertical asymmetry.

Various methods have been developed with the purpose of utilizing finite element modelling as a prediction tool for surgical procedures and facial expressions. Using the techniques described it is possible to individualize and mathematically model the soft and hard tissues to facilitate the prediction of hard and soft tissues as planned prior to surgery. The challenge is to refine these techniques for routine use and simulate speech and other facial actions.

#### **KL 05**

*Thursday, June 17 - at 8:30*

##### **SELF-LIGATING BRACKETS: CUI BONO?**

**Frank Weiland, Austria**

The first ligature free brackets were introduced in 1935. During the following decades numerous designs were patented but hardly ever reached production status. In recent years, however, self-ligating brackets have been the main focus of interest. All major orthodontic companies market their own designs. According to the promotional sales material, each company has the bracket system that may solve all clinical difficulties. This lecture will present a critical overview, based on clinical cases and scientific analyses, of the most frequently used self-ligating bracket systems on the market and point out the discrepancies between the promises of companies and scientific proof.

#### **KL 06**

*Thursday, June 17 - at 9:30*

##### **INTEGRATION OF PASSIVE SELF-LIGATION TREATMENT IN EVERYDAY CLINICAL PRACTICE**

**Alan Bagden, USA**

The points to be covered in this keynote address will include a basic understanding of the philosophy of self-ligation in general, and a specific knowledge of the development and philosophy of passive self-ligation (the Damon System). This will include the mastery of a specific organized protocol to clinical treatment, as exemplified through numerous successful clinical cases, as well as follow-up retention review. The ultimate goal of the presentation will be for each participant to achieve the ability to allow integration of passive self-ligation treatment into their own practices.

**KL 07***Thursday, June 17 - at 11:00***UPRIGHTING MISCONCEPTIONS ABOUT SELF-LIGATION****Friedrich K. Byloff**, Austria

The over enthusiastic, far too commercial and almost religious reflections that some have expressed about self-ligation has created a counter-movement in orthodontics. The impression orthodontists get when attending recent congress presentations and examining recent publications is that the pendulum is swinging 'back' toward manual ligation. It almost seems to be fashionable to say that there are no advantages to self-ligation. But is this really reflective of clinical reality? This presentation will attempt to create a better understanding of the self-ligation as a whole and what role it may play in the future of orthodontics. Various bracket designs will be contrasted and relevant research will be reviewed. Finally, recommended treatment approaches will be discussed under the light of well established and well researched orthodontic principles to dispel some of the myths being put forth.

**KL 08***Thursday, June 17 - at 14:30***LINGUAL ORTHODONTICS WITH COMPLETELY CUSTOMIZED APPLIANCES****Dirk Wiechmann**, Germany

Despite the uncontested aesthetic advantages, the lingual technique using conventional lingual appliances has not yet had its breakthrough. High bond failure rates, difficult rebonding, compromised patient comfort and difficult finishing are some of the main reasons for this. Modern lingual appliances are completely customized. Their unique features offer solutions for the aforementioned problems. This is why since its introduction in late 2004, Incognito has become the most popular lingual appliance worldwide to date.

**KL 09***Friday, June 18 - at 8:30***A RATIONAL TREATMENT APPROACH TO CLASS III MALOCCLUSION: DELIMITING THE CHALLENGE****Tiziano Baccetti**, Italy

Orthopaedic treatment of Class III malocclusions still presents with a prevalence rate of failure in about 20 to 30 per cent of patients. The Class III craniofacial growth pattern is also re-established after treatment, making relapse a probability.

This keynote lecture will focus on the clinical and biological factors that can improve the outcomes of Class III treatment by means of maxillary expansion and protraction. Data collected at the completion of active craniofacial growth are available for the evaluation of orthopaedic treatment of Class III disharmonies in the long term.

The prediction of the results of therapy in the individual patient, the definition of optimal treatment timing, and a better understanding of the relative roles of orthodontics and orthognathic surgery in the comprehensive treatment approach to the malocclusion, will greatly help in reducing the challenge of Class III treatment.

**KL 10***Friday, June 18 - at 9:30***CLASS III CAMOUFLAGE TREATMENT: WHAT ARE THE LIMITS?****Peter Wing Hong Ngan**, USA

Class III malocclusion can be intercepted in young patients with facemasks and chin caps. In adult patients, this malocclusion can be camouflaged with fixed appliances and orthodontic tooth movement. What are the limits of orthodontic and orthopedic camouflage treatment? What are the advantages with early compared with late treatment? With the advent of skeletal anchorage and the use of mini-implants, do they increase the possibility of camouflage orthodontic treatment? Class III cases camouflaged by orthodontic or orthopaedic treatment will be presented and the advantages and disadvantages of early and late treatment will be contrasted. New possibilities in camouflaging Class III malocclusions with mini-implants and miniplates will also be discussed.

**KL 11***Friday, June 18 - at 11:00***CONTEMPORARY TREATMENT OF SKELETAL CLASS III DEFORMITIES****Nataša Ihan Hren**, Slovenia

Dentofacial deformities such as mandibular prognathism and maxillary retrognathism result in a Class III occlusion that requires a combination of orthodontic and surgical treatment. The basic points of pre- and post-surgical orthodontic treatment are well-known and surgical procedures are standardized and without serious problems. However, the problem of post-surgical stability and treatment of developing Class III deformities still remains due to the many influences on the facial bones and teeth but, in particular, the facial soft tissue characteristics. Three-dimensional, ultrasound and speech analysis provides morphological and physiological knowledge that can improve the overall treatment of Class III orthognathic deformities.

**KL 12***Friday, June 18 - at 14:30***THE CHALLENGE OF APPLIANCE SELECTION...WITH SO LITTLE EVIDENCE****David L. Turpin**, USA

As patients become decision-makers there is more pressure on the professional health care provider to assist in making intelligent choices. High on the list should be knowledge of the evidence needed to justify a particular treatment approach. Based on the most complete in vivo studies, what do self-ligating brackets add to the correction of malocclusion? What is the role of Invisalign in your practice? Recent studies make it clear what can and cannot be accomplished with these very different appliances. There is little doubt that the continuum of a contemporary orthodontic practice will include an evidence-based component. Knowing how to search for and apply the evidence will become the driving force behind the 21st century practice.

**KL 13***Friday, June 18 - at 16:30***CLASS III TREATMENT - WHAT CAN WE DO?****Moriyon Uria**, Spain

Class III malocclusion has always been a challenge for the orthodontist. No matter how simple a Class III growing patient may seem there is always a certain amount of uncertainty in its treatment outcome. This is why many orthodontists are very reluctant to treat Class III cases before growth is finished or at least before the so called "pubertal growth spurt" has taken place, especially if there is genetic background, relying more on the orthognathic approach than on possibilities of a pure orthodontic solution. In the present lecture I will review some basic knowledge about Class III malocclusion and about that how different Class III clinical situations, ranging from early interceptive to adult treatment can be handled. From this information I will also try to draw some conclusions on what can be done and what should be avoided in order to render our patients a worthy service.



## ORAL PRESENTATIONS

### OP 01

Wednesday, June 16 - at 11:30

#### CRITICAL APICAL DIAMETER AND AUTOTRANSPLANTATION: A HISTOLOGICAL STUDY

**W Laureys**<sup>1</sup>, G De Pauw<sup>1</sup>, L Dermaut<sup>1</sup>, C Cuvelier<sup>2</sup>, Departments of  
<sup>1</sup>Orthodontics and <sup>2</sup>Anatomic Pathology, University of Gent, Belgium

**AIM:** Autotransplantation of immature teeth is an evidence based procedure and a valuable treatment option when making an orthodontic treatment plan. It has a high success rate when performed under ideal circumstances. This also means it has a restricted indication area and is age-related. The recovery of pulp tissue after transplantation of mature teeth is only 15 per cent. Expanding the indication area is possible by opening the apical foramen by apicoectomy of mature teeth. A critical diameter for the apical foramen is described in this literature. This critical diameter is measured on two-dimensional radiograph of immature teeth. Beneath this diameter of 1 mm the possibility of revascularisation after transplantation decreases. The aim of this study was to find histological evidence of this critical diameter.

**MATERIALS AND METHOD:** Fifteen single rooted mature teeth from three adult dogs were extracted, apicoectomized, the apical foramen digitally photographed and transplanted in infra-occlusion. The foramina varied from 0.24 to 1.09 mm. The observation period was 90 days post-transplantation. The histological slices were stained with haematoxylin and eosin.

**RESULTS:** Twelve of the teeth (80%) had new blood vessels filled with erythrocytes in at least one-third of the pulp chamber. Six of the teeth (40%) had new blood vessels up to the pulp horn. The apical foramina ranged from 0.31 to 1.09 mm at their widest.

**CONCLUSION:** Mature apicoectomized teeth with an apical foramen smaller than 1 mm can revascularise after autotransplantation with considerable success. If there exists a critical diameter for the apical foramen to allow revascularisation after transplantation, then it is smaller than 1 mm. Other factors such as occlusion (infra-occlusion after transplantation) may be more important than the size of the apical foramen when considering revascularisation of mature teeth after transplantation.

### OP 02

Wednesday, June 16 - at 11:40

#### CAN CONE BEAM COMPUTED TOMOGRAPHY PREDICT THE DIAMETER OF UNERUPTED TEETH?

**M Orellana**, E Nguyen, P Cubas, Department of Orofacial Sciences, School of Dentistry, University of California, San Francisco, USA

**AIM:** To test the accuracy and reproducibility of cone beam computed tomography (CBCT) in predicting the mesiodistal diameter (MDD) of unerupted teeth. A secondary aim was to determine the accuracy and reproducibility of three different viewing methods using two different CBCT softwares, InVivo dental 4.0 (Anatomage, San Jose, California) and CBWorks 3.0, (CyberMed Inc., Seoul, Korea). In measuring the MDD of teeth in models simulating unerupted teeth.

**MATERIALS AND METHOD:** CBCT data was collected on the CB MercuRay™ (Hitachi Medico Technology, Tokyo, Japan). Models of unerupted teeth (n = 25) created by embedding 25 samples into a polydimethylsiloxane polymer with similar density to tissues surrounding teeth, were scanned and measured by two investigators. Repeated MDD measurements of each sample were made using three CBCT viewing methods; InVivo sections view, InVivo volume render and CBWorks 3.0 volume render. These measurements were then compared with the MDD physically measured using a digital calliper before the teeth were embedded and scanned.



**RESULTS:** All three of the new methods had mean measurements that were statistically significantly less ( $P < 0.001$ ) than the physical method, adjusting for investigator and tooth effects. Specifically, InVivo sections view measurements were 0.273 (95% CI -0.36, -0.18) less than calliper measurements, InVivo volume render measurements were 0.49 less (95% CI -0.58, -0.40) than with the calliper, and CBWorks volume render measurements were 0.36 less (95% CI -0.24, -0.11) than calliper measurements.

**CONCLUSION:** Overall, there were high correlations amongst the three viewing methods indicating that CBCT can be used to measure the MDD of unerupted teeth. The InVivo section view method had the largest correlation with the calliper method.

### OP 03

Wednesday, June 16 - at 11:50

#### THREE-DIMENSIONAL AIRWAY VOLUME CHANGES IN ORTHOGNATHIC SURGERY PATIENTS

**N Stefanović<sup>1</sup>**, H El<sup>2</sup>, D Chenin<sup>3</sup>, D Nikolin<sup>4</sup>, J M Palomo<sup>2</sup>, <sup>1</sup>University of Belgrade, Serbia, <sup>2</sup>Case Western Reserve University, Cleveland, Ohio and <sup>3</sup>Anatomage Inc., San Jose, California, USA, <sup>4</sup>Private Practice, Stara Pazova, Serbia

**AIM:** Treating patients with dentofacial deformities commonly requires combined orthodontic-surgical treatment. Skeletal tissue movements within orthognathic surgery influence the shape and dimensions of the surrounding soft tissues and upper airways. The purpose of this study was to evaluate oropharyngeal (OP) airway volume changes in patients treated with either mandibular setback, or mandibular advancement surgery.

**MATERIALS AND METHOD:** Cone beam computed tomographic (CBCT) images of seven patients who underwent mandibular advancement (group A) and five patients who underwent mandibular setback surgery (group B). CBCT scans were taken before and after surgery using the CB MercuRay scanner (Hitachi Medical, Tokyo, Japan), with an average of one month between the two scans. DICOM images were analyzed using InVivo Dental Software (Anatomage Inc., San Jose, California, USA). For OP airway calculation, the palatal plane was used as the superior border, and the most antero-inferior point of the second cervical vertebrae as the inferior border. Volumes were calculated by a single operator, using automatic segmentation. All measurements were repeated for reliability tests. The Statistical Package for Social Sciences, version 17.0 (SPSS Inc., Chicago, Illinois, USA) was used for statistical analysis. Differences between the pre- and post-surgical groups were tested using a paired samples t-test, considering a significance of  $P < 0.05$ .

**RESULTS:** Intraoperator results showed good reproducibility. Data analysis showed an increase in OP volume in group A, and a decrease in group B. Comparative statistical analysis showed the change in group A was statistically significant ( $P < 0.05$ ), while those in group B were not ( $P > 0.05$ ).

**CONCLUSION:** Changes in OP airway volume occur after mandibular surgery. However, further investigation is required to determine the significance and amount of airway volume change in relation to mandibular movement. The change in airway volume may be an important factor to consider when planning mandibular antero-posterior skeletal movement.

### OP 04

Wednesday, June 16 - at 12:00

#### IMPORTANCE OF AGE IN ORTHODONTICS; A STUDY OF CRANIOFACIAL MUSCLE MATURATION

**K Proczek**, M Lewis, R Shah, N Hunt, UCL Eastman Dental Institute, London, England

**AIM:** Several treatment modalities currently employed to correct malocclusion and facial deformity incite an adaptive response in the muscles of the face and jaws. An understanding of the mechanisms involved in the adaptive response of the muscles is therefore important. Included in this is an appreciation of whether muscle response varies with age. Clinically, this could influence the choice of specific treatment approaches in patients of differing age. The aim of this study was to investigate the effect of age on the response of rabbit masseter muscle derived cells within a three-dimensional environment (3D).

**MATERIALS AND METHOD:** Myogenic muscle precursor cells (MPCs) were obtained from masseter muscle explants of four 10 month old, female, New

Zealand white rabbits. A known number of cells ( $4 \times 10^6$ ) were placed in a 3D collagen gel model anchored by metal floatation bars suspended in a chamber slide. As the MPCs proliferated and matured into myotubes (primitive muscle fibres), the development of the muscle contractile apparatus caused increased tension within the model, resulting in bowing of the gel. The degree of bowing of the respective gels was measured daily over a 14 day period and compared using a Mann Whitney test. MPCs derived from the same explants were also observed in a two-dimensional milieu over a similar time period to check the corresponding level of muscle differentiation.

**RESULTS:** The mean bowing of the gel based on young MPCs was statistically significantly higher ( $P < 0.001$ ) than that of mature MPCs at all times.

**CONCLUSION:** There are statistically significant differences in the behaviour of young and mature masseter muscle-derived cells, which may have clinical implications.

## OP 05

Wednesday, June 16 - at 12:10

### OPEN BITE TREATMENT WITH ZYGOMATIC ANCHORAGE

**M Sarioğlu**, H Türkkarman, Suleyman Demirel University, Isparta, Turkey

**AIM:** To evaluate the dentoalveolar and skeletal effects of open bite treatment with zygomatic anchorage.

**SUBJECTS AND METHOD:** Two groups were matched, a treatment and a control. Each group comprised 20 patients. The mean chronological age for the treatment group was  $16.68 \pm 2.80$  years and for the control group  $16.63 \pm 2.83$  years. The mean treatment period was  $1.00 \pm 0.31$  years. The subjects in control group were observed for  $0.95 \pm 0.14$  years. The study was carried out on 80 lateral cephalometric radiographs and dental casts taken before and at the end of the treatment or observation. Titanium integrated miniplates were placed on the lower contours of each zygomatic process using a minor surgical procedure. Nickel titanium closed coil springs were applied from the miniplates to the posterior teeth for intrusion force. A rigid hyrax screw was used for transverse control of the posterior teeth.

**RESULTS:** Anterior open bite closure was achieved in all patients. The amount of intrusion was  $3.59 \pm 1.34$  mm at the first maxillary molars ( $P < 0.001$ ). The overbite increased  $4.82 \pm 1.53$  mm ( $P < 0.001$ ). A closure of  $2.25 \pm 1.91$  degrees was determined for the mandibular plane angle (Go-Gn-SN) ( $P < 0.001$ ). Anterior face heights decreased as the mandible closed.

**CONCLUSION:** Zygomatic anchorage can be used effectively for open bite correction through posterior dentoalveolar intrusion.

## OP 06

Wednesday, June 16 - at 15:00

### CATEGORIZATION OF FACIAL FORM

**A Toma**<sup>1</sup>, A Zhurov<sup>1</sup>, R Playle<sup>1</sup>, D Marshall<sup>2</sup>, S Richmond<sup>1</sup>, <sup>1</sup>Dental School and <sup>2</sup>School of Computer Science, Cardiff University, Wales

**AIM:** To characterize normal variation of facial morphology in a young population based on three-dimensional (3D) analysis of facial soft tissue landmarks.

**SUBJECTS AND METHOD:** Three hundred and fifty British-Caucasian children (166 females, 184 males) aged 15½ years were recruited from the Avon Longitudinal Study of Parents and Children. 3D facial images were obtained for each individual using two high-resolution Konica/Minolta laser scanners. Twenty-one reproducible facial landmarks were identified on each image resulting in 63 X, Y, and Z coordinates. Generalized Procrustes analysis was used to provide the best fit of the 21 facial landmarks, considering both size and shape variations, equally important for 3D analysis of facial morphology. Principal component analysis was used to highlight important components that describe normal variation of facial morphology.

**RESULTS:** Thirteen principal components were identified that describe facial variation within the population, with the first three components explaining 46.6 per cent of the total variance in facial form.

**CONCLUSION:** The findings challenge traditional categorisations of facial morphology. Facial variation can be considered as a continuum. This method of facial assessment may be useful to identify and classify faces and facial changes that occur as a result of growth, and inform clinicians of appropriate healthcare interventions for specific facial types.



### PALATAL IMPLANT SUPPORTED VERSUS CONVENTIONAL DENTAL ANCHORAGE IN ADOLESCENTS. A RANDOMIZED CONTROLLED TRIAL

**G Borsos**<sup>1</sup>, T Gedrange<sup>2</sup>, A Végh<sup>1</sup>, <sup>1</sup>Department of Orofacial Orthopaedics and Orthodontics, Heim Pal Children's Hospital, Budapest, Hungary and <sup>2</sup>Department of Orthodontics, Preventive and Pediatric Dentistry, University of Greifswald, Germany

**AIM:** To clinically compare osseointegrated palatal implant (PI) supported anchorage with conventional dental anchorage (DA) for two-phase retraction, in extraction cases requiring 'maximum anchorage', in growing patients after the post-pubertal growth spurt.

**SUBJECTS AND METHOD:** Thirty patients with homogeneous facial skeletal characteristics (mean age  $14.22 \pm 1.37$  years) randomly allocated to two groups. In the PI group ( $n = 15$ , mean age  $14.15 \pm 1.2$  years) Orthosystem® implants were placed in the palate for absolute anchorage and, in each case, a  $1.2 \times 1.2$  mm thick rigid transpalatal arch (TPA) was fixed to the implant and to the molar bands by laser welding. In the DA group ( $n = 15$ , mean age  $14.3 \pm 1.57$  years), maximal anchorage was provided by a TPA and a  $0.017 \times 0.025$  inch stainless steel (SS) utility arch. A superelastic closed-coil spring was used for canine retraction. Sequential activation of the 'teardrop' closing loop of the SS contraction arch was used for incisor en masse retraction. The main outcome measures were: duration of the orthodontic treatment phases and cephalometric analysis of maxillary first molar movement.

**RESULTS:** An insignificant difference ( $P = 0.47$ ) was observed between the groups regarding the duration of extraction gap closure. In the PI group the duration of anterior retraction ( $P < 0.001$ ) and total treatment time was shorter ( $P < 0.05$ ). No significant difference in mesial molar movement was found in either treatment phase even if the net molar anchorage loss was  $4.35 \pm 1.69$  mm in the DA group while  $3.7 \pm 1.94$  mm in the PI group ( $P = 0.35$ ).

**CONCLUSION:** A shorter orthodontic treatment period was achieved by fastening the anchorage teeth to a PI compared with DA. The use of PI supported anchorage was without complication, representing a safe alternative in growing adolescents requiring maximum anchorage.

### EFFECTS OF VASCULAR ENDOTHELIAL GROWTH FACTOR AND DANSHEN ON OSTEOBLASTIC CELL ACTIVITIES

**Y Yang**, Y Tan, A Chin, R W K Wong, A B M Rabie, Orthodontics, University of Hong Kong, SAR China

**AIM:** The search for an ideal bone grafting material remains a challenge in modern surgery. Sufficient bone support is essential for effective orthodontic tooth movement. Osteogenesis and angiogenesis are closely correlated. This study aimed to investigate the effects of vascular endothelial growth factor (VEGF) and the medicinal herb, Danshen, on osteoblastic cell activities.

**MATERIALS AND METHOD:** An osteoblastic cell line, MC3T3-E1, was cultured in vitro with or without VEGF and Danshen. The cells in both control and test groups were collected at different culture time points of 24, 48 and 72 hours. Real-time polymerase chain reaction was carried out to quantify the expression of VEGF, alkaline phosphatase (ALP), osteopontin (OPN), osteocalcin (OCN), osteoprotegerin (OPG) and the receptor activator of nuclear factor kappa B ligand (RANKL) at the mRNA level.

**RESULTS:** The expression of VEGF was down-regulated by extrinsic VEGF ( $P < 0.05$ ), while upregulated by extrinsic Danshen ( $P < 0.05$ ). The expression of ALP was upregulated by both VEGF and Danshen ( $P < 0.05$ ), but Danshen showed a more continuous effect. Both VEGF and Danshen upregulated the expression of OPN and OCN ( $P < 0.05$ ), but the effect of Danshen appeared earlier than that of VEGF. Both VEGF and Danshen regulated the expression of OPG, while only Danshen regulated the expression of RANKL.

**CONCLUSION:** Danshen enhances expression of VEGF in osteoblastic cells, but VEGF shows negative feedback of auto-regulation. Both VEGF and Danshen promote bone remodelling by direct effects on osteoblastic cells via regulating gene expression of ALP, OPN, OCN and OPG.

Supported by Seed Funding Programme for Basic Research HKU 200803159006.

**FABRICATION AND BIOCOMPATIBILITY OF COMPOSITE SCAFFOLDS FOR BONE TISSUE ENGINEERING**

V D'Antò<sup>1</sup>, M G Raucci<sup>2</sup>, R Valletta<sup>1</sup>, R Martina<sup>1</sup>, A Luigi<sup>2</sup>, <sup>1</sup>University of Naples and <sup>2</sup>Institute for Composite and Biometical Materials – CNR, Italy

**AIM:** Tissue engineering is a novel field of research that aims to repair damaged tissues as well as to create replacement organs. In orthodontics, tissue-engineered temporomandibular joints, periodontium, and cranial sutures may have a dramatic impact on treatment possibilities. Tissue-engineered bone can also be useful in patients who need bone grafts or cleft palate repair. The aims of this project were to fabricate a novel composite scaffold for bone tissue engineering and to study the proliferation and differentiation of human mesenchymal stem cells (hMSCs) in this matrix.

**MATERIALS AND METHOD:** Hydroxyapatite/polycaprolactone (HA/PCL) composite scaffolds were prepared by the wet-chemical method at room temperature. Physico-chemical properties of the composite materials were preliminary characterized by X-ray diffraction analysis, Fourier transform infrared spectroscopy and X-ray photoelectron spectroscopy analysis. Moreover, the scaffold morphology was investigated by scanning electron microscopy and energy dispersive X-ray spectroscopy to validate the process used for the synthesis. Finally, the response of two different lines of hMSCs in terms of cell proliferation and differentiation into osteoblastic phenotype was evaluated using Alamar blue assay, scanning electron microscopy, and alkaline phosphatase activity.

**RESULTS:** Microstructural analysis indicated that the HA particles were distributed homogeneously within the PCL matrix. Biological results revealed that the HA/PCL composite scaffolds were well recognized from MSCs in vitro, supporting cell proliferation and differentiation of bone marrow derived MSC.

**CONCLUSION:** Tissue engineering by means of composite HA/PCL scaffolds represents a new therapeutic strategy to repair craniofacial bone defects and may have a strong impact in the treatment of complex interdisciplinary orthodontic patients.

This study was supported from the Ministero dell'Università e della Ricerca by funds of Rete Nazionale di Ricerca TISSUENET n. RBPR05RS and PRIN 20072L2KRX.

**BONE MARROW-DERIVED STEM CELLS IN PALATAL WOUND HEALING**

J Verstappen<sup>1</sup>, C Katsaros<sup>2</sup>, R Torensma<sup>3</sup>, H Von den Hoff<sup>1</sup>, <sup>1</sup>Department of Orthodontics and Oral Biology, Radboud University Nijmegen Medical Centre, Nijmegen, Netherlands, <sup>2</sup>Department of Orthodontics and Dentofacial Orthopedics, University of Bern, Switzerland, <sup>3</sup>Department of Tumor Immunology, Nijmegen Centre for Molecular Life Sciences, Netherlands

**AIM:** Myofibroblasts are responsible for contraction and scarring after cleft palate repair. This may lead to growth disturbances in the upper jaw. The hypothesis tested was that circulating stem cells originating from the bone marrow are recruited to palatal wounds and differentiate into myofibroblasts. If confirmed, this might offer new opportunities for anti-scarring therapy in cleft palate surgery, by preventing the recruitment of specific myofibroblast precursors. The aim of this study was to investigate the contribution of bone marrow-derived stem cells (BMDSCs) to the myofibroblast population in experimental wounds in the palate.

**MATERIALS AND METHOD:** Bone marrow from five green fluorescent protein (GFP)-transgenic Sprague-Dawley rats was transplanted into 10 lethally irradiated wild-type rats. Five weeks later, 89 ± 8.9 per cent of the mononuclear cells in the blood of the recipient rats were GFP-positive. Seven weeks after the bone marrow transplantation, 4 mm experimental wounds were made in the palatal mucoperiosteum. Wound and adjacent control tissue was harvested with a 5 mm biopsy punch. The tissue samples were fixed in 4 per cent paraformaldehyde for 24 hours, and embedded in paraffin. The contribution of GFP expressing BMDSCs to the myofibroblast population in both the wound and control areas was investigated by immunostaining. In addition, the contribution of BMDSCs to other cell

populations was estimated with specific markers.

**RESULTS:** At two weeks post-wounding,  $8.4 \pm 7.4$  per cent of all cells in the wound area were GFP-positive and  $5.0 \pm 4.0$  per cent of the myofibroblasts, which was significantly higher than in the control area. Similar percentages were found for activated fibroblasts and endothelial cells, but for myeloid cells it was considerably higher ( $22 \pm 9\%$ ).

**CONCLUSION:** BMDSCs contribute to palatal wound healing, but are not the main source of myofibroblasts. In small wounds, local precursor cells are probably sufficient to replenish the defect. This study was supported by an EOS Research Grant.

## OP 11

Wednesday, June 16 - at 15:50

### A NEW METHOD FOR AUTOMATIC LOCALIZATION OF CEPHALOMETRIC LANDMARKS ON DIGITAL IMAGES

**R Leonardi**, D Giordano, F Maiorana, M Greco, University of Catania, Italy

**AIM:** Efforts have been made to completely automate cephalometric analysis by automatic landmarking. However, the accuracy obtained was poorer than manual identification. Analogue-to-digital conversion of radiographs has been reported to be the main problem. Therefore the aim of this investigation was to evaluate the accuracy of the Cellular Neural Networks approach for automatic location of cephalometric landmarks on soft copies of direct digital cephalometric radiographs

**MATERIALS AND METHOD:** Forty-one, direct digital lateral cephalometric radiographs were obtained with a Siemens Orthophos DS cephalostat. Ten landmarks (N, point A, Ba, Po, Pt, point B, Pg, PM, UIE, LIE) were the object of automatic landmark identification. The mean errors and standard deviations from the best estimate of cephalometric points were calculated for each landmark. Differences in the mean errors of automatic and manual landmarking were compared with one-way analysis of variance.

**RESULTS:** Statistical analysis indicated that the differences were very small, and they were found at most within 0.59 mm. While a few of these differences were statistically significant, they were so small as to be, in most instances, clinically meaningless.

**CONCLUSION:** The use of radiographic files with respect to scanned radiographs improves landmark accuracy of automatic detection. Investigations on soft copies of digital cephalograms, to search for more landmarks in order to enable complete automatic cephalometric analysis are necessary.

## OP 12

Wednesday, June 16 - at 17:00

### AIRWAY OBSTRUCTION: NEW DATA ON MORPHOLOGICAL CHANGES INDICATE EARLIER INTERVENTION

**A Macari**<sup>1</sup>, J Ghafari<sup>1,2,3</sup>, <sup>1</sup>Division of Orthodontics and Dentofacial Orthopedics, Department of Otolaryngology Head and Neck Surgery, American University of Beirut Medical Center and <sup>2</sup>Lebanese University School of Dentistry, Lebanon and <sup>3</sup>Department of Orthodontics, Dental College, New York, USA

**AIM:** Airway blockage leads to dentofacial changes culminating in the long-face syndrome or 'adenoid facies', which is difficult to treat to the normal skeletal morphology. The aim of this study was to evaluate dentofacial changes following nasal obstruction and determine indicators for optimal timing of adenoidotomy that would avoid or minimize dentofacial alterations.

**MATERIALS AND METHOD:** Cephalographs of 200 Caucasian children (ages: 1.71-12.62 years) referred for evaluation of airway obstruction. Conventional angular and linear cephalometric measurements were used to determine relationships among cranial base, maxilla, and mandible. The distances from the soft palate, shortest to the adenoid (SAD), and to the most convex point on the adenoid (CAD) measured airway clearance, also graded (1-3) independently through a common rating system of the adenoid blocking various percentages of the airway. The occlusal relationships were noted. Statistical analyses included t-tests and analyses of variance for differences in continuous measurements between groups (by age or comparisons with available Caucasian norms); the Pearson product moment correlation coefficient for associations between various parameters and linear and multiple regressions for associations and prediction models.

**RESULTS:** Dysmorphology compensatory to nasal obstruction as early as age 1.7 years, the maxilla often affected first with a postero-inferior tilt of the palatal plane to the horizontal (PP/H  $-7.7 \pm 3.4^\circ$ ), greatest (For all subjects a negative overjet, mesiocclusion, immature swallowing pattern and incorrect tongue posture on the mouth floor were diagnosed at baseline. A Class III facial appearance was also observed ( $8.90 \pm 3.6^\circ$ ) between 4-5 years of age. Other changes occurred separately or together: mandibular antegonial notching, steeper mandibular plane, anterior symphyseal reduction, increased lower face height. Malocclusions included: increased overjet with normal or distocclusion, open bite, anterior and posterior crossbite. SAD and airway grading were highly correlated ( $r = 0.92$ ). SAD in combination with maxillary length (ANS-PNS) and inclination (PP/H) accounted for nearly 65 and 60 per cent of the variation in mandibular size, respectively.

**CONCLUSION:** The upper airway is more likely to be blocked by adenoid hypertrophy when SAD measures 0-2 mm. Adenoid removal is recommended early (age 2-3 years) to arrest or reverse long-face development.

## OP 13

Wednesday, June 16 - at 17:10

### CERVICAL VERTEBRAE ANOMALIES ASSESSED ON LATERAL CEPHALOGRAMS AND CONE BEAM COMPUTED TOMOGRAPHS

**D Bebnowski**<sup>1</sup>, M Hänggi<sup>1</sup>, G Markic<sup>1</sup>, M Roos<sup>2</sup>, T Peltomäki<sup>3</sup>, <sup>1</sup>Clinic for Orthodontics and Pediatric Dentistry and <sup>2</sup>Institute of Social and Preventive Medicine, Biostatistics Unit, University of Zurich, Switzerland and <sup>3</sup>Dental and Oral Diseases Outpatient Clinic Tampere University Hospital, Finland

**AIMS:** A high prevalence of cervical spine anomalies has been associated with various malocclusions by Sonnesen et al. These findings have recently been challenged due to difficulties in reliably determining vertebral spine anomalies on a single lateral cephalogram (Koletsis, 2009). The aims of this research were to study the prevalence of cervical spine anomalies on lateral cephalograms in Class II subjects and to compare the findings with those obtained from cone beam computed tomographs (CBCT).

**MATERIALS AND METHOD:** Standardized cephalograms of 238 patients (129 males, 109 females) with at least a three-quarter Class II molar relationship and overjet  $\geq 4$  mm were analyzed. Vertebral anomalies were identified by visual assessment and structural tracing. Both a cephalogram and CBCT were available for an additional 21 subjects. Cephalometric values were correlated with vertebral morphology, and logistic regressions were calculated. Intraobserver agreement was evaluated using the kappa index.

**RESULTS:** Inspection of the lateral cephalograms excluded an anomaly of the cervical vertebrae in 90.3 per cent of the subjects, while 9.7 per cent showed potential fusion. No posterior arch deficiency or block fusion was found. The measurements showed good repeatability (kappa 80). No correlations were found between the cephalometric values and potential vertebral anomalies. In the 21 patients with both a CBCT and a lateral cephalogram, visual assessment of the lateral cephalogram yielded a possible fusion in nine cases, while none could be confirmed with the CBCT.

**CONCLUSION:** A low number of potentially fused cervical vertebrae could be detected using lateral cephalograms in Class II patients. The possible fusions did not correlate with any cephalometric values nor could they be confirmed in those patients with a CBCT, the gold standard for assessing cervical vertebrae anomalies. Visual examination of a single cephalogram may result in a false-positive finding and does not allow reliable diagnosis of cervical vertebral fusion.


## OP 14

Wednesday, June 16 - at 17:20

### VITAMINS AND CLEFTING: PUTATIVE BIOLOGICAL MECHANISMS

**M Ackermans**, H Von den Hoff, Department of Orthodontics and Oral Biology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

**AIM:** In the aetiology of cleft lip and palate (CLP), both genetic and environmental factors are involved. While the genetic aetiology of CLP is starting to become unravelled, the environmental factors are still not



completely understood. Understanding their action could be of value for the prevention of CLP. In this context, dietary supplements such as vitamins, are of particular interest. The aim of this research was to identify the putative biological mechanisms underlying the association between maternal vitamin use and CLP in the offspring. The main focus was on folate and retinoic acid (RA).

**MATERIALS AND METHOD:** A systematic review was performed using PubMed (1953-December 2009) to review the effects of folate and retinoic acid on cellular processes involved in palatogenesis and CLP.

**RESULTS:** Retinoic acid (RA) and folate interfere with several signalling pathways that are also involved in the regulation of palatogenesis. RA inhibits EGF, bone morphogenetic protein, and platelet derived growth factor-C signalling, which regulate mesenchymal proliferation. The transforming growth factor (TGF)- $\beta$ 3 and Wnt pathway, which are crucial for epithelial fusion, are also affected. Furthermore, RA directly inhibits the synthesis of extracellular matrix components, and the formation of filopodia and desmosomes. It further inhibits disintegration of the epithelial seam by reducing matrix metalloproteinase activity and apoptosis. Low folate status disturbs homocysteine metabolism and affects cell proliferation, adhesion, and apoptosis. This might be mediated by inhibition of TGF- $\beta$ , IGF, Sonic hedgehog, and EGF signalling.

**CONCLUSION:** RA overexposure and folate underexposure interfere with important signalling pathways in palatogenesis. The inhibition of EGF, TGF- $\beta$ , and Wnt pathways might be crucial for the development of CLP. Better understanding of vitamin action on palatogenesis is vital for CLP prevention.

## OP 15

Wednesday, June 16 - at 17:30

### SIMULATION OF ORTHOGNATHIC SURGERY USING CONE BEAM COMPUTER TOMOGRAPHY AND DIGITAL MODELS

**J Seeberg**, B Melsen, M Dalstra, T K Pedersen, J Buhl, Department of Orthodontics, School of Dentistry, Aarhus University, Denmark

**AIM:** To develop a routine by which orthognathic surgery can be simulated and the necessary wafers produced by integrating cone beam computer tomography (CBCT) with digital models.

**MATERIALS AND METHOD:** CBCT images and O3DM<sup>®</sup> virtual models (Ortolab, Czestochowa, Poland) in STL format of patients being prepared by means of orthodontics for orthognathic surgery visualized in Mimics (Materialise, Leuven, Belgium). Alginate impressions served as the basis for generation of the virtual models and the data were converted to STL format and merged with the three-dimensional (3D) CBCT image dataset of the craniofacial skeleton. This was done by superimposing tooth structures from the CBCT on those from the virtual model using a minimum of eight points on both models. The planned surgical intervention (simulating a bilateral sagittal split osteotomy and Le Fort I surgery) was performed using the simulation tools of Mimics in such a way that the desired dental relationship was obtained. Minor occlusal adjustments were corrected using 3D software. Space was generated for the wafer by rotating the mandible around a centre corresponding to the condyle as in an individual articulator. The wafer could then be produced with computer controlled milling or a 3D printing process.

**RESULTS:** The combination of CBCT and the virtual models resolved the scattering artefacts of teeth with fillings or brackets in the CBCT images, visualised necessary tooth movement in the upper and lower arch, and allowed for an iterative process of 3D simulation of the surgery mimicking cuttings in the anatomically correct positions.

**CONCLUSION:** A method allowing for optimal utilization of the information from 3D images of both the face and model was developed. The high resolution of the 3D models allows for satisfactory precision in the production of the surgical wafer.

## OP 16

Wednesday, June 16 - at 17:40

### ROOT RESORPTION FOLLOWING 15 DEGREES TIP, 15 DEGREES TORQUE AND A ROTATING FORCE OF 225 GRAMS – A MICRO-COMPUTED TOMOGRAPHIC STUDY

S V Neitzert<sup>1</sup>, T Turk<sup>2</sup>, C Colak<sup>2</sup>, **S Elekdag-Turk<sup>2</sup>**, M A Darendeliler<sup>1</sup>, <sup>1</sup>Discipline of Orthodontics, Faculty of Dentistry, University of Sydney, Australia and <sup>2</sup>Department of Orthodontics, Faculty of Dentistry, University of Ondokuz Mayıs, Samsun, Turkey

**AIM:** To investigate and compare the amount of root resorption following the application of 15 degrees tipping, 15 degrees torque and 225 g rotating orthodontic forces using micro-computed tomography ( $\mu$ CT).

**SUBJECTS AND METHOD:** Thirty patients who required bilateral extractions of maxillary first premolars divided into three equal groups. In group TQ, the maxillary first premolar on one side of each patient had 15 degrees of distal root tipping force applied while the contralateral side was subjected to 15 degrees buccal root torque. In group TR, the maxillary first premolar of each patient had 15 degrees of distal root tipping force applied to one side and 225 g of rotating force to the other side. In group QR, the maxillary first premolar tooth on one side of each patient was subjected to 15 degrees of buccal root torque and the contralateral side to 225 g of rotating force. After four weeks the premolar teeth were extracted. The volume of root resorption craters was analyzed using  $\mu$ CT and measured utilizing specially designed software. Univariate analysis of variance (ANOVA) was performed.

**RESULTS:** There was a statistically significant difference in the extent of root resorption following tipping, torquing and rotational movement ( $P = 0.002$ ). In a pairwise comparison there was a significant difference between rotation and torque ( $P = 0.001$ ), torque and tip ( $P = 0.044$ ) but not between tip and rotation ( $P = 0.082$ ). The location of root resorption differed following the different tooth movements. Age and gender had no significant effect on the amount of root resorption.

**CONCLUSION:** (1) 225 g of rotating force causes more root resorption than 15 degrees torquing force (2) A 15 degree tipping force causes greater amounts of root resorption than 15 degrees torquing force (3) 225 g of rotating force causes similar amounts of root resorption as 15 degrees of tipping force (4) Overall, root resorption is found in areas of periodontal ligament compression while minimal resorption is found in areas of tension

## OP 17

Wednesday, June 16 - at 17:50

### THE INFLUENCE OF THE CENTRAL AND PERIPHERAL NERVOUS SYSTEMS ON HUMAN CRANIOFACIAL DEVELOPMENT

I Kjaer, Department of Orthodontics, School of Dentistry, Copenhagen University, Denmark


**AIM:** Close developmental interrelationship between the central (CNS) and peripheral (PNS) nervous system has been demonstrated in human embryological and foetal pathological material (Kjaer, 1998). Several research projects have since been performed, elucidating this interrelationship in the postnatal craniofacial area. This presentation is an overview of these results and demonstrates the interrelationship between the human CNS/neurocranium and CNS/PNS/dentition.

**SUBJECTS AND METHOD:** CNS/neurocranium: Profile radiographs of children/adults with different CNS disorders (lumbosacral myelomeningocele, Cri-du-Chat syndrome, Down syndrome, Fragile X syndrome, solitary median maxillary central incisor). In several cases the CNS diagnoses were based on magnetic resonance analyses and cephalometry and visual evaluation of sella turcica morphology. CNS/PNS/dentition: Dental pantomographs of mentally normal children. Histological sections of the permanent teeth. Evaluation of tooth formation and eruption. Histological localization of innervation.

**RESULTS:** CNS/neurocranium: Deviations in the cranial dimensions and morphology of sella turcica showed that disorders in the hemispheres were associated with deviations in the anterior wall of sella turcica and a short nasion-sella distance, while a deviation in the posterior wall was associated with disorders in the cerebellum. CNS/PNS/dentition: Agenesis and arrested eruption caused by inborn or acquired innervations were documented. Peripheral nerves were localized in the periodontal membrane. Furthermore, immunohistochemical analysis of osteoblasts indicated that compensatory and dysplastic craniofacial growth is under the control of peripheral nerves (Kjaer and Nolting, 2008).

**CONCLUSION:** There is a close interrelationship between CNS/PNS and the craniofacial area. This should be considered in future studies and, if available.

Kjaer I 1998 Neuro-osteology. Critical Reviews in Reviews in Oral Biology and Medicine 9: 224-244 Kjaer I, Nolting D 2008 Immunohistochemical PGP 9.5 positivity in human osteoblasts may indicate that compensatory and dysplastic craniofacial growth are under control by peripheral nerves. Orthodontics and Craniofacial Research 66:342-200



**IN VIVO COMPARISON OF TWO CONE BEAM COMPUTED TOMOGRAPHY SYSTEMS VERSUS PANORAMIC IMAGING**

**A Alqerban**<sup>1</sup>, R Jacobs<sup>2</sup>, S Fieuw<sup>3</sup>, G Willems<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Oral Imaging Center, Katholieke Universiteit Leuven and <sup>3</sup>Interuniversity Institute for Biostatistics and Statistical Bioinformatics, Leuven, Belgium

**AIM:** Mismanagement and inaccurate diagnosis of impacted maxillary canines may cause complications during development and eruption. The most frequent adverse effect of canine impaction is resorption of the adjacent incisor. The aim of this study was to compare the radiographic diagnostic accuracy for the localization of maxillary impacted canines and the detection of root resorption in adjacent incisors between conventional radiographic procedures using a two-dimensional (2D) panoramic radiograph with that of two three-dimensional (3D) cone beam computed tomography (CBCT) systems.

**MATERIALS AND METHOD:** The clinical records of 60 consecutive patients who had impacted or ectopically erupting maxillary canines were identified from patients seeking orthodontic treatment. For each case, two sets of radiographic information were obtained. The sample was divided into two groups group A (n = 30) included those for whom a panoramic and CBCT obtained with a 3D Accuitomo Tomograph was available, and group B (n = 30) who had a panoramic and CBCT obtained with a Scanora 3D CBCT. Panoramic and CBCT images were produced and subsequently analyzed by 11 examiners.

**RESULTS:** There was a highly significant difference observed between the 2D and 3D images for the width of the canine crown, canine location, and canine angulation to the occlusal plane. Moreover, there was a highly significant difference between the panoramic and Scanora CBCT images in canine angulation to the midline ( $P < 0.001$ ). The presence or absence of root resorption and the severity of the lateral incisor was also significantly different between 2D and 3D in both groups. Detection of central incisor root resorption was significantly different between the Accuitomo and panoramic images.

**CONCLUSION:** The use of CBCT imaging for the assessment of impacted canines has a potential diagnostic effect and may reduce treatment time, complexity, complications, and cost. CBCT is a reliable method for detecting root resorption of adjacent teeth and more accurate for the different diagnostic tasks than panoramic radiography.

**MEASUREMENTS OF DENTAL DIMENSIONS ON DIGITAL MODELS ARE SUPERIOR TO CONE BEAM COMPUTED TOMOGRAPHIC MEASUREMENTS**

**H Breuning**<sup>1</sup>, F Rangel<sup>1</sup>, O van Vlijmen<sup>1</sup>, T Maal<sup>2</sup>, A M Kuijpers-Jagtman<sup>1</sup>, Department of <sup>1</sup>Orthodontics and Oral Biology and <sup>2</sup>Oral and Maxillofacial Surgery, Radboud University Nijmegen Medical Centre, Netherlands

**AIMS:** Recent technological breakthroughs, such as digital models and cone beam computed tomographs (CBCT) enable digital measurement of the dentition. The aim of this study was to determine the accuracy of measurements on digital models of scanned impressions, volume rendered CBCT and teeth separated out of CBCT images. A further aim was to study the time needed for the digital measuring.

**MATERIALS AND METHOD:** Digital models of 10 patients with a complete permanent dentition were selected. Three digital datasets were available: a Digimodel™ (A), a CBCT image (B) and a segmented model of the dentition of a CBCT image, AnaTomodel™ (C). A commercially available program (Digimodel™) was used for measurements on the digital model. Another commercially available program (Maxillim™) was used for measurements on images B and C. Differences between similar measurements on models A and B and models A and C were statistically analyzed. Two groups of two researchers measured each distance 10 times, so every measurement was performed 40 times. The time needed for the measurements was recorded and analyzed.

**RESULTS:** Measurements on digimodels and CBCTs were statistically different. On average, measurements on CBCT images were larger. Measurements on digital models were more accurate. Measurements on segmented CBCT models were more accurate than measurements on

CBCTs. The mean measuring time on CBCT images was 14.5 minutes longer.  
**CONCLUSION:** Measurements on CBCT images are less accurate compared with those on Digimodels™. The segmentation of the dentition on an AnatoModel™ increases the accuracy of the measurements. The measurement procedure for CBCT images takes more time compared with measurements on Digimodels™.

## OP 20

Thursday, June 17 - at 9:20

### STABILITY OF SURGICALLY ASSISTED RAPID MAXILLARY EXPANSION WITH A BONE-BORNE DISTRACTOR

**Y Deleurant**<sup>1</sup>, A Zimmermann<sup>2</sup>, T Peltomäki<sup>3</sup>, <sup>1</sup>Clinic for Orthodontics and Pediatric Dentistry, Institute of Dentistry, University of Zürich and <sup>2</sup>Private practice, Zürich, Switzerland, <sup>3</sup>Tampere University Hospital, Finland

**AIM:** Tooth-borne surgically assisted rapid maxillary expansion (SARME) is widely used treatment modality in adolescent and adult patients with maxillary transverse deficiency. To prevent undesired dental side-effects bone-borne expansion devices have been introduced. The aim of the study was to evaluate the stability of maxillary expansion carried out by SARME with a bone-borne distractor.

**SUBJECTS AND METHOD:** Fifty patients (35 females, 15 males who had undergone SARME because of a bilateral (n=33) or unilateral (n = 13) crossbite, or no crossbite but who required maxillary expansion to facilitate mandibular advancement in a second stage osteotomy (n = 4). The surgical technique involved an osteotomy on the lateral maxillary walls and in the palatal midline. A bone-borne distractor (TPD, surgi-tec, B-Brugge) was secured in the region of the second premolars/first molars. After 5 days the patients activated the appliance once a day, i.e. 0.333 mm/day. The necessary expansion (no over expansion) was determined by clinical need, leading to an activation period of 15-27 days, and a retention period of 12 weeks. Study models were obtained before surgery (T0, n=50, mean age 19.0 years, range 14.6-35.4 years), once expansion and orthodontic treatment were completed (T1, n = 50, mean T0-T1 interval 1.6 years, range 1.3-2 years), and at follow-up (T2, n = 38, mean T0-T2 interval 2.2 years, range 2.0-6.8 years). The models were photographed and transverse distances between the canines and the 1st molars measured gingivally and occlusally on the computer screen.

**RESULTS:** Mean expansion between the canines was 3.6 mm occlusally and 3.9 mm gingivally, and between the first molars 4.3 and 4.2 mm, respectively. The mean relapse varied between 1.0 and 1.3 mm at T2.

**CONCLUSION:** Despite the use of bone-borne distractors for SARME, some relapse was found. Therefore, over expansion is still recommended.

## OP 21

Thursday, June 17 - at 10:00

### FACIAL SYMMETRY ANALYSIS IN WELSH ADOLESCENTS USING OPTICAL LASER SURFACE SCANNING

**J Djordjević**<sup>1</sup>, A Toma<sup>2</sup>, A Zhurov<sup>2</sup>, B Glišić<sup>1</sup>, S Richmond<sup>2</sup>, Schools of Dentistry, <sup>1</sup>University of Belgrade, Serbia and <sup>2</sup>Cardiff University, Wales

**AIM:** Different diagnostic approaches, mainly two-dimensional, have been used for facial symmetry analysis. Recent advances in laser scanning technology provide the opportunity to examine human faces more accurately in three dimensions. The aim of this study was to determine the extent of facial symmetry in healthy adolescents and to explore if there is any difference between the genders.

**SUBJECTS AND METHOD:** Laser scanning was performed on 270 Welsh adolescents (123 males, 147 females) of Caucasian origin, average age 15.5 years. They were randomly selected from a larger longitudinal study known as ALSPAC. Images were transferred into commercial software, processed and landmarked. Facial symmetry data were collected using an in-house developed subroutine. For each subject, a mirror facial shell was generated and superimposed on the original shell to create a symmetric face and establish a true mid-sagittal plane. A coincidence of the original face to the symmetric face (percentage of symmetry) was measured, with a 0.5 mm tolerance level. In addition, three angular and 14 linear measurements were made.

**RESULTS:** The average distance of an original shell to a mirror shell was higher in males (0.63 ± 0.16 mm) than in females (0.55 ± 0.13 mm) and statistically significant (P < 0.01). There was also a statistically significant

difference ( $P < 0.01$ ) in the extent of three-dimensional facial symmetry between males ( $53.49 \pm 10.73\%$ ) and females ( $58.46 \pm 10.25\%$ ). Two of three angular measurements showed statistically significant differences ( $P < 0.01$ ). None of the 14 linear measurements were statistically different. Point pogonion had the highest deviation from the mid-sagittal plane ( $1.05 \pm 0.9$  mm in males and  $0.93 \pm 0.73$  mm in females).

**CONCLUSION:** Facial symmetry can be analysed accurately using laser scanning. Male faces were less symmetric than those of females.

## OP 22

Thursday, June 17 - at 10:10

### ALVEOLAR BONE LOSS DURING ORTHODONTIC TREATMENT: FINDINGS WITH CONE BEAM COMPUTED TOMOGRAPHY

**K Hansen**<sup>1</sup>, H Lund<sup>2</sup>, <sup>1</sup>Department of Orthodontics, University Clinics of Odontology and <sup>2</sup>Department of Oral and Maxillofacial Radiology, The Sahlgrenska Academy, University of Gothenburg, Sweden

**AIM:** Cone beam computerised tomography (CBCT) has given orthodontics a new diagnostic dimension. The Department of Orthodontics, Gothenburg, Sweden has an ongoing study for monitoring root resorption during standardized orthodontic treatment in 170 patients with Class I molar relationship. The analyses of hard tissue changes started after 100 patients had finished their treatment. The aim of this study was to determine buccal and lingual/palatal alveolar bone level changes in patients treated with full fixed appliances.

**SUBJECTS AND METHOD:** CBCT scanning (3D Accuitomo FPD, Morita, Japan) was performed before and after treatment of 101 Angle Class I patients with bimaxillary crowding. The standardised treatment plan included full fixed appliances in both jaws and extraction of four premolars. The buccal and lingual/palatal bone levels were measured for all teeth from first molar to first molar in both jaws and the buccal-lingual root position in the alveolar process was measured. Twenty patients had one or more tooth/teeth displaced out of the alveolar process to the extent that at least half of the root was displaced outside the alveolar process with no detectable bone covering the apical foramen. The most common teeth were the mandibular incisors and, in some patients, the buccal roots of the maxillary molar. The patients with displaced lower incisors were compared with a group of patients with no displaced incisors after treatment.

**RESULTS:** Statistically significant differences were found in the degree of proclination of the lower incisors that occurred during treatment. The displaced incisors were on average proclined 10 degrees more than the non-displaced incisors.

**CONCLUSION:** CBCT reveals information in new dimensions and changes that previously were not visible can now be analysed in detail. The clinical significance of these new findings is difficult to quantify as the clinical treatment procedures have remained more or less unchanged for decades.

## OP 23

Thursday, June 17 - at 10:20

### CAN PIEZOELECTRIC SURGERY FACILITATE ORTHODONTIC TREATMENT?

**M Finotti**<sup>1</sup>, M Roberto<sup>1</sup>, M Del Torre<sup>2</sup>, F A Miotti<sup>1</sup>, <sup>1</sup>Orthodontic Department, University of Padova, and <sup>2</sup>Private Practice, Padova, Italy

**AIM:** Intrusion and bodily lower molar distalization are complex orthodontic movements with frequently a low success rate due to biomechanical problems. Various techniques, such as miniscrews, have been proposed to improve system efficiency. The aim of this presentation is to illustrate methods of distalising lower molars and intruding overerupted molars avoiding side-effects.

**SUBJECTS AND METHOD:** Twenty five adult patients treated with the support of piezo-microsurgery techniques. One distalization procedure and two parallel horizontal osteotomies are carried out distal to the last molar to be distalized. Two further osteotomies are carried out perpendicular to the previous ones, one distal and the other mesial, 2 mm from the molar. The arches are bonded with the straightwire technique. The distalizing device consists of a 0.016 x 0.022 inch wire with an open coil spring. A force of 100 g is applied to allow distalization of the molar along the bone defect. 2. Intrusion procedure: two osteotomies are carried out mesial and distal to the overerupted element. Three millimetres apical to the molar two horizontal osteotomies are performed at a distance corresponding to the molar intrusion required, and the cortical bone is removed. The arches

are bonded with straightwire technique and at the end of the surgical procedure a 0.016 x 0.022 or 0.018 x 0.022 inch Ni-Ti wire is positioned.

**RESULTS:** Distalization in 15 subjects ranged from 1.5 to 4.5 mm, depending on the amount of distal movement required and was obtained in 16-20 weeks. The required intrusion in 10 patients (from 2.0 to 3.0 mm) was obtained in 12-16 weeks. No side-effects such as anchorage loss, opening of the bite or damage to the periodontal tissues were observed during the procedures.

**CONCLUSION:** The corticotomy technique reduces treatment time, avoiding the side-effects of traditional methods. The follow-up results appear to confirm the validity and stability of the method obtained in a single sitting of piezo-microsurgery.

## OP 24

Thursday, June 17 - at 11:30

### CHANGES IN NASAL CAVITY VOLUME FOLLOWING RAPID MAXILLARY EXPANSION – A THREE-DIMENSIONAL EVALUATION

**S Ozkaya,** A Acar, A Haralambidis, M Ateş, N Küçükkeleş, Department of Orthodontics, Dental Faculty, Marmara University, Nişantaşı, İstanbul, Turkey.

**AIM:** To evaluate the permanence of the changes in nasal cavity volume induced by rapid maxillary expansion (RME) one year after completion of the retention phase of expansion, using three-dimensional (3D) models obtained from computed tomographs (CT).

**SUBJECTS AND METHOD:** Twenty-seven subjects (11 males, 16 females) aged between 11 and 17 years (mean 14.6 years). All patients presented with transverse maxillary deficiency and a bilateral posterior crossbite. The RME procedure was carried out with an acrylic cap splint type Hyrax appliance. Routine orthodontic records, as well as axial tomograms, were obtained before expansion, at the end of the 3 month retention period after active expansion was completed, and one year after the retention period. CT were assessed and analyzed using Mimics 10.11 software by reconstructing 3D images of the nasal cavities in order to calculate volume changes following RME.

**RESULTS:** A significant difference was found between the mean pre-expansion, post-expansion and 1-year follow-up measurements. There was a significant increase of 11.3 per cent in the volume of the nasal cavity due to RME as measured at the end of the 3 month retention period. However follow-up results after 1 year showed that there was a significant decrease in the volume of the nasal cavity compared with the post-expansion measurements. Nevertheless, when compared with the pre-expansion values, there was still a significant ( $P < 0.05$ ) increase in the volume of the nasal cavity 1 year after RME.

**CONCLUSION:** Taking significant relapse into consideration, RME should not be undertaken with the sole purpose of improving nasal respiration, without the presence of a transverse maxillary deficiency.

## OP 25

Thursday, June 17 - at 11:40

### BONE DENSITY OF THE MIDPALATAL SUTURE SEVEN MONTHS AFTER RAPID PALATAL EXPANSION IN ADULTS

**W Harzer,** S Petrick, E Tausche, Orthodontics, Technical University of Dresden, Germany

**AIMS:** To analyze changes in bone density of the midpalatal suture after surgically assisted rapid palatal expansion (SARPE) with the bone-borne Dresden distractor by means of computed tomography (CT), and comparison of pre-surgical findings with a control group.

**SUBJECTS AND METHOD:** Sixteen adult patients (mean age 24.5 years) who underwent axial CT scans before and 7 months after SARPE. CT image fusion was performed for the bone of the midpalatal suture. Sixty-six controls (mean age 25.7 years) served for comparison of age-related bone density. Bone structure and density were assessed in the coronal plane at the anterior, median and posterior level.

**RESULTS:** The highest density was found in the posterior part [1046 Hounsfield units (HU)] before expansion. Bone density 7 months after SARPE was only 48 per cent (anterior), 53 per cent (median) and 75 per cent (posterior) in comparison with pre-surgical values. The values in the control group were fairly equal (889 to 900 HU) for all aspects.

**CONCLUSION:** Seven months after SARPE the density of the midpalatal

suture is only half to three-quarters in comparison with pre-treatment values. For preservation of the resistance against the forces from the non-split posterior part, the time for retention should be lengthened.

## OP 26

Thursday, June 17 - at 11:50

### ASSESSMENT OF TONGUE POSTURE AND FUNCTION IN CLINICAL ORTHODONTICS – IS IT RELIABLE?

**S Melink**<sup>1</sup>, J Volk<sup>1</sup>, I Hočevnar-Boltežar<sup>2</sup>, M Mušič<sup>3</sup>, M Ovsenik<sup>4</sup>,  
Department of Orthodontics, University Medical Centre, <sup>2</sup>Department of Otorhinolaryngology University of Ljubljana, <sup>3</sup>Institute of Oncology and <sup>4</sup>Department of Orthodontics, University of Ljubljana, Slovenia

**AIM:** To assess the reliability of clinically diagnosing tongue posture and tongue function using ultrasonography.

**SUBJECTS AND METHOD:** Tongue posture and swallowing pattern were clinically assessed in 26 children (14 girls, 12 boys; mean age 5.5 years) with a unilateral posterior crossbite by an orthodontist (O) and an otolaryngologist (ENT). Ultrasound (US) analysis of tongue posture and swallowing pattern was carried out for all children using the Voluson 730 Expert US system with a 3D convex transducer RAB 2-5 MHz. The data were statistically analysed using the Wilcoxon signed ranks test.

**RESULTS:** O and ENT clinical assessments of tongue posture were statistically significantly different from the results of the US analysis ( $P = 0.002$  and  $P = 0.05$ , respectively). Assessment of swallowing pattern by the O showed similar results to the US analysis ( $P = 0.527$ ), while the assessment of the swallowing pattern by the ENT was statistically significantly different from the results of US analysis ( $P = 0.001$ ).

**CONCLUSION:** Clinical assessment of tongue posture and function is not reliable in children in the primary dentition period. Assessment of tongue posture and function during clinical examination in small growing children is unreliable. A reliable, non-invasive, objective, quick and easy method for observation of tongue posture and function could, therefore, become an important diagnostic tool.

## OP 27

Thursday, June 17 - at 12:00

### THREE-DIMENSIONAL EVALUATION OF TOOTH- AND BONE-BORNE SURGICALLY ASSISTED RAPID MAXILLARY EXPANSION

**R Nada**<sup>1,2</sup>, Y Mostafa<sup>1</sup>, R Novraz<sup>2</sup>, S Bergé<sup>3</sup>, A M Kuijpers-Jagtman<sup>2</sup>,  
<sup>1</sup>Department of Orthodontics and Dentofacial Orthopedics, Cairo University, Egypt and Departments of <sup>2</sup>Orthodontics and Oral Biology and <sup>3</sup>Oral and Maxillofacial Surgery, Radboud University Nijmegen Medical Centre, Netherlands

**AIM:** Dentoalveolar changes following surgically assisted rapid maxillary expansion (SARME) are particularly challenging to assess on frontal radiographs. Previous studies have generally relied on dental models, which provided information about the dentition rather than the alveolar bone. Cone beam computed tomography (CBCT) scans may allow a more detailed evaluation of dental and alveolar changes following maxillary expansion using bone-borne compared with tooth-borne expansion devices. The aim of this study was to evaluate three-dimensionally changes in dentoalveolar structures following SARME using a tooth-(Hyrax) or a bone- [Transpalatal distractor (TPD)] borne expansion device.

**SUBJECTS AND METHOD:** The study design was a prospective two group controlled clinical trial. Thirty skeletally mature patients with transverse maxillary hypoplasia undergoing SARME were included. In group A, expansion was performed using a Hyrax expander with bands on the first premolars and molars, while for group B a TPD was used. CBCT scans were obtained before treatment and 19 ( $\pm 5$ ) months later, at the end of fixed appliance treatment. Three-dimensional (3D) models were constructed from the CBCT data and superimposed using voxel based matching. Distance maps between the superimposed models were used to evaluate maxillary alveolar changes. Dental linear measurements such as intermolar, interpremolar and intercanine width were also measured on the 3D CBCT models.

**RESULTS:** There was a statistically significant positive correlation between the amount of alveolar change measured on the superimposed models and dental expansion in both groups ( $P = 0.007$ ). There was no statistically significant difference between the Hyrax and TPD in either parameter ( $P > 0.05$ ).

**CONCLUSION:** After a period of fixed appliance therapy, dentoalveolar changes produced by SARME are comparable when using either a tooth- or bone-borne expansion device.

## OP 28

Thursday, June 17 - at 12:10

### INTERCEPTIVE TREATMENT OF PALATALLY DISPLACED CANINES - A RANDOMIZED CLINICAL STUDY

**J Naoumova<sup>1</sup>**, H Kjellberg<sup>1</sup>, J Kuroi<sup>1</sup>, R Palm<sup>2</sup>, Departments of

<sup>1</sup>Orthodontics and <sup>2</sup>Oral and Maxillofacial Radiology, Institute of Odontology at the Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden

**AIM:** Interceptive treatment of palatally displaced canines (PDC) has, from an evidence-based point of view, not been sufficiently evaluated. The aim of this study was to evaluate and analyse the effect of extracting the primary canine to intercept the palatal displacement of maxillary canines

**SUBJECTS AND METHOD:** Twenty-four consecutive patients, out of 60 scheduled, participated in the study. The patients were referred from 22 general dental practices. All met the following inclusion criteria: 10-13 years of age, uni- and/or bilateral PDC(s), no previous orthodontic treatment, no crowding in the posterior part of the maxilla, no resorption of the incisor(s) or lateral(s). The patients were randomized into two groups with either extraction or non-extraction of the primary canine. Four girls and five boys had unilaterally displaced canines, and in seven girls and eight boys, the canines were bilaterally displaced. In the latter group, the right or left primary canine was randomized to either extraction or non-extraction. The outcome measures that were assessed were: changes in the angulation of the permanent canine to the vertical plane from the frontal and sagittal view, and the amount of eruption vertically and buccally. All measurements were performed with a cone beam computed tomograph (3D Accuitomo FPD) at baseline and after six and 12 months. In this preliminary evaluation, the results from the six month control are presented. Two-way ANOVA was used for statistical comparison of the groups.

**RESULTS:** There was a statistical improvement in the angulation from a frontal view and the amount of eruption buccally was seen in the extraction group compared with the control group.

**CONCLUSION:** Extraction of the primary canine to intercept a PDC is effective in creating a favourable eruption path. The 15 bilateral cases showed uniform results – only improvement of the palatal position on the extraction side.

## OP 29

Thursday, June 17 - at 12:20

### CURRENT REMARKS ON BOLTON ANALYSIS – A METRIC STUDY

**S Wriedt**, A Jaroni, H Wehrbein, Department of Orthodontics, J. Gutenberg-University, Mainz, Germany

**AIM:** To re-evaluate the results of Bolton's tooth size discrepancy (TSD) analysis and to find the factors influencing TSD.

**MATERIALS AND METHOD:** At the start and the end of orthodontic treatment, the tooth widths of 200 consecutive patients were measured using model casts and a digital calliper. Patients with missing teeth, prosthetic restoration or fillings in the interdental area were excluded. The Peer Assessment Rating (PAR) index was measured and calculated. Statistical analysis (descriptive, Pearson's correlation) was undertaken using the Statistical Package for Social Sciences, version 17.0 (SPSS Inc., Chicago, Illinois, USA).

**RESULTS:** Compared with the original Bolton standards (overall 91.3 ± 1.91%; anterior 77.2 ± 1.65%) slightly higher values were found: overall 91.98 ± 1.79 per cent; anterior 78.06 ± 2.15 per cent. The new values for anterior ratio showed better correspondence with the curve according to Gauß, while using the original Bolton standards more patients were found in the category >2 SD. A group of 63 patients, who had a PAR Index = 0 at the end of treatment, showed values for overall ratio of = 91.95 ± 1.76 per cent and of anterior ratio = 78.01 ± 1.94 per cent. For 50 patients who had tooth shaping or composite build-up during treatment their values for overall ratio changed from 92.10 ± 1.87 per cent to 91.50 ± 1.94 per cent, and anterior ratio from 78.17 ± 2.40 per cent to 77.12 ± 2.35 per cent. The widths of the upper lateral incisors, lower second premolars and the lower first molars had an influence on the values of anterior and overall ratio.

**CONCLUSION:** TSD standards vary at different times and between different populations. Narrow upper incisors are the reason for new standards, especially the anterior ratio. In addition to the Tonn relationship, the width of the lower first molars can be an indicator for TSD. Treatment outcome of PAR = 0 seems to be influenced more by other factors than Bolton standards.

### OP 30

Thursday, June 17 - at 12:30

#### STABILITY OF SURGICALLY ASSISTED RAPID MAXILLARY EXPANSION AND RAPID MAXILLA EXPANSION

**A T Altug**<sup>1</sup>, G Kurt<sup>2</sup>, M S Atac<sup>3</sup>, H A Karasu<sup>4</sup>, Departments of <sup>1</sup>Orthodontics and <sup>4</sup>Oral and Maxillofacial Surgery, Ankara University, <sup>2</sup>Department of Orthodontics, Erciyes University, Kayseri, <sup>3</sup>Department of Oral and Maxillofacial Surgery, Gazi University, Ankara, Turkey

**AIMS:** To evaluate the changes in dental, skeletal, nasal structures and nasopharyngeal airway 5 years after rapid maxillary expansion (RME) and surgically assisted (SARME), and to compare these changes with a control group.

**SUBJECTS AND METHOD:** Group 1 comprised 10 patients (6 males, 4 females) with a mean age of 15.51 years (range 13.33-17.58 years), treated with RME, and group 2, 10 patients (7 males, 3 females) with a mean age of 19.01 years (range 16.25-25.58 years) treated with SARME. All subjects had maxillary arch width deficiency with bilateral crossbites. The control group consisted of 10 untreated, skeletal Class 1 subjects (6 males, 4 females) with a mean age of 15.27 years (range 13.42-17.00 years) matched to group 1 for gender and age. Lateral cephalometric and postero-anterior films were taken before expansion (T1), post-expansion (T2) and three (T3) and five (T4) years after the retention period. Analysis of variance (ANOVA) and Duncan's tests were used to compare the changes between the groups.

**RESULTS:** After RME and SARME, significant increases were observed for dental, skeletal and nasal transverse widths ( $P < 0.01$ ). No statistically significant differences were found in the measurements related to soft palate morphology or nasopharyngeal dimensions among the SARME, RME or control groups; however, increases in soft palate angulation, and superior and inferior pharyngeal spaces following expansion/control were greater in the SARME group than in other groups. At T3, some of the increases in both transverse and sagittal dimensions of the nasomaxillary complex were lost, but without any statistical significance. The changes between T3 and T4 were almost negligible.

**CONCLUSION:** Both RME and SARME procedures remained stable, more stable and effective with SARME, at the 5 year follow-up with some amount of relapse, compared with the control group.

### OP 31

Thursday, June 17 - at 15:00

#### DOES CONE BEAM COMPUTED TOMOGRAPHY DEPICT THE ANATOMICAL TRUTH? A COMPARATIVE STUDY ON LOWER ANTERIOR ALVEOLAR BONE

**R Patcas**<sup>1</sup>, L Müller<sup>1</sup>, L Signorelli<sup>1</sup>, O Ullrich<sup>2</sup>, T Peltomäki<sup>3</sup>, <sup>1</sup>Clinic for Orthodontics and Pediatric Dentistry and <sup>2</sup>Institute of Anatomy, Faculty of Medicine, University of Zurich, Switzerland and <sup>3</sup>Dental and Oral Diseases Outpatient Clinic, Tampere University Hospital, Finland

**AIMS:** To determine the accuracy of cone beam computed tomography (CBCT) by measuring the alveolar bone covering the lower anterior teeth, and to compare radiographic with physical measurements.

**MATERIALS AND METHOD:** CBCT in two resolutions (0.125 and 0.4 mm Voxel) and analogue lateral cephalograms were obtained of eight decapitated heads. A digital calliper was used to measure the width of the attached gingiva on all lower anterior teeth ( $n = 51$ ). After removing the gingiva, the vertical position of the anterior alveolar bone margin (ABM) was measured. Additionally, two sites per tooth were selected to determine the thickness of the alveolar bone. The physical bony measurements were compared with the CBCT measurements and with the ABM identified on the lateral cephalograms. The data was evaluated using the Bland-Altman method and independent t-tests. Spearman's correlation was used to compare the physical findings (bone and gingiva) and the cephalometric data.

**RESULTS:** Measurements obtained with CBCT differed only slightly from the physical findings with a mean difference of 0.37 mm (high resolution,

CI 99% 0.22-0.55 mm) and 0.7 mm (low resolution, IC 99% 0.37-1.01 mm). These differences were, however, not statistically significant. The identified ABM on the lateral cephalogram did not correlate with the physical measurements ( $P = 0.65$ ). The vertical position of the ABM was highly significantly correlated with the vertical localization of the mucogingival junction ( $P < 0.001$ ). On average, this line was 1.6 mm more apical than the ABM (CI 95% 1.35-1.97 mm).

**CONCLUSION:** Both CBCT resolutions provided highly accurate data and reliably depict the anatomical truth. The identified ABM on the lateral cephalogram does not seem to correspond with the physical measurements. The mucogingival junction is a good landmark to ascertain correct localization of the ABM.

## OP 32

Thursday, June 17 - at 15:10

### A RANDOMIZED CONTROLLED TRIAL FOR ONE YEAR AFTER ORTHODONTIC TREATMENT, ASSESSED BY QUANTITATIVE LIGHT FLUORESCENCE

A Adeyemi, G Komarov, S M Pender, S M Higham, N Pender, University of Liverpool, England

**AIMS:** A variety of techniques can be used to determine demineralisation of teeth after orthodontic appliance treatment. Quantitative light fluorescence (QLF) allows non-invasive longitudinal monitoring (Higham et al., 2009). Lesions found by QLF are not always clinically evident. The aim of this randomised controlled trial was to detect the prevalence of demineralising enamel after orthodontic treatment in a patient cohort treated with fixed appliances, and to longitudinally monitor the effect of three toothpastes with differing fluoride concentrations on the enamel over a 12 month period.

**SUBJECTS AND METHOD:** Patients aged 14 years and over at debond were recruited until 90 had completed 12 months of post-treatment data collection on six occasions at debond, 2 weeks and 1, 3, 6 and 12 months. Digital and QLF images were obtained of the labial surfaces of the six upper and six lower anterior teeth. The patients were stratified into cells by the number of presenting lesions. Thirty patients drawn from similar strata were given toothpaste with fluoride at either 250, 1100 or 2800 ppm blind from the clinical trial team and patient. The inclusion criterion was at least one demineralised lesion detected by QLF.

**RESULTS:** Twenty-four per cent had 1-2; 55 per cent had 3-6 and 21 per cent had 7-12 demineralized lesions. The lesion size, measured in terms of area of tooth surface, reduced significantly over 12 months for each of the three toothpaste trial groups by 30-31 per cent. There was no difference between the groups after 12 months, which was confirmed by an ANCOVA model. The earliest significant remineralisation was at 3 months with both 250 and 1100 ppm ( $P < 0.05$ ). At 6 months, the 2800 pm toothpaste had a significant effect ( $P < 0.01$ ).

**CONCLUSION:** QLF discriminates changes in enamel demineralisation not clinically apparent. All patients using fluoride toothpastes showed remineralised enamel over a 12 month period. The effect is detected later when 2800 ppm fluoride is used.

Higham S, Pender N, de Josselin de Jong E, Smith P 2009 Application of biophysical technologies in dental research. Journal of Applied Physiology (in press)

## OP 33

Thursday, June 17 - at 15:20

### COMPARATIVE EFFECTS OF MANDIBULAR INCISOR INTRUSION USING MINI-IMPLANTS AND A UTILITY ARCH


E Aydogdu<sup>1</sup>, Ö Polat-Özsoy<sup>2</sup>, M Coskun<sup>3</sup>, B F Oduncuoglu<sup>4</sup>,

<sup>1</sup>Private Practice and Departments of <sup>2</sup>Orthodontics, <sup>3</sup>Radiology and

<sup>4</sup>Periodontology, Faculty of Dentistry, Baskent University, Ankara, Turkey

**AIM:** To evaluate dentofacial, periapical, and gingival effects of mandibular incisor intrusion obtained using mini-implants, and to compare these with the effects of an intrusion utility arch.

**SUBJECTS AND METHOD:** Twenty-six individuals at the post-pubertal growth period who had an increased overbite of more than 5 mm. The subjects were randomly divided into two equal groups. In group 1 mandibular incisor intrusion was achieved using a segmented archwire



supported by two mini-implants placed between the roots of the lateral incisors and canines. In group 2 mandibular incisor intrusion was carried out using a utility arch. The dentofacial and periodontal effects were evaluated on lateral cephalograms, periapical radiographs, computed tomograms and orthodontic models. Gingival index scores were also recorded during treatment. Parametric and non-parametric statistical analyses were used to determine any significant differences between and within the groups at the 95 per cent confidence interval.

**RESULTS:** Both intrusion methods were found to be successful for mandibular incisor intrusion. In both groups protrusion of incisors were observed. Distal tipping of the mandibular molars was noted in group 2. In group 1, since there was no posterior anchorage unit, intrusion did not cause a significant change in molar position. Both techniques caused a minimal amount of apical root resorption and the difference between the groups was not significant. Tomographic evaluation revealed a significant decrease in the density of apical root material of the mandibular left lateral incisor in group 2. The increase in scores for probing depth, keratinized gingival and plaque index, were similar between the groups after treatment.

**CONCLUSION:** Mandibular incisor intrusion with mini-implant anchorage or an intrusion utility arch results in similar changes in dentofacial, periapical, and gingival regions.

### OP 34

Thursday, June 17 - at 15:30

#### MANDIBULAR INCISOR ALIGNMENT AND DENTAL ARCH CHANGES AFTER EXTRACTION OF PRIMARY CANINES

**A Sjögren**<sup>1,2</sup>, J Huggare<sup>2</sup>, <sup>1</sup>Department of Orthodontics, Postgraduate Dental Education Centre, Örebro and <sup>2</sup>Department of Dental Medicine, Division of Orthodontics, Karolinska Institutet, Huddinge, Sweden

**AIM:** To investigate the early effects on mandibular incisor alignment and changes in dental arch dimensions after extraction of the primary canines.

**SUBJECTS AND METHOD:** Seventy-three Swedish children in the early mixed dentition, were randomized into one extraction ( $n = 32$ ) and one control ( $n = 41$ ) group. Impressions for dental casts were taken at baseline (T0) and after 1 year (T1). Changes over time and between groups were investigated for mandibular incisor irregularity index, rotation and maxillary and mandibular arch dimensions (arch width, length and circumference), overjet and overbite.

**RESULTS:** Mandibular incisor irregularity decreased more in the extraction group than in the control group (1.2 versus 0.7 mm;  $P < 0.01$ ), although with a wide range for both groups. A 50 per cent reduction or more of irregularity was recorded for 10/32 children in the extraction group and 1/41 in the control group. Changes in rotation  $>10$  degrees for the lateral incisors were twice as common in the extraction group (42 versus 20%;  $P < 0.01$ ), while the central incisors showed minor changes between groups ( $P = 0.02$ ). A significant decrease ( $P < 0.01$ ) in maxillary and mandibular arch length and arch circumference from T0 to T1 was recorded in the extraction group along with a significant increase in maxillary and mandibular arch circumference and mandibular arch width in the control group. Subjective assessment of alignment in the extraction group revealed that 84 per cent improved while 16 per cent showed no change. However, the association with objective assessment of changes in rotation was only moderate.

**CONCLUSION:** Extraction of the primary canines gave significant but small improvements in mandibular incisor alignment after 1 year. However arch length and circumference were reduced leading to a decrease in available space for the permanent dentition.

### OP 35

Thursday, June 17 - at 15:40

#### SKELETAL AND DENTAL EFFECTS OF A NEW MAGNETIC FUNCTIONAL APPLIANCE IN CLASS II CORRECTION

**N E Tarraf**<sup>1</sup>, P Taylor<sup>1</sup>, M A Darendeliler<sup>1</sup>, T Baccetti<sup>2</sup>, <sup>1</sup>University of Sydney, Australia and <sup>2</sup>University of Florence, Italy

**AIM:** To evaluate the skeletal and dental effects of a new magnetic fixed functional appliance (Magnoglide) compared with an untreated control group.

**SUBJECTS AND METHOD:** A prospective study of 42 consecutive Class II division 1 subjects treated with the Magnoglide compared with 30 untreated Class II controls matched for age gender and cervical vertebral

maturation stage (CS 3-4). Lateral cephalograms were obtained prior to the start of treatment (T1) and at the end of the functional appliance stage (T2). Cephalometric analysis was undertaken using the SO analysis of Pancherz, and linear and angular measurements.

**RESULTS:** There was no statistical difference between the two groups at T1. Class II correction was achieved in all 42 subjects. Pancherz's analysis showed that, on average, overjet correction of 4 mm was achieved. The correction can be attributed to 2.5 mm (62.5%) mandibular change and 1.5 mm (37.5%) maxillary change. The mandibular changes consisted of 2.2 mm (55%) skeletal advancement the chin level and 0.4 mm (10%) of lower incisor advancement. The maxillary changes consisted of 0.5 mm (15%) of restrained growth and 0.9 mm (22.5%) incisor retraction. There was a significant gain in mandibular length as measured by the change in Co-Gn with the treatment group showing 3.3 mm more increase in mandibular length than the control group, as well as a reduction in ANB angle of 1.4 degrees as opposed to an increase of 0.3 degrees in the untreated controls. Changes in lower incisor angulation were statistically insignificant

**CONCLUSION:** The new magnetic functional appliance (Magnoglide) is effective in orthopaedic Class II treatment with minimal dental side-effects.

## OP 36

Thursday, June 17 - at 15:50

### SECONDARY AUTOGENOUS ALVEOLAR BONE GRAFTING: A PROSPECTIVE AUDIT

**H Jeremiah<sup>1</sup>**, R Rimes<sup>1</sup>, M Cooke<sup>1</sup>, D Adlam<sup>2</sup>, <sup>1</sup>Orthodontic Department and <sup>2</sup>Oral and Maxillofacial Surgery, Addenbrooke's Hospital, Cambridge, England

**AIM:** Secondary bone grafting of the alveolar cleft with autogenous bone is an accepted procedure in oro-facial rehabilitation of cleft lip and palate (CLP) patients. The aim of this research was to re-evaluate the clinical process and success of autogenous secondary alveolar bone grafting (ABG).

**MATERIALS AND METHOD:** This was a prospective re-audit of medical case notes and radiograph outcome analysis of ABG using the modified Bergland scoring at Addenbrooke's Hospital, Cambridge, UK. Audit Standards 100 per cent availability of records; 100 per cent ABG to be carried out at ideal age; 81 per cent radiographic ABG success. Data collected from 29 consecutive children with a complete unilateral CLP or a complete bilateral CLP who underwent ABG in 2008 were compared with data from consecutive cases in 2007 and 2006. ABG success was defined by modified Bergland scores of 1 and 2. A single oral and maxillofacial surgeon carried out all ABG.

**RESULTS:** One hundred per cent of notes were available in 2008, 2007 and 2006. The average age at ABG was 10.7 years in 2008, 11.1 years in 2007 and 10.9 years in 2006. Radiograph availability/quality for scoring: 97.1 per cent (34/35) in 2008, 92.7 per cent (38/41) in 2007 and 100 per cent (25/25) in 2006. ABG success of the available sample was 94.1 per cent (32/34) in 2008, 86.8 per cent (33/38) in 2007 and 81.5 per cent (22/27) in 2006.

**CONCLUSION:** ABG success compares favourably with standards. Prospective audit is important for clinical transparency. Re-audit should be undertaken annually.

## OP 37

Friday, June 18 - at 9:00

### ORTHOPAEDIC EFFECTS OF MAXILLARY PROTRACTION WITH OR WITHOUT PALATAL EXPANSION

**A Cesta**, C Berto, T Avventi, F A Miotti, Department of Orthodontics, University of Padova, Italy

**AIM:** Maxillary protraction has become a major treatment option for the correction of developing Class III malocclusions. The frequent maxillary retrusion in such patients, with the limited possibility to influence mandibular growth, has increased the popularity of facemask therapy. The aim of this study was to analyze the skeletal effects of maxillary protraction, with and without rapid palatal expansion (RPE).

**SUBJECTS AND METHOD:** Forty-seven subjects, 15 (group 1) treated with a facemask and RPE, 2) 18 treated with a facemask only and 3) 14 untreated patients with a Class III malocclusion (control group). Radiographs were taken before and immediately after treatment (over a similar time interval for the control group) and at about 2 years post-treatment (33 cases). Statistical analysis was employed (ANOVA).



**RESULTS:** No statistically significant differences between the expansion and non-expansion groups in cephalometric variables or in overall treatment time were observed. Facemask therapy was effective in correcting the Class III malocclusion. Comparison of the control and treated groups showed significant treatment effects (ANB: group 1 +3.06<sup>o\*\*\*\*</sup>, group 2 +3.36<sup>o\*\*\*\*</sup>), mainly characterized by maxillary advancement (A-McNamara group 1 +2.9 mm\*, group 2 +1.61 mm\*) and a restraining effect on mandibular growth (Harvold Md: group 1 +3.02 mm ns, group 2 +1.19 mm ns, group 3 +5.85 mm\*). A slight increase in the intermaxillary vertical relationship (due to a combination of counterclockwise maxillary rotation and clockwise mandibular rotation) was found in the treated groups, especially in group 1 (Harvold Hf +4.26 mm\*). Long-term, the treated patients tended to resume a Class III growth pattern, though the correction of the malocclusion was maintained.

**CONCLUSION:** A Class III malocclusion can be successfully treated with a facemask. The need for palatal expansion, in the absence of a transverse maxillary discrepancy, was not supported by the findings of this study.

### OP 38

Friday, June 18 - at 9:10

#### INFLUENCE OF TONGUE POSTURE ON JAW MORPHOLOGY – A THREE-DIMENSIONAL EVALUATION

F Farčnik<sup>1</sup>, J Primožič<sup>1</sup>, S Richmond<sup>2</sup>, A Drevenšek<sup>3</sup>, M Pisek<sup>3</sup>, <sup>1</sup> Orthos, Institute of orthodontics and maxillofacial orthopaedics and <sup>3</sup>Department of Dental and Jaw Orthopaedics, University of Ljubljana, Slovenia and <sup>2</sup>Department of Orthodontics, University of Cardiff, Wales

**AIM:** To evaluate the influence of tongue posture on jaw morphology in two groups of untreated patients, one with a Class III (C3) and the other with a Class I malocclusion (control group-CG).

**SUBJECTS AND METHOD:** Twenty C3 patients (9 males, 11 females, mean age 19.2 ± 4.64 years) and 20 CG patients (6 males, 14 females, mean age 17.4 ± 1.68 years). Inter canine and intermolar transverse widths at the cusp and gingival levels were measured on three-dimensional digital images of study casts. Surface area and volume of the palate and the mouth floor were calculated. Tongue position relative to the palatal contour was assessed on lateral cephalograms. ANOVA was used to calculate the differences between the two groups. Correlation between tongue posture and the morphological characteristics of the jaws was calculated using Spearman's correlation coefficient.

**RESULTS:** The C3 patients had statistically significantly greater intermolar transverse mandibular widths at the cusp and gingival level ( $P = 0.047$  and  $0.004$  respectively), and a significantly greater surface area and volume of the mouth floor ( $P = 0.002$ ) than the CG subjects. C3 patients had a statistically significantly lower tongue position relative to the palate compared with the CG ( $P < 0.001$ ). However, there were no statistically significant differences between the two groups, either for palatal volume or for palatal surface area ( $P > 0.05$ ). A significant correlation ( $P = 0.028$ ) was found between the surface area and volume of the mouth floor and the tongue position relative to the palatal contour.

**CONCLUSION:** Tongue posture could influence the morphology of the lower jaw. It is considered to be an important aetiological factor for the transverse dimensions of the lower jaw.

### OP 39

Friday, June 18 - at 9:20

#### LONG-TERM EVALUATION OF MAXILLARY EXPANSION AND FACEMASK THERAPY: A CONTROLLED STUDY

L Franchi, C Masucci, T Baccetti, Università degli Studi di Firenze, Italy

**AIM:** Long-term data on the effectiveness of maxillary expansion and protraction in Class III patients are lacking. The aim of this study was to evaluate the stability of maxillary expansion and facemask therapy (E/FM) in Class III patients at a long-term observation, compared with the effects of growth in well-matched untreated Class III controls.

**SUBJECTS AND METHOD:** The treated group (TG) consisted of 21 patients (11 females, 10 males) with dentoskeletal Class III malocclusions treated with a two-phase protocol consisting of E/FM therapy followed by fixed appliance therapy, while the control group (CG) comprised 18 untreated subjects (12 females, 6 males) with similar dentoskeletal Class III features. Lateral cephalograms of both the TG and CG were analyzed at three time

points, before treatment (T1); after two-phase treatment (T2); and long term (T3). All patients started treatment (T1) before the pubertal growth spurt (CS1 or 2, mean age  $9.5 \pm 1.6$  years), and had completed active skeletal maturation at T3 (CS6, mean age  $18.7 \pm 2.1$  years).

**RESULTS:** At T3 the TG exhibited greater values for ANB angle, Wits appraisal, overjet and molar relationship when compared with the CG. No significant difference between the two groups was found for the changes occurring from T2 to T3.

**CONCLUSION:** Favourable long-term dentoskeletal outcomes were induced by E/FM and fixed appliances in Class III patients. These outcomes remained stable at T3 in 72.5 per cent of the treated subjects. The untreated Class III malocclusion group did not show any tendency toward self-improvement during the post-pubertal interval.

## OP 40

Friday, June 18 - at 10:00

### LONG-TERM FOLLOW UP IN SAGITTAL SPLIT OSTEOTOMY

**I Ilhan**<sup>1</sup>, T Alcan<sup>1</sup>, M Z Guzel<sup>2</sup>, <sup>1</sup>Department of Orthodontics, Faculty of Dentistry, Marmara University, Istanbul and <sup>2</sup>Department of Plastic, Reconstructive, and Esthetic Surgery, Cerrahpasa Faculty, Istanbul University, Turkey

**AIM:** To investigate whether stability of Class III surgical cases is influenced by factors such as the type of surgery, amount of dentoskeletal correction, vertical pattern, or gender.

**SUBJECTS AND METHOD:** Dentoskeletal and profile changes were followed up in 50 sagittal split osteotomy patients (34 females, 16 males) until 2.43 years post-treatment. Two subgroups (double and single jaw) were formed according to the type of surgical correction. The mean age of the study group was 20.87 years (double jaw: 20.17, single jaw: 21.58 years). Rigid fixation with triple miniscrews was used in all subjects, while miniplates were used at the maxillary osteotomy site. Surgery was performed by the same maxillofacial surgeon. Cephalograms were obtained initially, pre- and post-operatively, and at the end of treatment. Ten sagittal, eight vertical, and five soft tissue parameters were analyzed on hand-traced radiographs.

**RESULTS:** Subgroup comparisons revealed no statistically significant differences in the amount of relapse for any of the parameters. For anteroposterior measurements, no significant relationship was found between dentoskeletal correction and the amount of post-treatment relapse. For vertical and soft tissue measurements, some parameters showed negative correlations between treatment changes and relapse amount. Long, short, and normal faced individuals showed no significant difference in the amount of relapse, except for upper incisor inclination (U1-SN). Gender comparisons revealed statistically significant differences in only two parameters, ANB and lower incisor angle (L1-SN).

**CONCLUSION:** Inclusion of a Le Fort I osteotomy does not affect the stability of a sagittal split osteotomy. The amount of relapse was found to decrease with the increasing amount of correction in the vertical and soft tissue parameters. Initial vertical pattern and the extent of relapse, however, seem not to be correlated. Gender also does not play a role in the amount of relapse.

## OP 41


Friday, June 18 - at 10:10

### THREE-DIMENSIONAL ASSESSMENT OF MAXILLARY AND MANDIBULAR CHANGES WITH BONE-ANCHORED MAXILLARY PROTRACTION

**T Nguyen**, M Cornelis, L Cevidanes, J F C Tulloch, H DeClerck, University of North Carolina, Chapel Hill, USA

**AIM:** Conventional treatment for young Class III patients involve extra-oral devices designed to either protract the maxilla or restrain mandibular growth. These methodologies have limited success because the treatment change is primarily dentoalveolar rather than skeletal. The use of skeletal anchorage offers a promising alternative to obtain orthopaedic results with less dental compensations. The aim of this study was to evaluate three-dimensional (3D) changes in the maxilla and mandible of Class III patients treated with bone-anchored maxillary protraction (BAMP).

**SUBJECTS AND METHOD:** Twenty consecutive skeletal Class III patients between the ages of 9 and 13 years (mean:  $11.10 \pm 1.1$  years) treated using Class III intermaxillary elastics and bilateral miniplates (two in the



infra-zygomatic crests of the maxilla and two in the anterior mandible). The patients had cone beam computed tomograms (CBCTs) taken before initial loading (T1), and one year following treatment (T2). 3D models were generated from the CBCTs, registered on the anterior cranial base and analyzed using colour-maps.

**RESULTS:** Sagittal changes in the maxilla, zygoma and mandible were observed in all patients. An average maxillary protraction of 3.75 mm (SD  $\pm$  1.20) with infra-orbital changes of 3.11 mm (SD  $\pm$  0.79) was observed in the BAMP treated patients. Posterior displacement of the mandible at T2 was observed for all subjects (mean 2.9 mm (SD  $\pm$  0.86) for posterior ramus, 2.0 mm (SD  $\pm$  0.79) for the condyles and 0.56 mm (SD  $\pm$  1.55) for the chin. Remodelling of the glenoid fossa at the anterior eminence (0.89  $\pm$  1.38 mm) apposition and resorption at the posterior wall (0.72  $\pm$  0.45 mm) was observed in most patients.

**CONCLUSION:** This new treatment approach offers a promising protraction alternative for maxillary deficient patients as well as restraint of mandibular growth for Class III patients with a component of mandibular prognathism. Future studies with long-term follow-up and comparisons to facemask and chincup therapies are needed to better understand the treatment effects.

## OP 42

Friday, June 18 - at 10:20

### DENTOFACIAL EFFECTS OF BONE-ANCHORED MIDFACE PROTRACTION: A CONTROLLED STUDY\*\*\*

**H De Clerck**<sup>1</sup>, L Cevidanes<sup>1</sup>, T Baccetti<sup>2</sup>, J F C Tulloch<sup>1</sup>, T Nguyen<sup>1</sup>,

<sup>1</sup>Department of Orthodontics, University of North Carolina, Chapel Hill, USA and <sup>2</sup>Department of Orthodontics and Pediatric Dentistry, University of Florence, Italy

**AIM:** To evaluate the treatment effects of bone-anchored maxillary protraction (BAMP; with miniplates in the maxilla and mandible connected by Class III elastics) in Class III malocclusion subjects.

**SUBJECTS AND METHOD:** Twenty-one Class III patients (mean age 11.10  $\pm$  1.8 years) were consecutively treated with the BAMP protocol, and re-evaluated 1 year later. The treated group was compared with a matched control group of 18 untreated Class III subjects. Significant differences between the cephalometric changes in both groups were assessed with an independent sample *t*-test ( $P < 0.05$ ).

**RESULTS:** Intermaxillary skeletal measurements revealed highly significant improvements (Wits appraisal +6.7 mm, SD  $\pm$  1.7). Point A advanced 3.9 mm (SD  $\pm$  1.4) more than in untreated controls, with significant protraction effects at points, Orbitale (+2.9 mm, SD  $\pm$  0.5) and pterygomaxillare (+3.7 mm, SD  $\pm$  1.1). Significant improvements of overjet and molar relationship were recorded, as well as mandibular skeletal measurements at points B (-2.7 mm, SD  $\pm$  1.6) and Pogonion. The mean difference in gonial angle changes between both groups was -4.1 degrees (SD  $\pm$  -1.3). Vertical skeletal changes (NL-ML -0.8°, SD  $\pm$  1.4) and dental compensations were negligible, except for a significant proclination of the lower incisors (L1-ML +1.9°, SD  $\pm$  1.6) in the treated group. Significant soft tissue changes reflected the skeletal modifications.

**CONCLUSION:** Compared with growth of untreated Class III subjects, the BAMP protocol induced an average increment of skeletal and soft-tissue advancement of maxillary structures of approximately 4 mm, and favourable mandibular changes exceeded 2 mm.

## OP 43

Friday, June 18 - at 11:30

### SELF-ESTEEM AND CURRENT WELL-BEING OF PATIENTS WITH CLASS III PROFILES

**M Schulze**<sup>1</sup>, B Kahl-Nieke<sup>1</sup>, I Springer<sup>2,3</sup>, <sup>1</sup>Department of Orthodontics, Center for Dental and Oral Medicine, University Medical Center Hamburg-Eppendorf and <sup>2</sup>Department of Oral, Maxillofacial and Facial Plastic Surgery, University of Kiel, Germany and <sup>3</sup>Aesthetic Clinic, Oslo, Norway

**AIM:** The degree of satisfaction with one's facial or body image may depend on perceptual, developmental and socio-cultural influences, as well as self-esteem. This study addressed the influence of a Class III profile on a patient's facial or body image in randomly selected participants. The objective was to determine whether a Class III profile is associated with an increased likelihood of having a negative facial or body image, impaired well-being and social functioning and a greater willingness to undergo

aesthetic surgery in comparison with subjects with Class I or Class II profiles.

**SUBJECTS AND METHOD:** Three hundred and twenty five subjects (162 females, 163 males, 18-30 years) were photographed, asked to complete an adjective mood scale (AMS) and to rate 46 statements regarding their own appearance and their impact on social functioning as well as their willingness to undergo aesthetic surgery on a visual analogue scale. The subjects were separated into three groups, according to their profile type.

**RESULTS:** The average ratings of one's appearance were shown to be more positive in subjects with a Class I profile as compared with those with Class II or Class III profiles. Items regarding the impact of one's appearance on social functioning were answered more negatively in subjects with impaired well-being as compared with those with normal well being; this was not dependent on profile type. A Class III profile did not influence the well-being of the subjects. Subjects with a Class III profile did not show an increased willingness to undergo aesthetic surgery.

**CONCLUSION:** A Class III profile may cause an impaired facial image even if well-being is normal. Willingness to undergo aesthetic surgery was the same in subjects with Class I, II or III profiles.

#### OP 44

Friday, June 18 - at 11:40

##### SKELETAL RESPONSE TO EARLY MAXILLARY PROTRACTION IN PATIENTS WITH CLEFT LIP AND PALATE

**R S Tindlund**, Bergen Cleft Palate Team, Faculty of Medicine and Dentistry, University of Bergen, Norway

**AIM:** During the last 30 years CLP patients with maxillary deficiency had an interceptive orthopaedic treatment phase to correct anterior and/or lateral crossbites in the primary and early mixed dentition periods. The objective of this study was to evaluate dentofacial growth and development after maxillary protraction before 10 years of ages.

**SUBJECTS AND METHOD:** Seventy-two subjects with various types of cleft types and an anterior crossbite, treated to an acceptable normal occlusion by maxillary protraction using a Delaire facemask. Standardized records, including lateral cephalograms and plaster models, were collected immediately before and after the active treatment period, and at 15 years of age. Two cephalometric variables were chosen for evaluation of the sagittal skeletal effect 1) ANB angle representing the combined maxillo-mandibular change and 2) forward movement of the maxilla represented by an increase in the distance NSP-maxp parallel to the nasion-sella-line. A numerical change greater than or equal to 1.5 (degrees/mm respectively) was classified as a 'fair response' versus a 'poor response' by a change of less than 1.5.

**RESULTS:** During protraction the sagittal intermaxillary relationship showed a mean increase of ANB of +2.5 degrees, however with a range from 0.5 to 7.5 degrees. A fair response was found in 63 per cent of the cases. A fair response of forward movement of the maxilla was found in 44 per cent, with a linear increase of 4.1 mm versus 0.4 mm in the poor response group. There was a close relationship between soft tissue lip changes and the underlying hard tissue changes.

**CONCLUSION:** The skeletal effect of early maxillary protraction is favourable in most cases, but highly individual.

#### OP 45

Friday, June 18 - at 11:50

##### STABILITY OF DEEP OVERBITE CORRECTION IN CLASS II DIVISION 1 ADULTS WITH AND WITHOUT OSTEOTOMY

**R B Kuitert**, A Zentner, Department of Orthodontics, ACTA, Universiteit van Amsterdam and VU University, Amsterdam, Netherlands

**AIM:** Comparison of the stability of overbite correction after surgical and non-surgical correction of Class II division 1 malocclusions with deep bite, hypothesizing more stability after surgical correction.

**SUBJECTS AND METHOD:** Fifty-three 53 adults with Class II division 1 malocclusions and with increased overbite ranging 3.5-11.5 mm, 25 underwent orthodontic treatment only and 28 had surgical-orthodontic treatment including a bilateral sagittal split osteotomy. Pre- and post-treatment and long-term cephalograms eight years after treatment were analyzed by measuring dental and skeletal variables.

**RESULTS:** Pre-treatment overbite was larger in the surgery group. Post-treatment and long-term no differences in overbite were found between



the groups, however during treatment the non-surgery group showed significantly less correction and post-treatment significantly more overbite relapse (57 versus 21 per cent in the surgery group). Both groups showed post-treatment overbite changes between -2.0 and +4.0 mm. The non-surgery group showed stability in 33 per cent, relapse in 60 per cent, and a decrease in 7 per cent. In the surgery group these were 30, 56 and 14 per cent, respectively. No variables showed significant differences between post-treatment changes. The non-surgery group showed significant uprighting of the upper and lower incisors and a decrease in interalveolar distance. During treatment both groups showed similar proclination of the upper and lower incisors and a similar increase of the interalveolar distance, suggesting more stability in the surgery group. Correlation analyses in the non-surgery group did not show any relationship between post-treatment overbite changes and changes in other variables during or after treatment, or with any variable before or after treatment. Post-treatment overbite changes in the surgery group were mainly correlated with vertical skeletal changes during and after treatment

**CONCLUSION:** Surgical-orthodontic treatment seems to provide somewhat better overbite stability after correction of a Class II division 1 deep bite.

#### OP 46

Friday, June 18 - at 12:00

##### LEVELS OF INTERLEUKIN-1 $\beta$ IN ADULTS WITH PERIODONTITIS DURING ORTHODONTIC TREATMENT

**J E Vlachojannis**, S Efstratiadis, U Ersoy, T Cangialosi, J Grbic, Columbia University School of Dental and Oral Medicine, New York, USA

**AIM:** Remodelling of the bone matrix is reflected by cytokine-expression in the gingival crevicular fluid (GCF). The total amount of mediators expressed corresponds to the amount of tooth movement. An increased problem in adults is the occurrence of periodontitis. In humans with severe periodontal disease significantly higher levels of interleukin (IL)-1 were found compared with controls. The purpose of this study was to examine GCF levels of IL-1 $\beta$  during adult human orthodontic tooth movement.

**SUBJECTS AND METHOD:** Eight patients, four with a healthy periodontium and four with a history of periodontal disease. All four molars had bonded molar tubes and were uprighted continuously with a 0.017 X 0.025 TMA uprighting spring using a force of 30 cN. GCF was sampled at all pressure sites twice within two weeks of separation as observation/control before activation and at 1, 7, 14 and 21 days. IL-1  $\beta$  levels were determined with enzyme-linked immunosorbent assay.

**RESULTS:** IL-1 $\beta$  can be detected in the GCF of healthy adult patients and in those with a history of periodontitis. The increased GCF volume of adult patients with a history of periodontitis could either be a marker for subclinical periodontitis or for a genetic predisposition for IL-1 $\beta$  expression. The low mechanical force used might explain why IL-1 $\beta$  remained unaffected.

**CONCLUSION:** Patients with history of periodontitis displayed a higher amount of fluid volume and a higher concentration of IL-1 $\beta$  without significant changes being induced through orthodontic movement. The mechanical stimulation of 30 cN seems to be too low to stimulate IL-1 $\beta$  production. The overall mean total amount of IL-1 $\beta$  remained stable over time, and also when compared with the antagonistic teeth. The findings of this pilot study suggests for future research the inclusion of a greater number of patients, the use of a higher mechanical force, refinement of the GCF sampling and to use as control baseline cytokine levels at experimental teeth.

#### OP 47

Friday, June 18 - at 12:10

##### HO2 MODULATES WOUND CONTRACTION AND INFLAMMATION: IMPLICATIONS FOR CLEFT PALATE PATIENTS

**F Wagener**<sup>1,2</sup>, A Scharstuhl<sup>2</sup>, B Pennings<sup>2</sup>, R Rheden<sup>1</sup>, F Russel<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and Oral Biology and <sup>2</sup>Pharmacology and Toxicology, Radboud University Nijmegen Medical Centre, Netherlands

**AIM:** Cleft palate patients undergo several surgical interventions, resulting in the formation of scars, and increased morbidity in these patients. Following dermal wounding, the severity of scar formation is inversely related to the speed of wound contraction. Therefore, the aim of this study

was to identify novel factors that are involved in dermal wound contraction.

**MATERIALS AND METHOD:** It has previously been demonstrated that heme oxygenase (HO)-activity, via generation of carbon monoxide and bilirubin, modulate wound contraction *in vitro*. In this research HO was studied in more detail using *in vivo* models of wound contraction. Four excisional wounds (4 mm) were made on the dorsal skin of HO2 knock-out (KO) mice and wild type controls (WT), and additionally, wounds were made in hyperbilirubinemic Gunn-UDPTGTJ and WT Gunn-UDPTGT+ rats. Wound size was photographically documented daily up to day 7.

**RESULTS:** In HO2 KO mice wound contraction was significantly slower on days 1, 2, 5 and 7 compared with WT. In contrast, in hyperbilirubinemic rats wound contraction was significantly faster than in WT rats in the early phases of wound healing, suggesting that HO-effector molecules can also modulate wound contraction *in vivo*. Leukocyte infiltration and HO1 expression in the wound area were significantly higher in HO2 KO animals compared with WT at day 7 following wounding. Serum bilirubin levels in HO2 KO mice were significantly higher at baseline than in WT controls and also HO1 mRNA levels on day 2 were significantly increased when compared with WT. Since abrogation of HO2 results in delayed dermal wound closure and hampers resolution of inflammation despite increased HO1 expression, the data suggest that not only HO1, but also HO2 proteins are crucial in the resolution of inflammation.

**CONCLUSION:** Both HO isoforms and their effector molecules are involved in dermal wound contraction and therefore most likely in the process of scar formation.

## OP 48

Friday, June 18 - at 12:20

### CRANIOFACIAL STABILITY IN CROUZON AND APERT PATIENTS ONE YEAR POST LE FORT III DISTRACTION

**J H Reitsma**<sup>1</sup>, E M Ongkosuwito<sup>1</sup>, P H Buschang<sup>2</sup>, B Pahl-Andersen<sup>1</sup>, Departments of Orthodontics, <sup>1</sup>Academic Centre of Dentistry (ACTA), Amsterdam, Netherlands and <sup>2</sup>Baylor College of Dentistry, Dallas, Texas, USA

**AIM:** Le Fort III external distraction osteogenesis (DO) is currently used as a technique for the correction of midface hypoplasia. The technique has been applied in craniofacial surgery of Crouzon and Apert patients. The prevalence of Crouzon syndrome is very low, with about 16 per one million live births and is even lower for Apert syndrome. The aim of this study was to evaluate the post-operative stability of Le Fort III external DO in Crouzon and Apert patients.

**SUBJECTS AND METHOD:** Seven Apert and six Crouzon consecutive patients. Surgery was undertaken between 2003 and 2008. Cephalometric measurements were taken from lateral cephalograms shortly before surgery (T1), at removal of the distraction device, 4 months after surgery (T2), and 1 year after removal of the distraction device (T3). Statistical analysis using a Z-test was performed to compare the syndromic patients with control subjects (n = 482) from the Nijmegen Growth Study.

**RESULT:** At T1, the syndromic patients and control subjects differed significantly for horizontal and vertical measurements. Treatment changes (T1-T2) were significant for all sagittal measurements, with no differences between the Apert and Crouzon patients. The increased vertical values before treatment did not change during DO or at T3. Treatment (T1-T3) showed a significant increase for sagittal measurements; the vertical measurements changes that occurred were not significantly different from control values.


**CONCLUSION:** A Le Fort III external DO in Crouzon and Apert patients seems stable in the sagittal direction at 4 months and one year post-operatively. It appears that for both syndromes the vertical dimensions did not change during treatment.

## OP 49

Friday, June 18 - at 12:30

### CLINICAL IMPLICATIONS OF CRANIOFACIAL ANATOMY IN CLASS III MALOCCLUSIONS

**R Haddad**<sup>1</sup>, J Ghafari<sup>1,2,3</sup>, <sup>1</sup>Division of Orthodontics and Dentofacial Orthopedics, Department of Otolaryngology Head and Neck Surgery, American University of Beirut Medical Center, <sup>2</sup>Lebanese University School of Dentistry, <sup>3</sup>Department of Orthodontics, Dental College, New York University, USA



**AIM:** Variations from normal craniofacial anatomy observed in Class III malocclusions include two features unreported to date in the literature, namely clockwise rotation of the palatal plane and a higher position of sella turcica. The aim of this study was to evaluate morphological characteristics of Class III malocclusion, specifically the relationship between anterior crossbite, cant of the palate and anterior cranial base.

**MATERIALS AND METHOD:** Lateral cephalometric radiographs of a Class III group (80 males; 62 females) and a Class II, division 1 group (41 males, 43 females). The radiographs were analyzed following a standardized method. Cephalometric measurements gauged the overjet, ANB, SN/horizontal, saddle angle, palatal cant to SN and to H. Statistical methods included t-tests and analyses of variance for group differences in continuous measurements, Pearson product moment correlation coefficient for associations between various parameters, and linear and multiple regressions for prediction models.

**RESULTS:** In the Class III sample, the anterior cranial base inclination was more inclined than the average, more in males ( $5.21 \pm 4.3^\circ$ ) than females ( $7.36 \pm 3.43^\circ$ ;  $P = 0.002$ ). The opposite direction was noted in Class II patients, who were compared with a Class III subgroup matched for age. When both malocclusion groups were merged to evaluate the continuum of severity from opposite occlusions, high correlations were noted between overjet and PP/H ( $r = -0.82$ ); SN/H and PP/H ( $r = -0.77$ ), and overjet and SN/H ( $r = 0.67$ ).

**CONCLUSION:** Development and/or severity of Class III dysmorphology relate to intragrowth orthopaedics a sustained anterior crossbite generates forces against the maxilla causing retrognathism that otherwise may not exist. Maxillary retrognathism is apparently more prevalent than mandibular prognathism in mesiocclusion.

## OP 50

Friday, June 18 - at 15:00

### DOES ORTHOGNATHIC SURGERY HAVE AN EFFECT ON TEMPOROMANDIBULAR DISORDERS IN PATIENTS WITH DENTOFACIAL DEFORMITIES?

**C Abrahamsson**<sup>1</sup>, L Bondemark<sup>1</sup>, T Henrikson<sup>2</sup>, E C Ekberg<sup>3</sup>, Departments of <sup>1</sup>Orthodontics <sup>3</sup>Stomatognathic Physiology, Malmö University and <sup>2</sup>Private Practice, Malmö, Sweden

**AIM:** In a systematic review (Abrahamsson et al., 2007) considering alterations in temporomandibular disorders (TMD) after orthognathic surgery, heterogeneous results were found. The aim of this research was to investigate whether patients with dentofacial deformities benefit from orthognathic surgery considering TMD, and furthermore compare if the prevalence of TMD may change during the same period of time in a control group.

**SUBJECTS AND METHOD:** One hundred and twenty one consecutive patients, mean age  $22.5 \pm 7.4$  years, 70 (58%) females and 51 males, referred for orthognathic surgery examined by means of a questionnaire and a clinical examination regarding signs and symptoms of TMD. Three years after treatment, 98 (81%) of the patients completed a follow-up examination. A control group was formed of 56 age- and gender matched individuals, mean age  $23.4 \pm 7.4$  years, 33 (59%) females and 23 males. Thirty-eight (68%) were reached for the follow-up examination. TMD diagnoses according to Research Diagnostic Criteria for Temporomandibular Disorders were used at both examinations.

**RESULTS:** Before treatment 47 (49%) patients had at least one diagnosis of TMD compared with 19 (19%) after treatment ( $P < 0.001$ ). The patients who improved most were those with myalgia without limited opening ( $P = 0.005$ ), disc displacement with reduction ( $P = 0.040$ ) and arthralgia ( $P < 0.001$ ). Before surgery the patient group had a higher frequency of diagnosed TMD ( $P = 0.007$ ) compared with the control group, however after surgery the patient group and control group were comparable regarding frequency of TMD ( $P = 0.934$ ).

**CONCLUSION:** Orthognathic surgery was beneficial in patients with dentofacial deformities considering the frequency of TMD.

### APICAL ROOT RESORPTION AFTER LEVELLING WITH SUPERELASTIC AND MULTISTRANDED STAINLESS STEEL WIRES

**K Alzahawi**<sup>1</sup>, M Mavragani<sup>1</sup>, E Færovig<sup>2</sup>, P Brudvik<sup>1</sup>, O E Bøe<sup>1</sup>,

<sup>1</sup>Department of Clinical Dentistry, Orthodontic Division, University of Bergen and <sup>2</sup>Department of Orthodontics, Faculty of Dentistry, University of Oslo, Norway

**AIM:** Orthodontic root resorption is influenced by both patient and treatment related factors. It can be identified in the early treatment stages. Superelastic archwires are now widely used in everyday clinical practice. However, their effect on orthodontic root resorption has not been thoroughly investigated. The aim of this prospective study was to compare root resorption between teeth treated with superelastic archwires and conventional multistranded stainless steel archwires after the levelling phase of treatment.

**SUBJECTS AND METHOD:** A total of 156 orthodontic patients were consecutively allocated to a treatment group, where levelling was performed either with superelastic heat-activated Sentalloy archwires, or conventional multistranded stainless steel archwires. Pre- as well as post-levelling periapical radiographs were taken for all maxillary and mandibular incisors. After exclusion due to unsatisfactory radiographs, a total of 82 patients (41 in each group) were included in the study. Variables recorded for each patient included gender, age, ANB angle, impacted canines, invagination and anterior crossbite. Crown and root lengths of all incisors were measured on pre- and post-levelling periapical radiographs corrected for image distortion. The percentage of root shortening and root length loss in millimetres was calculated.

**RESULTS:** No significant differences were observed between the two groups concerning apical root resorption for each incisor separately. It was found that apical root resorption of the maxillary lateral incisors was significantly associated with the presence of an anterior crossbite on the affected side.

**CONCLUSION:** Root resorption of maxillary and mandibular incisors after levelling did not differ between patients treated with superelastic and conventional stainless steel wires. An anterior crossbite seemed to be a risk factor for root resorption of the maxillary lateral incisors during the initial stages of orthodontic treatment.

### FRICTIONAL FORCES IN ARCH GUIDED TOOTH MOVEMENT OF CONVENTIONAL AND SELF-LIGATING BRACKETS

**C Bourauel**<sup>1</sup>, M Montasser<sup>2</sup>, S Reimann<sup>1</sup>, T El-Bialy<sup>3</sup>, A Jäger<sup>4</sup>, <sup>1</sup>Oral

Technology and <sup>4</sup>Department of Orthodontics, University of Bonn, Germany and Departments of Orthodontics, <sup>2</sup>University of Mansoura, Egypt, <sup>3</sup>University of Alberta, Edmonton, Canada

**AIMS:** Bracket design and the method of ligation have a crucial influence on frictional resistance in arch guided tooth movement. Manufacturers continuously try to improve the quality and effectiveness of their products and especially self-ligating brackets are reported to have optimised sliding properties. It was the aims of this research to further investigate the differences in the frictional behaviour between various self-ligating bracket systems and standard-edgewise brackets, and to compare the results with former studies.

**MATERIALS AND METHOD:** Three types of orthodontic brackets were investigated using the Orthodontic Measurement and Simulation System: (1) conventional ligating brackets (Victory Series and Mini-Taurus), (2) self-ligating brackets (SmartClip, passive self-ligating bracket, Time3 and Speed, active self-ligating brackets), and (3) a conventional low friction bracket (Synergy). All brackets had a nominal 0.022 inch slot size. The brackets were combined with three archwires (rectangular, 0.019 X 0.025 inch) (1) Remanium (stainless steel, Dentaurum), (2) Nitinol SE [nickel titanium NiTi], and (3) Beta III Titanium [titanium molybdenum (TMA), both 3M Unitek]. Stainless steel ligatures were used with the conventional brackets. Arch guided tooth movement was simulated over a retraction path of 4 mm using a superelastic NiTi coil spring (force 1 N).

**RESULTS:** Force loss due to friction was lowest for the Victory and SmartClip brackets in combination with the steel guiding arch (36 and

35%, respectively) and highest for the Speed and Mini-Taurus brackets in combination with the TMA wire (71 and 64%). Friction gradually increased by 10 per cent for each bracket type in combination with the different wires in the following sequence: steel, NiTi, TMA.

**CONCLUSION:** In agreement with former studies, self-ligating brackets did not show improved performance when compared with conventional brackets.

## OP 53

Friday, June 18 - at 15:30

### CLINICAL TRIAL EVALUATING THE EFFECTS OF CASEIN PHOSPHOPEPTIDE-AMORPH CALCIUM PHOSPHATE PASTE ON WHITE SPOT LESIONS

**G De Pauw**, V Noens, Department of Orthodontics, University of Ghent, Belgium

**AIM:** To investigate the effect of casein phosphopeptide-amorph calcium phosphate (CPP-ACP) paste on the evolution of white spot lesions in orthodontic patients, 4 months after the removal of fixed appliances.

**SUBJECTS AND METHOD:** A prospective, randomized, single-blind study with 19 patients (7 females, 12 males, average age 16 years 1 month; drop outs four). Group A (n = 10) used CPP-ACP paste once daily during 4 months while group C (n = 9) did not use any extra fluoride products beside a normal fluoride toothpaste. The Plaque Index (PI), Gingival Index (GI), surface area of the white spots and the white spot lesion (WSL) index were recorded for all teeth in the upper and lower arches from the left to the right first premolar at three time points. The GI and PI were evaluated with a Cochran Q-test for the surface area, and for the WSL index a *t*-test was used.

**RESULTS:** An improvement in the WSL index was observed in both groups but was statistically significant only in group A. The surface area decreased significantly in both groups during the observation period but was not clinically significant.

**CONCLUSION:** The use of CPP-ACP paste after removal of fixed appliances improves the clinical appearance of white spots and makes them less noticeable. CPP-ACP paste decreases the surface area of a WSL, but this is not clinically significant.

## OP 54

Friday, June 18 - at 15:40

### DENTAL ARCH RELATIONSHIP IN CHILDREN WITH A CLEFT AFTER ONE- AND THREE-STAGE SURGICAL PROTOCOLS

**P Fudalej**<sup>1</sup>, C Katsaros<sup>2</sup>, C Bongaarts<sup>3</sup>, Z Dudkiewicz<sup>1</sup>, A M Kuijpers-Jagtman<sup>3</sup>, <sup>1</sup>Institute of Mother and Child, Warsaw, Poland, <sup>2</sup>University of Bern, Switzerland and <sup>3</sup>Radboud University Nijmegen, Netherlands

**AIM:** To compare dental arch relationship in patients with complete unilateral cleft lip and palate following one- and three-stage surgical protocols.

**MATERIALS AND METHOD:** Dental casts of 61 children (mean age 11.2 years, SD 1.7), consecutively treated in the Warsaw Cleft Centre with one-stage closure of the complete cleft at 9.2 months (SD 2.0), were compared with a sample of 97 patients (mean age 8.7 years, SD 0.9) consecutively treated in the Nijmegen Craniofacial Unit with a three-stage protocol including delayed hard palate closure. The dental casts were assigned random numbers to blind their origin. Four raters graded dental arch relationship and palatal morphology using the EUROCRAN Index. The strength of agreement of rating was assessed with kappa statistics. Independent *t*-tests were run to compare the EUROCRAN scores between one- and three-stage samples and Fisher's exact tests were performed to evaluate differences of distribution of the EUROCRAN grades.

**RESULTS:** Intra- and inter-rater agreement was moderate to very good. Dental arch relationship in the one-stage sample was less favourable than in three-stage group (mean scores 2.58 and 1.97, respectively;  $P < 0.000$ ). Palatal morphology in the one-stage sample was more favourable than in the three-stage group (mean scores 1.79 and 1.96, respectively;  $P = 0.047$ ).

**CONCLUSION:** The dental arch relationship following one-stage repair was less favourable than the outcome of three-stage repair. Palatal morphology following one-stage repair, however, was more favourable than the outcome of three-stage repair.

### ANALYSES OF THE CONDYLAR CARTILAGE AFTER ANTERIOR MANDIBULAR DISPLACEMENT IN JUVENILE PIGS

**T Gredes**, A Spassov, C Kunert-Keil, P Proff, T Gedrange, EMAU Greifswald, Department of Orthodontics, Germany

**BACKGROUND:** The condylar cartilage of the mandible is considered a secondary growth center and represents a joint cartilage different from other cartilage structures. The condylar cartilage shows adaptive and compensatory effects depending on the functional demands and in response to mandibular protrusion after forward displacement appliances.

**AIM:** This study aimed to investigate histologically and molecular-biologically the condylar cartilage after anterior mandibular displacement similar to functional orthopedic treatment.

**METHODS:** Twenty pigs were randomly divided into an experimental group and a control group. The experimental animals were provided bilaterally with synthetic occlusal build-ups in the posterior area which induced anterior displacement of the mandible in terminal occlusion. After 4 weeks the temporomandibular structures were removed en bloc and the condylar cartilage was analyzed histologically and changes in the mRNA content of genes coding for different collagens (col1, col2, col10), matrix metalloproteinases (MMP8, MMP13) and vascular endothelial growth factor (VEGFA) were measured by Real-Time PCR.

**RESULTS:** The experimental animals displayed a significantly increased total cartilage thickness of the posterocranial mandibular condyle. The proliferative layer showed a significant increase, whereas significant differences in thickness were absent in the articular layer. Increased cell proliferation was not observed in the experimental group as compared to the controls. The mRNA expression of col1 and MMP13 during condylar adaptation coincided with that during natural condylar growth. The amount of the expression of col10 was significantly lower ( $p < 0.05$ ) during condylar adaptation, whereas the expression of col2, MMP8 and VEGFA was significantly higher ( $p < 0.05$ ).

**CONCLUSION:** It is suggested that condylar adaptation in growing pigs triggered by mandibular forward positioning is not only a passive adaptive but also a growth affected effect. Results showed that mechanical strain produced by mandibular advancement induced remodelling and revascularization in the posterocranial mandibular condyle. These results were proved by the histological and histomorphometrical analyses.

### THE MASTICATORY SYSTEM UNDER REDUCED FUNCTIONAL LOAD

**T Gruenheid**, G Langenbach, A Zentner, Academic Centre for Dentistry Amsterdam (ACTA), Netherlands

**AIM:** The ability of the masticatory system to adapt to varying functional requirements enables its normal function under changing environmental conditions. The aim of this research was to investigate adaptive changes in neuromuscular activity and phenotypic properties of jaw muscles and mineralization of mandibular bone in response to reduced functional load.

**MATERIALS AND METHOD:** Intramuscular electromyograms of the masseter, temporalis and digastric muscles were recorded in male rabbits between 7 and 20 weeks of age. Starting at 8 weeks of age the experimental animals were fed pellets significantly softer than those fed to the controls. At 20 weeks of age the myosin heavy chain (MyHC) composition and the cross-sectional area (CSA) of jaw-muscle fibres were determined using immunohistochemistry. The degree of mineralization of bone (DMB) of the mandible was determined using micro-computed tomography.

**RESULTS:** Reducing the food hardness significantly decreased the neuromuscular activity of the masseter muscle at activity levels that reflect muscle contractions during chewing. The decrease in neuromuscular activity was accompanied by a decrease in proportion and CSA of fibres expressing slow MyHC isoforms. In contrast, altered food hardness did not influence the activity or fibre properties of the temporalis and digastric muscles. The DMB of the mandible did not differ between experimental and control animals. However, it differed significantly among areas that are loaded directly or indirectly by jaw-muscle contractions in the animals that had been fed soft pellets.

**CONCLUSION:** Long-term reduction in masticatory functional load contributes to selective disuse resulting in functional and structural

adaptation of the masseter muscle. This adaptation is reflected in decreased neuromuscular activity, proportion and CSA of slow fibres. The associated alteration in the loading pattern might induce a more heterogeneous mineral distribution in mandibular bone.

Supported by an EOS research grant

## OP 57

Friday, June 18 - at 17:10

### ROLE OF PERIODONTAL LIGAMENT CELLS IN IMPAIRED TOOTH ERUPTION IN PATIENTS WITH CLEIDOCRANIAL DYSPLASIA\*\*\*

**A Jäger**, B Abou Jamra, R Abou Jamra, W Götz, S Lossdörfer, Department of Orthodontics, University of Bonn, Germany

**AIM:** Patients with cleidocranial dysplasia (CCD), which is caused by mutations in the runt related transcription factor 2 (RUNX2) gene, are characterized by anomalies of the clavicles, thorax, spine, pelvis and extremities, and by disturbances in skull and tooth development. Of orthodontic relevance are multiple supernumerary teeth associated with delayed tooth eruption. The present investigation was based on the hypothesis that an altered phenotypic expression of periodontal ligament (PDL) cells from CCD patients and a reduced ability of those cells to support the differentiation of bone resorbing osteoclasts might contribute to delayed tooth eruption.

**MATERIALS AND METHOD:** PDL cells from healthy donors and from two patients with molecular genetically diagnosed CCD were characterized for the basal and induced expression of osteoblastic marker genes, and the physiological relevance of the findings for the differentiation of osteoclasts was examined in a co-culture model of PDL cells and osteoclastic precursors.

**RESULTS:** Both CCD patients displayed missense mutations of the RUNX2 gene. *In vitro* experiments revealed an unaltered expression of RUNX2 mRNA as opposed to a reduced basal expression of mRNA for osterix, alkaline phosphatase, osteocalcin and the key regulatory gene for bone remodelling, RANKL, (receptor activator of NF- $\kappa$ B ligand). Furthermore, those factors were less inducible by stimulation of the cultures with 1 $\alpha$ ,25(OH) $_2$ D $_3$  as compared with the control cells from healthy donors. In the co-culture experiment, PDL cells from the CCD patients showed a reduced capacity to induce the differentiation of active osteoclasts.

**CONCLUSION:** PDL cells from CCD patients express a less differentiated osteoblastic phenotype resulting in an impaired ability to support osteoclastogenesis that might, in part, account for the delayed tooth eruption that can be observed clinically.

## OP 58

Friday, June 18 - at 17:20

### ASSOCIATION BETWEEN CONDYLAR MORPHOLOGY AND INFLAMMATION IN EXPERIMENTAL TEMPOROMANDIBULAR JOINT ARTHRITIS

**K D Kristensen**<sup>1</sup>, P Stoustrup<sup>1</sup>, A Küsel<sup>1</sup>, T K Pedersen<sup>1</sup>, T Herlin<sup>2</sup>,  
<sup>1</sup>Department of Orthodontics, Aarhus University and <sup>2</sup>Department of Pediatrics, Aarhus University Hospital, Denmark

**AIM:** In juvenile idiopathic arthritis involvement of the temporomandibular joints (TMJs) is often associated with severe mandibular growth deviations. The relationship between condylar growth deviations, inflammation severity, the micro-architectural composition, and bone quality has not previously been investigated. The aim of this study was to determine the effects of antigen-induced arthritis on the bony structures in rabbit mandibular condylar development, in particular the morphological changes and bone micro-architecture.

**MATERIALS AND METHOD:** Juvenile rabbits with ovalbumin-induced TMJ arthritis treated with intra-articular saline, intra-articular etanercept (an anti-TNF- $\alpha$  drug) or subcutaneous etanercept. One TMJ from each animal was scanned using micro-computed tomography and structural parameters were calculated. Three-dimensional reconstructions of the mandibular condyle were scored blindly by two independent observers as 'normal' or 'abnormal'. The TMJs were stratified for condylar morphology and evaluated against data on inflammation, trabecular structural parameters, and overall mandibular growth. Mineral apposition rate was measured using fluorochrome labelling.

**RESULTS:** Abnormal morphology was seen in 15 of 32 animals available for data analysis and was strongly related to the degree of inflammation. However, no differences in trabecular structural parameters or mineral apposition rate were seen according to treatment. Erosions were an uncommon finding. Abnormal condylar morphology was not associated with overall mandibular growth.

**CONCLUSION:** Severe inflammation in the TMJs during mandibular development is associated with morphological changes of the mandibular condyle. Morphological changes may occur because of deficient development of condylar cartilage and not because of resorptions during times of active inflammation, and is hence a temporary phenomenon.

## OP 59

Friday, June 18 - at 17:30

### THE QUESTIONABLE ROLE OF PERIODONTAL AND SUPRA-ALVEOLAR COLLAGEN FIBRES IN RELAPSE

S Henneman, R Reijers, J W Von den Hoff, **J C Maltha**, Department of Orthodontics and Oral Biology, Radboud University Nijmegen Medical Centre, Netherlands

**AIM:** Relapse is a common problem in orthodontics. Its aetiology is still unresolved, although most authors consider the periodontal and/or the supra-alveolar fibres responsible. Consequently, these fibres should persist in a stretched condition during orthodontic treatment. However, the question whether this is really the case is still unanswered. Therefore, the aim of the present study was to analyze the turnover rate of collagen fibres in different supra-alveolar and periodontal regions.

**MATERIALS AND METHOD:** Thirty male rats, 2 weeks old, received an intraperitoneal injection of 3H proline 3 times a week for 3 weeks to label all collagen. They were sacrificed at 0, 1, 4, 8, 15, 22, 29, 36, 57, 78, and 113 days after the last injection, and autoradiographs were prepared. Grains were counted in 19 predefined areas in the periodontal ligament and supra-alveolar structures to determine collagen turnover. The grain density was plotted against time, and half-lives were calculated with 95 per cent confidence intervals (CI). Multivariate analyses were performed to compare the half-lives in the different areas.

**RESULTS:** The half-life values were not different for the mesial and distal sides. However, significant differences were found along the roots. The half-lives were shortest at the top of the interradicular septum (mean = 2.4 days, 95% CI = 2.1-2.6), intermediate halfway along the roots (mean = 4.6 days, 95% CI = 4.3-4.8), and longest in the cervical and supra-alveolar region (mean = 9.6 days, 95% CI = 8.1-11.9).

**CONCLUSION:** The results point towards a more stable fibre system in the cervical and supra-alveolar regions. However, with a half-life of about 10 days, approximately 99 per cent of all fibres are remodelled within two months. This makes it very unlikely that stretched fibres persist throughout orthodontic therapy, indicating that probably other factors are involved in the aetiology of relapse.

## OP 60

Friday, June 18 - at 17:40

### ASSOCIATED PAX9 AND FIBROBLAST GROWTH FACTOR RECEPTOR 4 VARIANTS IN CLEFT LIP AND PALATE PATIENTS WITH OLIGODONTIA

O Matern<sup>1,2</sup>, B Grollemund<sup>1,2</sup>, P Tschill<sup>1,2</sup>, W Bacon<sup>2</sup>, **F Perrin-Schmitt<sup>1</sup>**,  
<sup>1</sup>UMR977-INSERM, UDS - Faculty of Medicine and <sup>2</sup>University Hospital, Functional Unit of Dentofacial Orthopedics, Strasbourg, France

**AIM:** Oral clefting is a common congenital malformation (1/700 lives), often associated with disturbed tooth development, as a result of misexpression of critical molecular actors involved in craniofacial development. Combined genetic and/or environmental factors are consensually used to explain their aetiology, and the roles of several genes in these defects have been documented both by mutant animal-model phenotypes and by direct identification of causative mutations that affect candidate-genes isolated from human patients. The aim of this investigation was to determine in patients presenting with hypodontia or oligodontia within a NSCLP cohort, the existence of specific allelic variant or mutant in selected genes.

**SUBJECTS AND METHOD:** The entire regional sample cohort comprised 50 cleft individuals and 50 unrelated controls. For each patient genomic DNA was extracted either from blood samples or from tissues obtained during surgery. Genotyping was carried out using polymerase chain reaction amplification on selected genes followed by direct sequencing.

**RESULTS:** Sequence variants in MSX1, MSX2, PAX9, DLX5, DLX6 and FGFRs were identified. In seven patients who presented with hypodontia/ oligodontia, a new association between the Ala240Pro-PAX9 and the Gly388Arg-FGFR4 variants was observed.

**CONCLUSION:** The combination of apparently minor polymorphisms in several proteins resulting in slightly imbalanced molecular pathways, might account for some causative mutations in developmental anomalies with a wide spectrum such as oral clefts.

## OP 61

Friday, June 18 - at 17:50

### ULTRASOUND EVALUATION OF TONGUE FUNCTION AND POSTURE IN CLASS III MALOCCLUSION

**J Volk**<sup>1</sup>, M Marolt Mušič<sup>2</sup>, S Melink<sup>1</sup>, M Ovsenik<sup>3</sup>, <sup>1</sup>Department of Orthodontics, University Medical Centre, <sup>2</sup>Institute of Oncology and <sup>3</sup>Department of Orthodontics, Faculty of Medicine, University of Ljubljana, Slovenia

**AIM:** To compare tongue function during swallowing and tongue posture among children with a Class III (CLIII) malocclusion and those with a normal dentition (ND) using two- and three-dimensional (3D) ultrasonography.

**SUBJECTS AND METHOD:** Swallowing cycles of 13 children with CIII (mean age:  $7 \pm 1$  years) and 14 children with ND (mean age:  $7.1 \pm 0.3$  years), were recorded three times using B- and M-mode ultrasound (US) techniques (SSA-770A with a convex transducer). The scan line was set through the middle of the tongue for the first two swallowing cycles and through the tip of the tongue in the last cycle. The duration of tongue movements of the middle third of the tongue were compared within each subphase and during the entire cycle between the CIII and the ND group, and the swallowing pattern for each child was assessed according to the action of the tip of the tongue. Voluson 730 Expert with a 3D convex transducer was used to obtain two 3D US images of the tongue in each child in both groups. Referential 3D US reconstructions were used for assessment of the tongue posture in each child.

**RESULTS:** The duration of the entire swallowing cycle ( $P < 0.001$ ) and of phase IIb ( $P < 0.01$ ) was prolonged in the CIII group. Sixty-nine per cent of the CIII children and 43 per cent of the ND children had a visceral swallowing pattern (Fisher exact test,  $P = 0.161$ ). Eighty-five per cent of the CIII children but only 36 per cent of the ND children demonstrated a tongue posture on the mouth floor according to the 3D US reconstructions ( $P < 0.001$ ).

**CONCLUSION:** There were significant differences in the duration of tongue movements between both groups of children. The tongues of the CLIII children were, to a greater extent, postured on the mouth floor compared with children with ND.

**CONCLUSION:** No conclusive evidence was found for a positive relationship between masticatory muscle volume and maximum cross-sectional area with vertical facial dimensions. However, subcutaneous tissue, especially at the chin area, may affect face height measurements and influence the accuracy of the results.



# SCIENTIFIC POSTERS



## LIST OF SCIENTIFIC POSTERS

### List of Scientific Posters

Abstract No	Title, Presentator, Country	Page
SP 001	ASSOCIATION BETWEEN ORTHODONTIC TREATMENT NEED AND MAXILLARY INCISOR TRAUMA A <b>Borzabadi-Farahani</b> , Iran	57
SP 002	NOCTURNAL ENURESIS AND RAPID MAXILLARY EXPANSION <b>F Bazargani</b> , Sweden	57
SP 003	CEPHALOMETRICS IN THE MAGNETIC FIELD*** <b>T Brandt</b> , Germany	57
SP 004	EVALUATION OF DENTAL AESTHETICS AND ORTHODONTIC TREATMENT NEED <b>E González Nieto</b> , Spain	58
SP 005	STRONTIUM DIMINISHES DELETERIOUS EFFECTS OF RUNX2 HAPLOINSUFFICIENCY IN HUMAN OSTEOBLASTS <b>P Römer</b> , Germany	58
SP 006	STRONTIUM COUNTERACTS TOOTH RESORPTION BY INHIBITING EXPRESSION OF RANKL IN PERIODONTAL LIGAMENT CELLS <b>P Römer</b> , Germany	58
SP 007	THREE-DIMENSIONAL COMPARISON OF AVERAGE FACES OF CHILDREN WITH ORAL CLEFTS <b>I Bugaighis</b> , U.K	58
SP 008	ASSESSMENT OF THREE METHODS TO ASSESS PLAQUE LEVELS AND WHITE SPOT LESIONS <b>N Pender</b> , England	59
SP 009	INFLUENCE OF THE VISUAL SYSTEM TO HORIZONTAL JAW RELATIONSHIP DETERMINATION – A PILOT STUDY <b>D Ohlendorf</b> , Germany	59
SP 010	THE IMPACT OF FORCED LEG LENGTH DISCREPANCIES ON BODY POSTURE AND JAW POSITION <b>D Ohlendorf</b> , Germany	59
SP 011	THREE-DIMENSIONAL MAXILLO-MANDIBULAR DENTAL CHANGES AFTER ORTHODONTIC TREATMENT WITH PREMOLAR EXTRACTIONS <b>E C K Wee</b> , Singapore	59
SP 012	INFLUENCE OF CORROSION OF SURFACE DEFECTS OF NICKEL TITANIUM WIRES <b>C Palazón Martínez</b> , Spain	60
SP 013	DO DIFFERENT TENSILE BOND STRENGTH STUDIES END IN THE SAME RESULTS? <b>S Hemmatpour</b> , Iran	60
SP 014	IN VITRO CORROSION BEHAVIOUR OF LINGUAL ORTHODONTIC ARCHWIRES <b>C Suárez</b> , Spain	60
SP 015	THE TEMPOROMANDIBULAR JOINTS IN PATIENTS WITH SAGITTAL MALOCCLUSIONS <b>I Skrypnik</b> , Ukraine	60
SP 016	RESVERATROL INHIBITS PERIODONTAL PATHOGENS IN VITRO <b>D J O'Connor</b> , SAR China	61
SP 017	USE OF AN ER,Cr:YSGG, WATERLASE LASER IN ORTHODONTIC PATIENTS WITH GINGIVAL OVERGROWTH <b>P Thongdee</b> , SAR China	61
SP 018	EFFECT OF DIET HARDNESS ON MANDIBULAR CONDYLAR CARTILAGE METABOLISM <b>P Pirttiniemi</b> , Finland	61
SP 019	COMPOSITE OCCLUSAL BITE BLOCKS BONDED WITHOUT LIQUID RESIN <b>A T H Tang</b> , SAR China	61
SP 020	DEVELOPMENTAL ANALYSIS OF ETHMOIDAL SINUS EMPLOYING SAGITTAL COMPUTED TOMOGRAPHIC SCANS*** <b>M Numata</b> , Japan	62
SP 021	STEM CELL DIFFERENTIATION INTO ANGIOGENESIS LINEAGE <b>A Almoqubel</b> , SAR China	62
SP 022	ELECTROMYOGRAPHIC AGE VALUES OF STOMATOGNATHIC MUSCLES <b>N Pankratova</b> , Russia	62
SP 023	SIGNIFICANCE OF AGE ON LOWER JAW MOTION – A THREE DIMENSIONAL STUDY <b>N Pankratova</b> , Russia	62
SP 024	PULPAL REACTION TO LOADING APPLIED TO UPPER TEETH IN PATIENTS WITH CLASS II MALOCCLUSIONS <b>M Szumielewicz</b> , Poland	63

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 025	CLEFT LIP AND PALATE PATIENTS AND HYPODONTIA – A SINGLE NUCLEOTIDE POLYMORPHISM STUDY <b>B Nemes</b> , Hungary	63
SP 026	ENZYMATIC ACTIVITY IN GINGIVAL CREVICULAR FLUID WITH SELF-LIGATING BRACKETS <b>A Caprioglio</b> , Italy	63
SP 027	COMPARISON OF CEPHALOMETRIC MEASUREMENTS: COMPUTERIZED VERSUS CONVENTIONAL ANALYSIS <b>I N Zetu</b> , Romania	64
SP 028	A COMPARATIVE STUDY ON THE ALIGNMENT EFFICIENCY OF A LINGUAL SYSTEM <b>S J Zhu</b> , SAR China	64
SP 029	ROLE OF FLUORIDE AND CARBONATED DRINKS IN THE CORROSION OF NICKEL TITANIUM WIRES <b>C Palazón Martínez</b> , Spain	64
SP 030	FINITE ELEMENT ANALYSIS OF ORTHODONTIC IMPLANTS WITH DIFFERENT DESIGNS <b>C Szuhaneck</b> , Romania	64
SP 031	EVALUATION OF CEPHALOMETRIC MEASUREMENTS OF PATIENTS AFTER ORTHOGNATHIC SURGERY <b>M Janega</b> , Czech Republic	65
SP 032	INFLUENCE OF ADHESIVE SURFACE ROUGHNESS ON STREPTOCOCCAL ADHESION FORCES <b>Y Ren</b> , Netherlands	65
SP 033	MATHEMATICAL APPRAISAL OF ARCH FORM IN SUBJECTS WITH CROWDING TREATED NON -EXTRACTION <b>B Vande Vannet</b> , Belgium	65
SP 034	RATE OF DISTALIZATION IN CLASS II SUBJECTS USING HIGH PULL HEADGEAR <b>W Stós</b> , Poland	65
SP 035	CERVICAL VERTEBRAL MATURATION STAGES AND CALCIFICATION OF THE MANDIBULAR SECOND MOLAR <b>M Bachleda-Księdzularz</b> , Poland	66
SP 036	MICRODAMAGE IN THE ALVEOLAR PROCESS AFTER ORTHODONTIC TOOTH MOVEMENT <b>T M Präger</b> , Germany	66
SP 037	EFFECT OF MINIMAL OSTEOPERFORATIONS ON BONE REMODELLING AND TOOTH MOVEMENT I <b>Chartres</b> , Germany,	66
SP 038	HUMAN FOETUS CADAVERS WITH A CLEFT LIP AND PALATE IN THE THIRD TRIMESTER: CASE REPORT <b>E Esenlik</b> , Turkey	66
SP 039	STABILITY OF IMMEDIATELY LOADED MINISCREW IMPLANTS ACCORDING TO FORCE LEVELS*** <b>Y-M Jo</b> , Germany	67
SP 040	MECHANICAL PROPERTIES OF NICKEL-TITANIUM ARCHWIRES AFTER USE IN THE ORAL ENVIRONMENT <b>M Sarul</b> , Poland	67
SP 041	THERMOCYCLING EFFECTS ON THE BOND STRENGTH OF BRACKETS BONDED WITH DIFFERENT LIGHT SOURCES <b>E Çerekja</b> , Turkey	67
SP 042	EFFECT OF OESTROGEN AND LOADING TO THE AMOUNT OF OESTROGEN RECEPTOR ALPHA AND MATRIX METALLOPROTEINASE-8 IN RAT CONDYLAR CARTILAGE <b>M Orajärvi</b> , Finland	67
SP 043	INVESTIGATION OF THE MOST PREFERRED MOUTH SHAPE DURING SMILING IN JAPANESE. PART I*** <b>J Funaki</b> , Japan	68
SP 044	INVESTIGATION OF THE MOST PREFERRED MOUTH SHAPE DURING SMILING IN JAPANESE. PART II CHANGES AFTER ORTHODONTIC TREATMENT*** <b>J Funaki</b> , Japan	68
SP 045	THE EFFECTS OF A FACEMASK/BITE BLOCK THERAPY WITH OR WITHOUT RAPID PALATAL EXPANSION <b>C Pavoni</b> , Italy	68
SP 046	WHICH FACIAL PROFILES MOSTLY DISCRIMINATE PERCEPTION OF MALE AND FEMALE ADOLESCENTS? <b>S Špalj</b> , Croatia	69
SP 047	SKELETAL CHANGES INDUCED BY RAPID MAXILLARY EXPANSION ASSESSED WITH SUTURAL DENSITY ANALYSIS ON LOW-DOSE COMPUTED TOMOGRAPHY <b>L Franchi</b> , Italy	69
SP 048	TREATMENT PLANNING FOR MAXILLARY CANINE DISPLACEMENT BASED ON INDIVIDUAL SKELETAL MATURITY <b>V Giuntini</b> , Italy	69
SP 049	A DIAGNOSTIC PERFORMANCE STUDY ON OVERJET IN CLASS II MALOCCLUSION SUBJECTS <b>A Vangelisti</b> , Italy	70
SP 050	NON-CONVENTIONAL BRACKETS VERSUS LIGATURES FOR THE ALIGNMENT OF BUCCALLY DISPLACED TEETH <b>A Vangelisti</b> , Italy	70
SP 051	THREE-DIMENSIONAL LOCALIZATION OF IMPACTED MAXILLARY CANINES AND EFFECTS ON ADJACENT INCISORS <b>K Mina</b> , Iran	70
SP 052	CRANIOFACIAL MORPHOLOGY OF CHINESE ADULTS WITH NORMAL OCCLUSION AND SKELETAL DIVERGENCE <b>Y Ren</b> , China	70
SP 053	RELATIONSHIP BETWEEN UPPER AIRWAY WIDTH AND FACIAL GROWTH CHANGES DURING ORTHODONTIC TREATMENT OF GROWING CHILDREN <b>Y-H Park</b> , Korea	71
SP 054	TWO SCORING SYSTEMS FOR TREATMENT EVALUATION OF BILATERAL CLEFT LIP AND PALATE PATIENTS <b>T Bartzela</b> , Switzerland	71
SP 055	ADULT CLASS II DIVISION 1 HERBST TREATMENT – IS IT STABLE? <b>N Bock</b> , Germany	71

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 056	EFFECT OF AUTOTRANSPLANTATION ON BONE REGENERATION IN ORTHODONTIC PATIENTS <b>E Hamza</b> , SAR China	71
SP 057	ANTERIOR DENTOALVEOLAR MORPHOLOGY IN RELATION TO NATURAL HEAD POSTURE <b>H Gokalp</b> , Turkey	72
SP 058	RELATIONSHIP BETWEEN ROOT RESORPTION AND DENTAL ANOMALIES <b>K Van Parys</b> , Netherlands	72
SP 059	FIXED FUNCTIONAL TREATMENT VERSUS PREMOLAR EXTRACTIONS IN YOUNG ADULTS <b>I Ilhan</b> , Turkey	72
SP 060	A PROTOCOL FOR IMPROVED STABILITY WITH HERBST APPLIANCE TREATMENT <b>D F Austin</b> , SAR China	72
SP 061	EDENTULOUS RIDGE ALTERATIONS FOLLOWING ORTHODONTIC TOOTH MOVEMENT <b>B Lindskog Stokland</b> , Sweden	73
SP 062	THE DYNAMIC FOURTH DIMENSION IN IMAGING AND ITS RELATION TO CRANIAL INCLINATIONS <b>S Usumez</b> , Turkey	73
SP 063	A NEW DIAGNOSTIC APPROACH TO ORTHODONTIC PATIENTS <b>L Sonnesen</b> , Denmark	73
SP 064	SURGERY-FIRST APPROACH IN SKELETAL CLASS III MALOCCLUSION TREATED WITH TWO-JAW SURGERY <b>S-H Baek</b> , South Korea	73
SP 065	CEPHALOMETRIC EVALUATION OF WITS ANALYSIS <b>I Buck</b> , Germany	74
SP 066	SURGICAL TREATMENT PLANNING OF PATIENTS WITH CLASS II MALOCCLUSIONS <b>K Kurakin</b> , Russia	74
SP 067	THE BIDIMENSIONAL TECHNIQUE EN MASSE RETRACTION USING SELF-LIGATING APPLIANCES <b>L Iacopini</b> , Italy	74
SP 068	CEPHALOMETRIC DECISION: WHETHER OR NOT TO EXTRACT IN ORTHODONTIC TREATMENT PLANNING <b>P Elmi</b> , Netherlands	75
SP 069	INFLUENCE OF FUNCTIONAL APPLIANCES ON POSTURAL BALANCE IN CHILDREN WITH MALOCCLUSION <b>I Rubleva</b> , Russia	75
SP 070	THE EFFECT OF LIP ADHESION ON EARLY MAXILLARY GROWTH IN BILATERAL CLEFT PATIENTS <b>G De Pauw</b> , Belgium	75
SP 071	FINITE ELEMENT ANALYSIS OF STRESS AND STRAIN DURING USE OF RAPID PALATAL EXPANDERS <b>D C Bratu</b> , Romania	75
SP 072	SKELETAL MATURATION IN CHILDREN <b>M Durka-Zajac</b> , Poland	76
SP 073	EPIDEMIOLOGICAL ASPECTS OF MISSING TEETH IN A ROMANIAN POPULATION <b>I N Zetu</b> , Romania	76
SP 074	CONE BEAM COMPUTED TOMOGRAPHY FOR ROUTINE TREATMENT PLANNING IN ORTHODONTICS <b>M Nitka</b> , Germany	76
SP 075	VALIDATION OF CONVENTIONAL RADIOGRAPHS EXAMINATIONS TO ASSESS THE MAGNITUDE OF EXTERNAL RESORPTION <b>I Marek</b> , Czech Republic	76
SP 076	A NEW CONCEPT FOR ANB ANGLE BASED ON FLOATING NORMS <b>K Faltina</b> , Brazil	77
SP 077	EFFECTS OF RAPID MAXILLARY EXPANSION ON CRANIOFACIAL SUTURES AND SPHENO-OCCIPITAL SYNCHONDROSIS – A COMPUTERIZED TOMOGRAPHIC STUDY <b>R Leonardi</b> , Italy	77
SP 078	CEPHALOMETRIC VALUES FOR A MOROCCAN POPULATION COMPARED WITH CAUCASIAN NORMS*** <b>K Arhouni</b> , Belgium	77
SP 079	ANALYSIS OF OCCLUSAL PLANE POSITION IN PATIENTS TREATED WITH DISTAL BITE CORRECTORS <b>M Markova</b> , Russia	77
SP 080	DIAGNOSTIC INVESTIGATIONS OF PATIENTS WITH OLIGODONTIA AND ECTODERMAL DYSPLASIA <b>M Zadurska</b> , Poland	78
SP 081	UNILATERAL POSTERIOR CROSSBITE CORRECTION IN THE MIXED DENTITION – A 3-YEAR FOLLOW-UP <b>S Petrén</b> , Sweden	78
SP 082	ESTIMATION OF DENTAL ALVEOLUS MORPHOLOGY AS A PROGNOSTIC INDICATOR OF GINGIVAL RECESSION <b>M Dominiak</b> , Germany	78
SP 083	AESTHETICS BEFORE AND AFTER ORTHODONTIC TREATMENT <b>A Muntean</b> , Romania	78
SP 084	A COMPARATIVE STUDY OF LOWER THIRD MOLAR SPACE IN SKELETAL CLASS III AND CLASS II ANOMALIES <b>M Popescu</b> , Romania	79
SP 085	ORTHODONTIC TREATMENT OF PATIENTS WITH CROWDING USING SELF-LIGATING BRACKETS <b>O Emeljanova</b> , Russia	79
SP 086	PREDICTION OF THE MESIODISTAL WIDTH OF PERMANENT TEETH <b>T Yavorovskaya</b> , Russia	79

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 087	HYOID BONE POSITION, MANDIBULAR GROWTH DIRECTION AND HEAD POSTURE IN MOUTH BREATHERS <b>G Curcieva-Cuckova</b> , Former Yugoslav Republic of Macedonia	79
SP 088	CONE BEAM COMPUTED TOMOGRAPHY OF THE TEMPOROMANDIBULAR JOINT IN THE SURGICAL TREATMENT OF CLASS III ANOMALIES <b>M Popescu</b> , Romania	80
SP 089	ASSESSMENT OF CROWDING WITH A SELF-LIGATING SYSTEM IN NON-EXTRACTION THERAPY <b>B Vande Vannet</b> , Austria	80
SP 090	VACUUM RETAINER TECHNIQUE IN THE TREATMENT OF ANTERIOR CROSSBITE <b>I Apruzzese</b> , Italy	80
SP 091	VALIDITY OF UPPER AIRWAY ASSESSMENT IN CHILDREN <b>K Piriälä-Parkkinen</b> , Finland	80
SP 092	AESTHETIC CHANGES OF THE PROFILE IN EXTRACTION AND NON-EXTRACTION CASES <b>A Muntean</b> , Romania	81
SP 093	ROOT DAMAGE AFTER SCREW INSERTION FOR ORTHODONTIC MINIPLATE PLACEMENT IN DOGS <b>M Cornelis</b> , Belgium	81
SP 094	THE ROLE OF MODERN THREE-DIMENSIONAL IMAGING IN ORTHODONTIC DIAGNOSIS AND TREATMENT PLANNING <b>G Rossini</b> , Italy	81
SP 095	EFFECTS OF SURGICALLY ASSISTED RAPID MAXILLARY EXPANSION ON CRANIOFACIAL MORPHOLOGY <b>A Y Gungor</b> , Turkey	81
SP 096	ORTHODONTIC-PERIODONTIC INTERRELATIONSHIP: A SYSTEMATIC REVIEW <b>P Christou</b> , Greece	82
SP 097	EARLY TREATMENT OF UNILATERAL POSTERIOR CROSSBITE – AN ELECTROGNATHOGRAPHIC EVALUATION <b>E Sever</b> , Slovenia	82
SP 098	ASSESSMENT OF CHEWING CYCLE MORPHOLOGY – A LONGITUDINAL STUDY IN CHILDREN FROM 5 TO 7 YEARS OF AGE <b>E Vehovec</b> , Slovenia	82
SP 099	INCOMPETENT LIP SEAL AND TONGUE POSTURE – IS THERE A CORRELATION? <b>V Božnik</b> , Slovenia	82
SP 100	THREE-DIMENSIONAL CHARACTERISTICS OF FACIAL AND JAW MORPHOLOGY IN THE EARLY MIXED DENTITION <b>D Malinar</b> , Wales	83
SP 101	ASSESSMENT OF THE OCCLUSAL CANT IN THE FRONTAL PLANE <b>O Popoviciu</b> , Romania	83
SP 102	FACIAL MORPHOLOGY CHANGES IN THE MIXED DENTITION PERIOD FROM 8 TO 9 YEARS OF AGE – A THREE-DIMENSIONAL EVALUATION <b>M Verdenik</b> , Wales	83
SP 103	BUCCALLY BLOCKED OUT CANINES – CLINICAL EVIDENCE OF THE THREE-BRACKET RELATIONSHIP <b>M Svejda</b> , Austria	83
SP 104	THE PREVALENCE OF WHITE SPOT LESIONS AFTER ORTHODONTIC FIXED APPLIANCE THERAPY <b>M Drevenšek</b> , Slovenia	84
SP 105	THREE-DIMENSIONAL CEPHALOMETRY FOR DIAGNOSIS AND TREATMENT PLANNING OF IMPACTED CANINES <b>I M Dagsuyu</b> , Turkey	85
SP 106	CORRELATION BETWEEN TOOTH SIZE AND JAW LENGTH IN CLASS III AND CLASS I MALOCCLUSION SUBJECTS <b>H R Fattahi</b> , Iran	85
SP 107	A COMPARISON OF FACEMASK TREATMENT EFFECTS ACCORDING TO CERVICAL VERTEBRAE MATURATION <b>B Yang</b> , Korea	85
SP 108	EFFECTS OF A FACEMASK COMBINED WITH A CHINCAP ON MAXILLARY GROWTH IN SKELETAL CLASS III SUBJECTS <b>H N İřcan</b> , Turkey	85
SP 109	EARLY TREATMENT OF PSEUDO CLASS III MALOCCLUSION – A 10-YEAR FOLLOW-UP STUDY <b>I M Anderson</b> , SAR China	86
SP 110	PREDICTION OF THE OUTCOME OF TREATMENT OF CLASS III MALOCCLUSION – A SYSTEMATIC REVIEW <b>P Fudalej</b> , Poland	86
SP 111	EFFECTS OF PALATAL PLANE ROTATION ON SOFT TISSUE CHANGES IN DOUBLE JAW SURGERY <b>Y Kim</b> , Korea	86
SP 112	MANAGEMENT OF SKELETAL CLASS III PATIENTS <b>V Tarmure</b> , Romania	86
SP 113	EFFECTS OF A FACEMASK COMBINED WITH A CHINCAP ON MAXILLARY GROWTH IN SKELETAL CLASS III SUBJECTS <b>H N İřcan</b> , Turkey	87
SP 114	CEPHALOMETRIC ASSESSMENT OF CHANGES IN THE FACIAL SKELETON IN CHILDREN WITH MANDIBULAR PROGNATHISM <b>M Pospiesznyńska</b> , Poland	87
SP 115	TONGUE VOLUME AND POSTERIOR AIRWAY SPACE CHANGES IN CLASS III TWO JAW ORTHOGNATHIC SURGERY <b>O Uslu Akçam</b> , Turkey	87
SP 116	STRUCTURAL CHANGE OF THE HYOID BONE AND UPPER AIRWAY AFTER ORTHOGNATHIC SURGERY USING THREE-DIMENSIONAL COMPUTED TOMOGRAPHY <b>H-S Yu</b> , Korea	87
SP 117	LONG-TERM OBSERVATION OF MASSETER MUSCLE CHANGES IN FACIAL ASYMMETRY <b>H-S Yu</b> , Korea	88

Abstract No	Title, Presentator, Country	Page
SP 118	COMPARISON OF FACEMASK OR REVERSE CHIN CUP THERAPY IN TREATING CLASS III MALOCCLUSIONS <b>A Jamilian</b> , Iran	88
SP 119	CERVICAL VERTEBRA ANATOMY IN PATIENTS WITH NORMAL OCCLUSION AND CLASS III MALOCCLUSION <b>M Pop Stefanova-Trposka</b> , Former Yugoslav Republic of Macedonia	88
SP 120	AN ETHNIC BACKGROUND-BASED CEPHALOMETRIC STUDY IN HUNGARIAN AND JAPANESE CLASS III SUBJECTS <b>I Csiki</b> , Japan	88
SP 121	EFFECTS OF FACEMASK TREATMENT ON SAGITTAL PHARYNGEAL DIMENSIONS IN CLASS III SUBJECTS <b>R Lione</b> , Italy	89
SP 122	PSYCHOLOGICAL CHARACTERISTICS OF PATIENTS WITH SKELETAL CLASS III MALOCCLUSIONS <b>A Kovalenko</b> , Russia	89
SP 123	MANDIBULAR FEATURES OF CLASS III PATIENTS WITH IMPLICATIONS ON TEMPOROMANDIBULAR DISORDERS <b>M-A Bencze</b> , Romania	89
SP 124	TOOTH-BORNE DISTRACTOR APPLICATION FOR ANTERIOR MAXILLARY POSITIONING IN CLEFT AND PALATE PATIENTS <b>A Showkatbakhsh</b> , Iran	90
SP 125	EFFECTIVENESS OF COMBINED SURGICAL-ORTHODONTIC TREATMENT: CEPHALOMETRIC ANALYSIS <b>J Szyper-Szczurowska</b> , Poland	90
SP 126	CORRELATION OF INCISOR INCLINATION AND THE SAGITTAL POSITION OF THE JAWS IN CLASS III SUBJECTS <b>M Stosic</b> , Serbia	90
SP 127	FACIAL SOFT TISSUE PROFILE CHANGES FOLLOWING BILATERAL SPLIT SAGITTAL OSTEOTOMY AND SUBCONDYLAR MANDIBULAR SETBACK <b>S Torkan</b> , Iran	90
SP 128	CONSIDERATIONS ON THE THERAPEUTIC INTERVENTION IN MIXED DENTITION CLASS III CASES <b>E Ionescu</b> , Romania	90
SP 129	CORRELATION BETWEEN MANDIBULAR AND CRANIAL BASE FEATURES IN SKELETAL CLASS III ANOMALIES <b>R Grigore</b> , Romania	91
SP 130	DENTAL EFFECTS OF MAXILLARY PROTRACTION WITH A FACEMASK, WITH OR WITHOUT PALATAL EXPANSION <b>A Cesta</b> , Italy	91
SP 131	MAXILLARY PROTRACTION WITH A FACEMASK: EFFECTS ON SAGITTAL AIRWAY DIMENSIONS <b>T Avventi</b> , Italy	91
SP 132	MAXILLARY PROTRACTION WITH OR WITHOUT PALATAL EXPANSION: CEPHALOMETRIC SAGITTAL EFFECTS <b>C Berto</b> , Italy	91
SP 133	VERTICAL SKELETAL CHANGES WITH FACE MASK THERAPY, WITH OR WITHOUT PALATAL EXPANSION <b>C Berto</b> , Italy	92
SP 134	A NOVEL METHOD OF ANALYSIS OF CLASS III PATIENTS USING THREE-DIMENSIONAL SOFT TISSUE ANALYSIS <b>M Božič</b> , Slovenia	92
SP 135	PRIMARY VERSUS PERMANENT DENTITION ANCHORAGE IN EARLY CLASS III ORTHOPAEDIC TREATMENT <b>E Frumusa</b> , Italy	92
SP 136	EFFECTS OF CHIN CUP THERAPY ON SYMPHYSIS MORPHOLOGY <b>G Görgülü</b> , Turkey	92
SP 137	CONDYLAR TRACINGS IN CLASS III MALOCCLUSION SUBJECTS BEFORE AND AFTER ORTHOGNATHIC SURGERY <b>G Frongia</b> , Italy	93
SP 138	REVERSE-SEQUENCING CHEWING PATTERNS BEFORE AND AFTER CLASS III ORTHOGNATHIC SURGERY <b>M G Piacino</b> , Italy	93
SP 139	A COMPARATIVE STUDY OF CLASS III TREATMENT WITH A TONGUE APPLIANCE AND REVERSE PULL CHIN CAP <b>A Showkatbakhsh</b> , Iran	93
SP 140	NATURAL HEAD POSITION IN CLASS III MALOCCLUSION SUBJECTS <b>Z Hedayati</b> , Iran	93
SP 141	FACIAL MORPHOLOGY OF CLASS III PATIENTS IN THE MIXED DENTITION USING THREE-DIMENSIONAL IMAGING <b>B Krneta</b> , Slovenia	94
SP 142	ASSESSMENT OF THE PHARYNGEAL AIRWAY IN SURGICALLY CORRECTED CLASS III PATIENTS <b>U Toygar Memikoğlu</b> , Turkey	94
SP 143	THE PREDICTION OF FUTURE ORTHOGNATHIC SURGERY NEED IN CHILDREN <b>G Schuster</b> , Germany	94
SP 144	HYPERDIVERGENT CLASS III AND MAXILLARY HYPOPLASIA: IS FACEMASK THERAPY NECESSARY? <b>P Lucchi</b> , Italy	94
SP 145	COMPARISON OF CRANIAL BASE ANGLE BETWEEN CLASS II AND CLASS III MALOCCLUSION SUBJECTS <b>Z Hedayati</b> , Iran	95
SP 146	THE CRANIAL BASE AND CRANIOFACIAL COMPLEX IN CLASS III MALOCCLUSION: A CEPHALOMETRIC STUDY*** <b>J Blaga</b> , Romania	95
SP 147	PREVALENCE OF DIFFERENT CLASS III MALOCCLUSIONS AND THE TIMING OF TREATMENT <b>A Toth Pirnovar</b> , Slovenia	95
SP 148	LONG-TERM STABILITY OF SKELETAL CLASS III PATIENTS TREATED WITH SPLINTS AND CLASS III ELASTICS <b>L Perillo</b> , Italy	95

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 149	LINEAR VARIABLES AND THE PRESENCE OF THIRD MOLARS IN CLASS III AND CLASS II DIVISION 2 PATIENTS <b>B Mady Maričić</b> , Croatia	95
SP 150	PREVALENCE OF CLASS III MALOCCLUSIONS – IS THERE SUFFICIENT EVIDENCE? <b>M Orellana</b> , USA	96
SP 151	PREVALENCE OF FUNCTIONAL DISORDERS IN PATIENTS WITH CLASS III MALOCCLUSIONS <b>M Mikac Cankar</b> , Slovenia	97
SP 152	LONG-TERM OUTCOME AND STABILITY IN TREATED CLASS I EXTRACTION AND NON-EXTRACTION PATIENTS <b>M R Gjetthamme</b> , Norway	97
SP 153	THE QUERCETIN EFFECT ON SOX9 AND COLLAGEN II EXPRESSIONS IN SPHENO-OCCIPITAL SYNCHONDROSES <b>A Veerawattanatigul</b> , China	97
SP 154	EFFECT OF QUERCETIN ON EXPRESSION OF BMP2 AND PTHRP IN MICE SPHENO-OCCIPITAL SYNCHONDROSES <b>K Wangwarunyoo, R Wong, B Rabie</b> , China	97
SP 155	OCCLUSAL FEATURES OF CHILDREN WITH ADENOID HYPERTROPHY IN COMPARISON WITH A CONTROL GROUP <b>K Lubinska</b> , Poland	98
SP 156	ADVANCED MORPHOMETRICS: REGISTRATION OF THREE-DIMENSIONAL FACIAL IMAGES <b>A I Zhurov, A M Toma, S Richmond</b> , Wales	98
SP 157	RADIOGRAPHIC STUDY OF SUPERNUMERARY TEETH CAUSING NON-ERUPTION OF PERMANENT INCISORS <b>M T Chew</b> , Singapore	98
SP 158	ACCURACY OF MECHANICAL TORQUE LIMITING GAUGES FOR MINI-SCREW PLACEMENT <b>M Schätzle</b> , SAR China	98
SP 159	CHANGES IN SKULL PARAMETERS IN CLASS II DIVISION 2 SUBJECTS <b>M Dushenkova</b> , Russia	99
SP 160	CORRELATION BETWEEN ANTEGONIAL NOTCH DEPTH WITH SAGITTAL AND VERTICAL PARAMETERS <b>H R Fattahi</b> , Iran	99
SP 161	EXTRAORAL JUDGEMENT OF SKELETAL CLASS <b>M Miguez Contreras</b> , Spain	99
SP 162	LONG-TERM SKELETAL AND SOFT TISSUE OUTCOMES OF BIONATOR THERAPY IN CLASS III MALOCCLUSION SUBJECTS <b>V Giuntini</b> , Brazil	99
SP 163	A SYSTEMATIC REVIEW OF INTERCEPTIVE TREATMENT OF PALATALLY DISPLACED MAXILLARY CANINES <b>J Naumova</b> , Sweden	100
SP 164	VEGETABLE ALTERNATIVES TO FIGHT AGAINST DENTAL MICROBIALS <b>I Ionescu</b> , Romania	100
SP 165	TESTING THE FIDELITY OF IMPRESSION MATERIALS USED IN ORTHODONTICS <b>G Tanase</b> , Romania	100
SP 166	ERRORLESS PHOTOMETRY – IS IT POSSIBLE? <b>Z Lendengolts</b> , Russia	100
SP 167	HUMAN MASTICATORY MUSCLE DIMENSIONS AND VERTICAL FACE HEIGHT - A MRI STUDY <b>P Y E Sham</b> , Singapore	101
SP 168	DEGREE OF MANDIBULAR MOLAR ROOT DEVELOPMENT IN RELATION TO CRANIOFACIAL GROWTH TYPE <b>J Andjelic</b> , Montenegro	101
SP 169	ELECTRON PROBE MICROANALYSIS OF TEN REPRESENTATIVE NICKEL TITANIUM ARCHWIRES <b>Y Bolender</b> , France	101
SP 170	DENTOSKELETAL AND SOFT TISSUE PROFILE CHANGES IN CLASS II HIGH MANDIBULAR ANGLE PREADOLESCENTS <b>T Boskovic-Brkanovic</b> , Montenegro	101
SP 171	COMPARATIVE STUDY OF BOND STRENGTH OF NEW AND RECONDITIONED BRACKETS AND ASSESSMENT OF RESIDUAL ADHESIVE <b>B Wendl</b> , Austria	102
SP 172	HOW WELL DO ORTHODONTISTS DIAGNOSE AND RECORD TEMPOROMANDIBULAR DISORDERS? <b>S Visram</b> , England	102
SP 173	SKELETAL MATURATION IN CHILDREN WITH A CLEFT LIP AND PALATE <b>M O Akcam</b> , Turkey	102
SP 174	CLEFT LIP/PALATE IN MAYO, IRELAND 1999–2007 <b>C M McNamara</b> , England	102
SP 175	EVALUATION OF ORAL HYGIENE IN PATIENTS TREATED WITH REMOVABLE APPLIANCES <b>P Pietrzak</b> , Poland	103
SP 176	CEPHALOMETRIC EVALUATION OF PATIENTS WITH OBSTRUCTIVE SLEEP APNOEA <b>H Turkkahraman</b> , Turkey	103
SP 177	USING ULTRAVIOLET LIGHT TO ARTIFICIALLY AGE ELASTOMERIC CHAIN <b>D Bister</b> , England	103
SP 178	IN VITRO PEAK INSERTION TORQUE VALUES OF SIX COMMERCIALY AVAILABLE MINI-IMPLANTS <b>C Z-Y Whang</b> , England	103
SP 179	CLEFT LIP/PALATE IN MAYO, IRELAND 1999–2007 <b>C M McNamara</b> , England	104

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 180	META-ANALYSIS OF STUDIES ON PAEDIATRIC CRANIOMANDIBULAR DISORDERS <b>A Pilotto</b> , Italy	104
SP 181	EFFECTS OF CONTINUOUS AND INTERMITTENT CONTROLLED FORCES ON CANINE DISTALIZATION <b>S Usumez</b> , Turkey	104
SP 182	STRUCTURAL ADAPTATION OF RABBIT JAW MUSCLES TO ALTERED MASTICATORY FUNCTIONAL LOAD <b>M Vreeke</b> , Netherlands	105
SP 183	CHARACTERIZATION OF AS-RECEIVED AND IN VIVO ORTHODONTIC STAINLESS STEEL BRACKETS <b>S Hoste</b> , Belgium	105
SP 184	EVALUATION OF SKELETAL CHANGES INDUCED BY SURGICALLY ASSISTED RAPID PALATAL EXPANSION <b>G Iodice</b> , Italy	105
SP 185	ANTERIOR MAXILLARY AND MANDIBULAR INTERRADICULAR DIMENSIONS AND CORTICAL BONE THICKNESS <b>M Fayed</b> , Egypt	105
SP 186	WHITE SPOT LESIONS AFTER ORTHODONTIC TREATMENT – THREE MONTHS USE OF CASEIN-PHOSPHOPEPTIDE AMORPHOUS-CALCIUM-PHOSPHATE FLUORIDE PASTE <b>M W Beerens</b> , Netherlands	106
SP 187	EFFECT OF CASEIN-PHOSPHOPEPTIDE AMORPHOUS-CALCIUM-PHOSPHATE FLUORIDE PASTE ON PLAQUE COMPOSITION AFTER ORTHODONTIC TREATMENT <b>N C W van der Kaaij</b> , Netherlands	106
SP 188	RE-BOND STRENGTH OF THE BONDED LINGUAL WIRE RETAINER <b>K van Westing</b> , Netherlands	106
SP 189	CEPHALOMETRIC ASSESSMENT OF RAPID MAXILLARY EXPANSION IN CLASS II DIVISION 1 MALOCCLUSION SUBJECTS*** <b>M Pacurar</b> , Romania,	106
SP 190	POSTERIOR CROSSBITE AND SUCKING HABITS IN CHILDREN FOLLOWED BETWEEN 3 AND 7 YEARS OF AGE <b>L Dimberg</b> , Sweden	107
SP 191	CLINICAL EFFICACY OF POLYVINYLPIRROLIDONE-SODIUM HYALURONATE FOR THE TREATMENT OF ULCERS CAUSED BY ORTHODONTIC APPLIANCES <b>S-M Lee</b> , Korea	107
SP 192	EFFECTS OF SALVIA MILTIORRHIZA ON OSTEOBLASTIC CELL LINE OF MC3T3-E1 <b>A Chin</b> , SAR China	107
SP 193	CEPHALOMETRIC ASSESSMENT OF RAPID MAXILLARY EXPANSION IN CLASS II DIVISION 1 MALOCCLUSION SUBJECTS*** <b>M Pacurar</b> , Romania,	107
SP 194	POSTURAL INVESTIGATIONS IN CHILDREN AND JUVENILES WITH AND WITHOUT A CROSSBITE <b>S Kopp</b> , Germany	108
SP 195	ORAL HYGIENE IN ORTHODONTIC PATIENTS – A RANDOMIZED CLINICAL TRIAL <b>L Golland</b> , Switzerland	108
SP 196	DOES ORTHODONTIC TREATMENT HAVE ANY EFFECT ON MORPHOPSYCHOLOGY? <b>A Karamifar</b> , Iran	108
SP 197	BUCCAL CORRIDOR RATIO – DOES IT FOLLOW A RULE? <b>S Saadat</b> , Iran	108
SP 198	CRANIOFACIAL MORPHOLOGY IN SLEEP APNOEA PATIENTS WITH DIFFERENT CERVICAL COLUMN FUSIONS <b>P Svanholt</b> , Denmark	109
SP 199	IMPACTED MAXILLARY CANINES – AN EPIDEMIOLOGICAL STUDY OF TREATED PATIENTS <b>J Cotič</b> , Slovenia	109
SP 200	FORCE DEGRADATION OF ELASTOMERIC LIGATURES AND ELASTIC SEPARATORS USED AS AN ACTIVE TIEBACK <b>F Mahmoudi</b> , Iran	109
SP 201	APPLICATION OF A ER:YAG LASER FOR ORTHODONTIC BONDING. A PRELIMINARY IN VITRO STUDY <b>G Di Carlo</b> , France	109
SP 202	BONDING AND REBONDING OF DIFFERENT CERAMIC BRACKETS <b>F Falkensammer</b> , Austria	110
SP 203	PROFILE CHANGES AFTER MAXILLARY POSTERIOR IMPACTION <b>A Karamifar</b> , Iran	110
SP 204	ASSESSMENT OF THE PERMANENT DENTITION IN ADULTS WITH AN IDEAL OCCLUSION <b>M Ribackova</b> , Russia	110
SP 205	PATHWAYS BETWEEN SUBJECTIVE ORAL HEALTH, FACIAL PAIN, TEMPOROMANDIBULAR DISORDERS AND OCCLUSAL FEATURES <b>J Rusanen</b> , Finland	110
SP 206	ARRESTED ERUPTION OF THE PERMANENT LOWER SECOND MOLAR <b>N Toseska-Spasova</b> , Former Yugoslav Republic of Macedonia	111
SP 207	INCIDENCE OF OROFACIAL CLEFTS IN THE CZECH REPUBLIC FROM 2000-2006 <b>W Urbanova</b> , Czech Republic	111
SP 208	EFFECTS OF FLUORIDE-CONTAINING CASEIN PHOSPHOPEPTIDE-AMORPHOUS CALCIUM PHOSPHATE ON SHEAR BOND STRENGTH OF ORTHODONTIC BRACKETS <b>Ö Polat-Özsoy</b> , Turkey	111
SP 209	QUALITY AND VARIATION IN ORTHODONTIC POSTGRADUATE EDUCATION IN EUROPE <b>E Santigli</b> , Denmark	112
SP 210	ORTHODONTIC FEATURES IN CEREBRAL PALSY <b>E Pozzani</b> , Italy	112

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 211	INFLUENCE OF ORTHODONTIC TREATMENT ON CANDIDA COLONISATION IN THE ORAL CAVITY <b>A Wojtaszek-Słomińska</b> , Poland	112
SP 212	ASSESSMENT OF STAINING OF ORTHODONTIC AESTHETIC BRACKETS DUE TO FOOD COLOURANTS <b>K Malkiewicz</b> , Poland	112
SP 213	THREE-DIMENSIONAL STUDY OF THE DUMBO RAT <b>M Charezinski</b> , Belgium	113
SP 214	AN IN VITRO STUDY OF FRICTION OF CONVENTIONAL LIGATURES WITH SELF-LIGATING BRACKETS <b>M R Ricchiuti</b> , Italy	113
SP 215	SELF-LIGATING VERSUS CLEAR ALIGNERS ANALYSIS OF DENTOALVEOLAR EFFECTS <b>G Laganà</b> , Italy	113
SP 216	THE SUSCEPTIBILITY TO STAINING OF ORTHODONTIC ELASTICS LIGATURES <b>K Malkiewicz</b> , Poland	113
SP 217	PHOTOGRAMMETRIC PROFILE TREATMENT CHANGES OF CLASS II DIVISION 1 MALOCCLUSIONS <b>A Moullas</b> , Greece	114
SP 218	FACIAL PROPORTIONS IN SERBIAN FEMALE ADULTS <b>N Nedeljković</b> , Serbia	114
SP 219	IS A SECULAR TREND IN CRANIAL FORMATION AN EXPLANATION OF THE INCREASE IN MALOCCLUSIONS? <b>M Svabova</b> , Czech Republic	114
SP 220	PERCEPTION OF PAIN DURING ORTHODONTIC TREATMENT <b>J Petrovska</b> , Former Yugoslav Republic of Macedonia	114
SP 221	FORCE SYSTEM OF REVERSE CURVE NICKEL TITANIUM WIRES IN CONVENTIONAL AND SELF-LIGATING BRACKETS <b>I Sifakakis</b> , Germany	114
SP 222	DEVELOPMENTAL ANALYSIS OF MANDIBULAR FOSSA CIRCUMFERENCE EMPLOYING SAGITTAL COMPUTED TOMOGRAPHIC IMAGES*** <b>R Oyake</b> , Japan	115
SP 223	HUMAN PRIMARY TEETH REVEAL THE INFLUENCE OF EPITHELIAL CELLS AND INNERVATION IN RESORPTION <b>M-L B Bille</b> , Denmark	115
SP 224	DENTOSKELETAL CHARACTERISTICS IN SUBJECTS WITH ORTHODONTIC TREATMENT NEED IN TOOTH AGENESIS CASES <b>G Laganà</b> , Italy	115
SP 225	IN VITRO INVESTIGATION OF THE TENSILE BOND STRENGTH OF THREE DIFFERENT BRACKET TYPES <b>R J Speel</b> , Netherlands	115
SP 226	CHANGES IN ARCH FORM AND DENTITION WITH DIFFERENT RETAINERS AFTER ORTHODONTIC TREATMENT <b>S-S An</b> , Korea	116
SP 227	ELECTRON BEAM IRRADIATION OF POLYMER BRACKETS <b>A Faltermeier</b> , Germany	116
SP 228	THE INFLUENCE OF ORTHODONTIC ADHESIVE FILLER LEVEL ON BACTERIAL ADHESION AND BOND STRENGTH <b>A Faltermeier</b> , Germany	116
SP 229	ROOT RESORPTION IN ORTHODONTICS: COMPARISON OF TWO TREATMENT TECHNIQUES <b>E Pontarolo</b> , Italy	116
SP 230	EFFECT OF ORAL APPLIANCES ON INFLAMMATORY MARKERS IN OBSTRUCTIVE SLEEP APNOEA <b>A Nizankowska-Jędrzejczyk</b> , Poland	117
SP 231	STABILITY OF OPEN BITE CORRECTION IN CLASS II DIVISION 1 ADULTS WITH AND WITHOUT OSTEOTOMY <b>R B Kuitert</b> , Netherlands	117
SP 232	DIMENSIONAL CHANGES OF THE MAXILLARY APICAL BASE IN UNTREATED CLASS I SUBJECTS <b>J van der Vlag</b> , Netherlands	117
SP 233	FACIAL TRANSVERSE AND VERTICAL DEVELOPMENT DURING PRE-PUBERTAL GROWTH IN YOUNG BRAZILIANS <b>K Faltin Jr.</b> , Brazil	118
SP 234	IN VITRO STUDY LABORATORY TEST OF THE SHEAR BOND STRENGTH OF METAL BRACKETS BONDED TO ENAMEL <b>S I Penco</b> , Italy	118
SP 235	LONG-TERM OUTCOME AND STABILITY IN TREATED ANGLE CLASS II DIVISION 1 PATIENTS <b>K Nordås</b> , Norway	118
SP 236	IMPACTED MAXILLARY CANINES: IMPLICATIONS FOR ORTHODONTIC TREATMENT ON ROOT LENGTH <b>E M G Brusveen</b> , Norway	118
SP 237	AUTOMATED TRACKING OF FACIAL LANDMARKS FOR THREE-DIMENSIONAL ANIMATED MODELS, USING STEREOPHOTOGRAMMETRY <b>E Di Gioia</b> , Italy	119
SP 238	PULPAL REPAIR AFTER SUBCUTANEOUS TRANSPLANTATION OF CRYOPRESERVED IMMATURE TEETH IN RATS <b>L Temmerman</b> , Belgium	119
SP 239	THIN-PLATE SPLINE ANALYSIS OF THE DIFFERENT FACIAL TYPES IN FRONTAL NORM TELERADIOGRAPHS <b>K Faltin Jr.</b> , Brazil	119
SP 240	A TWO-JAW ORTHODONTIC APPLIANCE FOR A CLASS III MALOCCLUSION PATIENT TREATMENT <b>M Postnikov</b> , Russia	119
SP 241	A SYSTEMATIC REVIEW OF THE CORRELATION BETWEEN THE STOMATOGNATHIC SYSTEM AND BODY POSTURE <b>G Perinetti</b> , Italy	120

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 242	THE POSITION OF THE HYOID BONE INVESTIGATED BY DIFFERENT METHODS OF ANALYSIS OF A RADIOGRAPH I <b>Kirovski</b> , Former Yugoslav Republic of Macedonia	120
SP 243	APPRAISAL OF THE GEOMETRIC EFFECTS ON TWO CEPHALOMETRIC INDICES OF SKELETAL CLASS E <b>Di Maria</b> , Italy	120
SP 244	IS CASEIN PHOSPHOPEPTIDE-AMORPHOUS CALCIUM PHOSPHATE BENEFICIAL TO ORTHODONTIC PATIENTS? S <b>Saadat</b> , Iran	120
SP 245	ORTHODONTIC STATUS AND ORTHODONTIC TREATMENT NEED AMONG SEVENTH GRADE STUDENTS E <b>Deljo</b> , Bosnia and Herzegovina	120
SP 246	BOTULINUM TOXIN INFLUENCES THE HISTOLOGY AND GENE EXPRESSION OF DYSTROPHIC MASSETER MUSCLE T <b>Gedrange</b> , Germany	121
SP 247	EXPRESSION OF GENES INVOLVED IN CALCIUM HOMEOSTASIS IN DYSTROPHIC MASTICATORY MUSCLES C <b>Kunert-Keil</b> , Germany	121
SP 248	INFLUENCE OF THE MAGNITUDE OF INTRUSIVE FORCE ON EXTERNAL APICAL ROOT RESORPTION AND CHANGES IN ALVEOLAR CREST HEIGHT I <b>Marek</b> , Czech Republic	121
SP 249	KNOWLEDGE ON BIOSTATISTICS OF ORTHODONTIC POSTGRADUATE STUDENTS IN EUROPE T <b>Eliades</b> , Greece	121
SP 250	PREVALENCE OF MALOCCLUSION IN PATIENTS WITH TURNER SYNDROME C <b>Misevska</b> , Former Yugoslav Republic of Macedonia	122
SP 251	SOFT TISSUE PROFILE CHANGES IN THE EARLY TREATMENT OF A DEEP BITE WITH MYOFUNCTIONAL APPLIANCES M <b>Jämsä</b> , Finland	122
SP 252	ECTOPIC ERUPTION OF THE MAXILLARY FIRST PERMANENT MOLAR M <b>Zuzelova</b> , Former Yugoslav Republic of Macedonia	122
SP 253	MAXILLARY SECOND AND THIRD MOLAR INCLINATION AFTER EXTRACTION OF THE FIRST MOLARS IN CLASS II DIVISION 1 SUBJECTS C <b>Livas</b> , Netherlands	122
SP 254	KNOWLEDGE ON BIOSTATISTICS OF ORTHODONTIC POSTGRADUATE STUDENTS IN EUROPE T <b>Eliades</b> , Greece	123
SP 255	NASOLABIAL ANGLE VARIATIONS DUE TO UPPER INCISOR TORQUE AND POSITIONAL CHANGES J <b>Seoane</b> , Spain	123
SP 256	INFLUENCE OF SELF-ESTEEM ON QUALITY OF LIFE IN CHILDREN SEEKING ORTHODONTIC TREATMENT E <b>De Baets</b> , Belgium	123
SP 257	A SURVEY OF LEVELS OF DENTAL ANXIETY AMONGST INDIVIDUALS WITH CLEFT LIP AND PALATE K <b>Al-Moammar</b> , England	124
SP 258	VERIFYING THE TRACING OF TRUE HORIZONTAL AND TRUE VERTICAL USING THE INTRACRANIAL REFERENCE PLANE V <b>Lomakina</b> , Russia	124
SP 259	TREATMENT EFFECTS ON THE SOFT TISSUE PROFILE IN PATIENTS TREATED WITH THE HERBST APPLIANCE V <b>Lomakina</b> , Russia	124
SP 260	QUANTITATIVE ASSESSMENT OF RELAPSE AFTER RAPID MAXILLARY EXPANSION L <b>Vaida</b> , Romania	124
SP 261	EN MASSE RETRACTION OF ANTERIOR TEETH WITH THE BIDIMENSIONAL TECHNIQUE A <b>Bodini</b> , Italy	125
SP 262	IS THERE A RELATIONSHIP BETWEEN ORTHODONTIC AND PERIODONTAL TREATMENT NEED? R <b>Nalcaci</b> , Turkey	125
SP 263	ALLERGIES IN ORTHODONTICS – FACTS AND MYTHS G <b>Stefanović</b> , Serbia	125
SP 264	COMPARISON OF PLAQUE REMOVAL OF TWO POWER BRUSH HEADS VERSUS A MANUAL TOOTHBRUSH C <b>Erbe</b> , Germany	125
SP 265	ROOT RESORPTION AFTER ORTHODONTIC TREATMENT E <b>Kowalska</b> , Poland	126
SP 266	PHOTOGRAPHIC PROFILE ANALYSIS OF PATIENTS WITH PIERRE-ROBIN SEQUENCE A <b>Ifert</b> , Germany	126
SP 267	EVALUATION OF THE CRANIAL BASE ANGLE IN ACHONDROPLASTIC DWARFS – A CEPHALOMETRIC ANALYSIS M <b>de Mezer</b> , Poland	126
SP 268	FAMOTIDINE AFFECTS HISTOMORPHOMETRIC AND BIOCHEMICAL PARAMETERS DURING TOOTH MOVEMENT A <b>Meh</b> , Slovenia	126
SP 269	RADIOGRAPHIC ASSESSMENT OF THE PRESENCE OF THIRD MOLAR GERMS IN MACEDONIAN CHILDREN G <b>Curciewa-Cuckova</b> , Former Yugoslav Republic of Macedonia	127
SP 270	DENTAL ANOMALIES IN CHILDREN REGISTERED WITHIN THE CLEFT PALATE CENTRE OF STRASBOURG O <b>Matern</b> , France	127
SP 271	VITAMINS B6, B12 AND HOMOCYSTEINE IN CLEFT LIP AND PALATE: BIOCHEMICAL ANALYSIS A <b>T Altug</b> , Turkey	127
SP 272	THREE DIMENSIONS OF THE MAXILLA IN UNILATERAL VERSUS BILATERAL CANINE IMPACTION Z <b>Ulku</b> , USA	128

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 273	THE BENDING LOAD CAPACITIES OF RAPID MAXILLARY EXPANSION SCREWS – A PILOT STUDY <b>A P Muchitsch</b> , Austria	128
SP 274	FINITE ELEMENT ANALYSIS OF MINI-IMPLANT AND PERIPHERAL BONE DURING UPPER CANINE RETRACTION I <b>Bavbek Karakaya</b> , Turkey	128
SP 275	INFLUENCE OF PACIFIER NECK DIMENSIONS ON FLEXIBILITY AND INTRAORAL FORCE DISTRIBUTION <b>M Bertl</b> , Austria	128
SP 276	THREE-DIMENSIONAL MOVEMENT OF THE MAXILLARY FIRST MOLAR WITH THE QUADHELIX APPLIANCE*** <b>K Terada</b> , Japan	129
SP 277	ROOT RESORPTION AFTER 2.5 AND 15 DEGREE BUCCAL ROOT TORQUE FOR 4 WEEKS: A MICRO-COMPUTED TOMOGRAPHIC STUDY <b>C Colak</b> , Turkey	129
SP 278	A CLINICAL STUDY OF THE DELAYED ERUPTION OF THE MAXILLARY FIRST MOLAR*** <b>N Sano</b> , Japan	129
SP 279	FINITE ELEMENT ANALYSIS OF MINI-IMPLANT AND PERIPHERAL BONE DURING UPPER CANINE RETRACTION I <b>Bavbek Karakaya</b> , Turkey	129
SP 280	NOSE CHANGES FOLLOWING BIMAXILLARY SURGERY IN TURKISH SUBJECTS WITH CLASS III MALOCCLUSIONS <b>F Aydin Mustafa</b> , Turkey	129
SP 281	PERMANENT TOOTH-CROWN DIMENSIONS IN PREMATURELY BORN CHILDREN <b>L Paulsson-Björnsson</b> , Sweden	130
SP 282	DYNAMICS OF THE ERUPTION OF PERMANENT TEETH IN DIABETES TYPE I CHILDREN <b>M Koselj</b> , Slovenia	130
SP 283	THE EFFECT OF DIABETES TYPE 1 ON ORAL HEALTH FACTORS IN CHILDREN IN THE MIXED DENTITION PERIOD <b>M Cilenšek</b> , Slovenia	130
SP 284	BIOCOMPATIBILITY OF ORTHODONTIC BANDS AFTER EXPOSURE TO DENTAL PLAQUE <b>S Hornikel</b> , Germany	131
SP 285	RELATIONSHIP BETWEEN DENTAL MALOCCLUSION AND OTITIS MEDIA IN CHILDREN: A DESCRIPTIVE ANALYSIS <b>M Fabbrizi</b> , Italy	131
SP 286	MAXILLARY ARCH WIDTH AND PALATAL DEPTH IN PATIENTS WITH MARFAN SYNDROME <b>E Vahle-Hinz</b> , Germany	131
SP 287	DOES LOCATION OF CONGENITALLY MISSING TEETH AFFECT DENTOFACIAL MORPHOLOGY? <b>A Y Gungor</b> , Turkey	131
SP 288	DIFFERENCES IN SOFT TISSUE ANALYSIS BETWEEN CONVENTIONAL AND COMPUTERIZED TRACING METHODS <b>S Erken</b> , Turkey	132
SP 289	FORCES IN COMPLEX MALALIGNMENT CORRECTION WITH CONVENTIONAL AND SELF-LIGATING BRACKETS <b>M Montasser</b> , Germany	132
SP 290	THE WORLD-WIDE-WEB, ORTHODONTICS AND TEMPOROMANDIBULAR DISORDERS: A BALANCED INFLUENCE ON PATIENT DECISION-MAKING? <b>F Luther</b> , England	132
SP 291	DENTO-SKELETAL EFFECTS OF RAPID MAXILLARY EXPANSION WITH THE FERRO PROTOCOL <b>M Sorrentino</b> , Italy	132
SP 292	TIME-DEPENDENT BIOMECHANICAL BEHAVIOUR OF THE PERIODONTAL LIGAMENT – AN IN VITRO STUDY <b>K Papadopoulou</b> , Greece	133
SP 293	ORTHODONTIC FORCED ERUPTION IN THE COMPROMISED TOOTH: WEEKLY VERSUS SINGLE FIBROTOMY I <b>Frezzato</b> , Italy	133
SP 294	FORCES IN COMPLEX MALALIGNMENT CORRECTION WITH CONVENTIONAL AND SELF-LIGATING BRACKETS <b>M Montasser</b> , Germany	133
SP 295	ANALYSIS OF SUPPORTING AREAS AND ANGLES ON DENTAL PANTOMOGRAMS USING A NEW PROGRAM <b>R Reinhardt</b> , Ukraine	133
SP 296	EFFECTIVENESS OF THE TRANSPALATAL ARCH – A COMBINED CLINICAL AND EXPERIMENTAL STUDY <b>C Bouraue</b> , Germany	134
SP 297	BONE GRAFTING FOR ORTHODONTIC PURPOSES <b>C Reichert</b> , Germany	134
SP 298	DEFORMATION AND DEACTIVATION CHARACTERISTICS OF NICKEL-TITANIUM WIRES <b>B Pseiner</b> , Austria	134
SP 299	COMPARISON OF BOND STRENGTH OF EIGHT CONTEMPORARY ORTHODONTIC ADHESIVE SYSTEMS <b>C Reicheneder</b> , Germany	135
SP 300	ORAL HYGIENE INDEX IN MALE AND FEMALE PATIENTS BEFORE ORTHODONTIC TREATMENT <b>M Albreht Zore</b> , Slovenia	135
SP 301	ROOT RESORPTION AFTER LIGHT AND HEAVY ROTATIONAL FORCES FOR 4 WEEKS – A MICRO- COMPUTER TOMOGRAPHIC STUDY <b>T Tamer</b> , Turkey	135
SP 302	CONE BEAM COMPUTED TOMOGRAPHIC BASED TREATMENT OUTCOME ANALYSES OF ACTIVE AND PASSIVE SELF-LIGATING BRACKETS <b>P Cattaneo</b> , Denmark	135
SP 303	CEPHALOMETRIC PARAMETERS AND SPEECH ABILITY IN DOWN SYNDROME <b>E Pozzani</b> , Italy	136

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
SP 304	mRNA EXPRESSION OF REGULATORY AND GROWTH FACTORS IN OROFACIAL MUSCLES OF DYSTROPHIC MICE <b>A Spassov</b> , Germany	136
SP 305	CEPHALOMETRIC MORPHOLOGY OF ADULTS WITH OBSTRUCTIVE SLEEP APNOEA <b>A M Enache</b> , Romania	136
SP 306	USE OF DELMOPINOL GEL IN PATIENTS UNDERGOING FIXED ORTHODONTIC TREATMENT <b>D Dalessandri</b> , Italy	136
SP 307	EFFICACY OF AN INDIRECT BONDING TECHNIQUE IN REDUCING PLAQUE RETENTION <b>L Laffranchi</b> , Italy	137
SP 308	MUSCULAR EVALUATION IN SKELETAL CLASS II AND III SUBJECTS AT THE END OF GROWTH <b>G Galbiati</b> , Italy	137
SP 309	CORRECTION OF ASYMMETRIC TRANSVERSE DISCREPANCIES <b>G Galbiati</b> , Italy	137
SP 310	BOND STRENGTH OF CERAMIC BRACKETS USING DIFFERENT CURING MODALITIES <b>V Grassia</b> , Italy	137
SP 311	ORTHODONTIC-SURGICAL TREATMENT: ELECTROMYOGRAPHIC AND ELECTROGNATHOGRAPHIC EVALUATION <b>L Giannini</b> , Italy	138
SP 312	FUNCTIONAL EVALUATION IN ADULT PATIENTS WITH OPEN AND DEEP SKELETAL BITES <b>L Giannini</b> , Italy	138
SP 313	THE TRANSVERSE SAGITTAL MAXILLARY EXPANDER: A CEPHALOMETRIC INVESTIGATION IN GROWING PATIENTS <b>S Tronca</b> , Italy	138
SP 314	COMPARISON OF ARCHAIC AND CONTEMPORARY ITALIAN SKULLS USING THREE-DIMENSIONAL COMPUTED TOMOGRAPHIC VOLUME RENDERING IMAGES <b>G Tufano</b> , Italy	138
SP 315	DIAGNOSTIC SET-UP AND OCCLUSAL SPLINT GUIDE <b>F Iazzetti</b> , Italy	139
SP 316	IS CARIES IN THE EARLY MIXED DENTITION A PREDICTOR OF CARIES RISK DURING ORTHODONTIC TREATMENT? <b>V Soumas</b> , Switzerland	139
SP 317	EVALUATION OF MICROLEAKAGE WHEN BRACKETS ARE CURED FROM DIFFERENT SIDES <b>S I Ramoglu</b> , Turkey	139
SP 318	META-ANALYSIS OF SKELETAL MANDIBULAR CHANGES DURING FRÄNKEL APPLIANCE TREATMENT <b>R Cannavale</b> , Italy	139
SP 319	COMPARISON IN THE ADHERENCE LEVEL OF STREPTOCOCCUS MUTANS AND STREPTOCOCCUS SOBRINUS ON SELF-LIGATING BRACKETS*** <b>C-J Hwang</b> , Korea	140
SP 320	EFFECTS OF AMORPHOUS CALCIUM PHOSPHATE CONTAINING COMPOSITE ON ENAMEL DEMINERALIZATION <b>M Amasyali</b> , Turkey	140
SP 321	THE IMPORTANCE OF HEATH EDUCATION OF PATIENTS WITH FIXED ORTHODONTIC APPLIANCES <b>S Matijević</b> , Montenegro	140
SP 322	EVALUATION OF MAXILLOMANDIBULAR WIDTH USING THEE-DIMENSIONAL CONE BEAM COMPUTED TOMOGRAPHIC IMAGES <b>K-M Lee</b> , Korea	140



## Biotechnology and the Future in Orthodontics

### SP 001

Wednesday, June 16 - at 8:30, Screen 1

#### ASSOCIATION BETWEEN ORTHODONTIC TREATMENT NEED AND MAXILLARY INCISOR TRAUMA

**A Borzabadi-Farahani<sup>1</sup>**, A Borzabadi-Farahani<sup>2</sup>, F Eslamipour<sup>3</sup>,  
<sup>1</sup>Department of Orthodontics, Birmingham Dental Hospital, England, <sup>2</sup>Private Practice, Tehran and <sup>3</sup>Department of Community Dentistry, Isfahan University of Medical Sciences, Iran

**AIM:** To investigate the association between gender, malocclusion complexity, orthodontic treatment need and the prevalence of maxillary incisor trauma (MIT).

**SUBJECTS AND METHOD:** Five hundred and two subjects (253 females, 249 males, aged 11-14 years), including one female and five males who were wearing an orthodontic appliance. The Index of Complexity, Outcome and Need (ICON) scores were calculated and the subjects were categorized into five groups according to ICON complexity (easy, mild, moderate, difficult and very difficult). Subsequently they divided into two groups according to orthodontic treatment need (subjects with an ICON >43 and those with an ICON <44). Fisher's exact test was performed to determine any gender differences in trauma experience. Logistic regression was undertaken to test for any differences in risk of trauma among subjects in different ICON complexity groups, and also to estimate the predictive value of gender, orthodontic treatment need and ICON scores for MIT

**RESULTS:** Overall 9 percent experienced incisor trauma. Of those, 93 and 7 per cent had maxillary and mandibular incisor injuries, respectively. Males had greater odds of MIT compared with females (OR = 2.16). Subjects with an orthodontic treatment need (ICON > 43) had greater odds of MIT compared with those with no orthodontic treatment need (ICON < 44) (OR = 2.37). Only subjects presenting with a difficult complexity grade (64 < ICON < 77) showed significantly higher odds of experiencing MIT (OR = 3.16) when compared with the easy complexity group (ICON < 29)

**CONCLUSION:** Female gender decreased the risk of MIT by 46.2 per cent. Every unit increase in ICON score increased the risk of MIT by 1.2 per cent. The prevalence of incisor trauma in this study was rather low in comparison with other research. The current finding may be important in terms of targeting and screening certain vulnerable groups during the course of orthodontic treatment.

### SP 002

Wednesday, June 16 - at 8:30

#### NOCTURNAL ENURESIS AND RAPID MAXILLARY EXPANSION

**F Bazargani<sup>1</sup>**, T Nevéus<sup>2</sup>, L Leissner<sup>3</sup>, S Rudblad<sup>4</sup>, J Kuroi<sup>5</sup>,  
<sup>1</sup>Department of Orthodontics, Postgraduate Dental Education Center, Örebro, <sup>2</sup>Paediatric Nephrology Uppsala University Children's Hospital, Departments of <sup>3</sup>Neurology and <sup>4</sup>ENT, Örebro University Hospital, <sup>5</sup>Institute of Odontology at the Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden

**AIM:** Nocturnal enuresis (NE) is not uncommon in children. Rapid maxillary expansion (RME) has the effect of widening the maxillary suture as well as the nasal width. There have been a few reports of RME having anti-enuretic effects. The aim of this study was to prospectively evaluate RME treatment in a group of severely therapy-resistant enuretic children with long-standing re-

sistance to medical therapy, considering prognostic factors and effects on sleep and respiration.

**SUBJECTS AND METHOD:** Eleven children with NE, aged 9-15 years old (mean age = 12), who wet their bed almost every night and had never been dry. Sleep registrations, rhinomanometric measurements and bladder diaries were completed before and after RME, with an extra bladder diary completed with the orthodontic appliance *in situ* but before expansion of the maxilla to evaluate the possible placebo effect of RME for a period of four weeks. After this period the maxilla was expanded for 10-14 days. Mean RME was 6.0 mm (range = 5-7 mm).

**RESULTS:** All but three of patients became dry or almost dry. The number of wet nights per week pre-study, with the appliance *in situ* and after maxillary expansion was  $4.6 \pm 1.43$ ,  $3.9 \pm 1.64$ , and  $1.4 \pm 1.69$ , respectively. The anti-enuretic effect was highly statistically significant [Within-subjects ANOVA  $F(2, 20) = 39.7$ ;  $P < 0.001$ ]. No prognostically significant factors were found among anamnestic data or bladder parameters, but the non-responders noted that the amounts of urine lost in bed during wet nights decreased during treatment.

**CONCLUSION:** Orthodontic RME is a new option for treating children with NE and may be a potentially curative treatment of therapy-resistant nocturnal enuresis.

### SP 003

Wednesday, June 16 - at 8:30

#### CEPHALOMETRICS IN THE MAGNETIC FIELD\*\*\*

**T Brandt<sup>1</sup>**, I Buck<sup>2</sup>, S Kopp<sup>2</sup>, <sup>1</sup>Clinic Orthodontics Wiesbaden and <sup>2</sup>Department of Orthodontics, University of Frankfurt, Germany

**AIM:** New cephalometry without X-rays by means of a 92 g head cap is an accurate alternative to classic cephalostats. The new cephalometry is based on three-dimensional (3D) registration in a magnetic field. The carbon head cap, in combination with a 3D magnetic scanner, represents a digital presentation and cephalometric analysis technology for orthodontic diagnostics. One problem of conventional cephalometry is the exact and reproducible positioning in the cephalostat and its verifiability. The aim of this study was to determine the measurement accuracy of the noXrayCeph<sup>®</sup> device for repeated measurements.

**SUBJECTS AND METHOD:** Ten patients were measured with the noXrayCeph<sup>®</sup> device and the reproducibility of the projection to the midsagittal plane was examined by 10 repeated measurements of the same patient on different days. The differences in accuracy of the lateral cephalogram and the noXrayCeph<sup>®</sup> were analysed with a Mann-Whitney U test.

**RESULTS:** The accuracy of the production of a cephalometric drawing with the new magnetic device does not depend on standard positioning of the patient. The device produces reproducible drawings with different positioning of the patient's head. The error of the mean quartal deviation was < 0.05.

**CONCLUSION:** Due to the construction of a new midsagittal plane with each measurement, repeated evaluations are possible.

### EVALUATION OF DENTAL AESTHETICS AND ORTHODONTIC TREATMENT NEED

**E González Nieto**, B González Nieto, R Martínez de Aragón Lancha, A M González Sanz, M Romero Maroto, Universidad Rey Juan Carlos, Madrid, Spain

**AIM:** To quantitatively determine orthodontic treatment need and dental aesthetics by evaluating the level of malocclusion in a group of young Spanish adults using the Index of Orthodontic Treatment Need (ION).

**SUBJECTS AND METHOD:** Sixty young adults aged 20-32 years (61.7% females), without a history of previous orthodontic treatment. Their mean age was 24.38 years.

**RESULTS:** Forty per cent had an IOTN level 3 whereas 46.7 per cent were not in need of treatment or presented a low level of ION. The mean of IOTN was 2.50 with a standard deviation of 0.983. For the aesthetic component, all subjects considered photograph 10 to be the least attractive and 36.7 per cent placed themselves as one code from the ideal. Comparing both components of the IOTN it was found that 93 per cent of the subjects presented very low or no treatment need, while 38.3 per cent were classified as a moderate treatment need.

**CONCLUSION:** Ninety three per cent of subjects presented very low or no treatment need, a finding that was statistically significant. The largest percentage of subjects answered that their tooth corresponded to numbers 1 or 2 of the stimulus of aesthetic component photographs. Forty per cent presented level 3 of the IOTN, with a moderate or no treatment need, whereas 46.7 per cent had a low level and were not in need of treatment. There was no treatment need in 18.3 per cent.

### STRONTIUM DIMINISHES DELETERIOUS EFFECTS OF RUNX2 HAPLOINSUFFICIENCY IN HUMAN OSTEOBLASTS

**P Römer**, C Reicheneder, A Faltermeier, P Proff, Department of Orthodontics, University of Regensburg, Germany

**AIM:** Runx2 is an important transcription factor for osteoblast differentiation as it promotes expression of osteoblast markers, such as alkaline phosphatase (ALP) or osteocalcin (OCN). Heterozygous mutations of Runx2 cause cleidocranial dysplasia (CCD), due to the low functional activity of this gene. This congenital disease is responsible for patent fontanels and supernumerary teeth, as well as aplasia or hypoplasia of the clavicles. In a cell biological trial the ability of strontium to activate remaining functional Runx2 protein was examined. This activation was proven by determination of gene expression of ALP and OCN and by estimation of biomineralization of the extracellular matrix (ECM).

**MATERIALS AND METHOD:** Runx2<sup>±</sup> osteoblasts were isolated from human alveolar bone of a CCD patient. Osteoblasts were grown in the presence or absence of strontium for 10 days. Total RNA isolation and real time polymerase chain reaction were performed according to molecular biology standard protocols. Biomineralization assay was performed according to Gregory et al. (2004).

**RESULTS:** Strontium-exposed Runx2<sup>±</sup> osteoblasts exhibited statistically significantly enhanced expression of ALP ( $P = 0.02$ ) and OCN ( $P = 0.049$ ) in comparison with non-exposed Runx2<sup>±</sup> osteoblasts. The results of the gene expression study were counter-checked using a biomineralization assay, where strontium exposed adherent cells exhibited statistically significant ( $P = 0.002$ ) enhanced biomineralization of the ECM.

**CONCLUSION:** An improved effect of biomineralization in strontium-exposed Runx2<sup>±</sup> osteoblasts was observed, which might have promising potential in minimizing the disease pattern of CCD.

### STRONTIUM COUNTERACTS TOOTH RESORPTION BY INHIBITING EXPRESSION OF RANKL IN PERIODONTAL LIGAMENT CELLS

**P Römer**, C Reicheneder, A Faltermeier, P Proff, Department of Orthodontics, University of Regensburg, Germany

**AIM:** The function of the periodontal ligament (PDL) includes the attachment of the tooth in the alveolar bone, and the sensation of mechanical strain, as well as the support of alveolar bone formation and resorption when the tooth is moved by an orthodontic force. The effect of strontium in PDL cells regarding cytotoxicity and expression of RANKL was examined. So far, the effect of strontium has been exclusively studied in osteoblasts, where strontium enhances osteoblastic replication and activity, while counteracting the expression of RANKL. In this investigation the ability of strontium to inhibit the expression of osteoclastogenesis promoting cytokines, such as soluble Receptor Activator of NF- $\kappa$ B Ligand (sRANKL), in PDL fibroblasts (cells) was examined. Moreover, PDL cells were incubated with different concentration of strontium and cell viability determined in order to study the optimal concentration of strontium.

**MATERIALS AND METHOD:** PDL cells were isolated from the root of an extracted tooth. A force was applied in vitro on PDL cells to stimulate the expression of osteoclastogenesis promoting cytokines. Strontium-exposed PDL cells were employed in order to examine the effect of strontium on sRANKL expression in PDL cells. Expression of mRNA was determined by means of real time polymerase chain reaction, while protein expression of the cytokines was determined by enzyme-linked immunosorbent assay.

**RESULTS:** The treatment of pressed PDL cells with strontium resulted in a statistically significant down-regulation of osteoclastogenesis promoting cytokines on mRNA and protein level. Molar concentration of strontium above 10 mM exhibited cytotoxic effects in the PDL cells.

**CONCLUSION:** Strontium diminishes RANKL production in PDL cells. This might have clinical interest regarding tooth movement and inhibition of root resorption.

### THREE-DIMENSIONAL COMPARISON OF AVERAGE FACES OF CHILDREN WITH ORAL CLEFTS

**I Bugaighis**<sup>1</sup>, C Mattick<sup>2</sup>, R Hobson<sup>3</sup>, B Tiddeman<sup>4</sup>, <sup>1</sup>Dental School, Arab Medical University, Benghazi, Libya, <sup>2</sup>Royal Victoria Infirmary and <sup>3</sup>Windmill Dental Suite, Newcastle-upon-Tyne, <sup>4</sup>School of Computing Science, University of St Andrews, U.K

**AIM:** The introduction of three-dimensional (3D) non-invasive imaging systems and the implementation of advanced statistical analyses have widened the perspectives of their application to assist in clinical diagnosis and evaluation of treatment outcome through visualization. The aim of this research was to discriminate between 3D average faces of children with oral clefts and matched controls.

**MATERIALS AND METHOD:** A prospective cross-sectional 3D stereophotogrammetric study of the facial morphology of 103 children with non-syndromic oral clefts; 40 unilateral cleft lip and palate (CLP); 23 unilateral cleft lip and alveolus, 19 bilateral CLP; 21 cleft palate, and 80 gender and age matched controls.

**RESULTS:** Significant morphological differences between the mean facial shapes of all groups were found ( $P < 0.005$ ). Shape variation between the five group means was visualized by superimposing each average face with the control group average face. The differences were mainly in the nasolabial region but not confined to it, and more pronounced in groups where the lip was affected. This suggests that cleft lip and soft tissue lip repair has a greater effect on facial shape than cleft palate repair alone. These results emphasize the fact that a cleft palate only has a different aetiology than a CLP.

**CONCLUSION:** 3D shape analysis allows both subjective and objective morphometric discrimination between subjects with craniofacial anomalies and control groups, and underlines the potential value of statistical shape analysis in assessing the outcomes of CLP surgery and orthodontic treatment.

## SP 008

Wednesday, June 16 - at 8:30

### ASSESSMENT OF THREE METHODS TO ASSESS PLAQUE LEVELS AND WHITE SPOT LESIONS

E R Thomas, S M Higham, N Pender, University of Liverpool, England

**AIM:** Demineralisation during fixed appliance treatment is estimated to affect between 2-96 per cent of patients. It is frequently subclinical but can progress to irreversible lesions requiring restoration. Early detection is vital to institute appropriate management of this reversible condition. Quantitative light induced fluorescence (QLF) has the ability to detect subclinical mineral loss by quantitating changes in the autofluorescence of enamel. However, it is expensive and requires sophisticated computer software. Toothcare™ is a simple, inexpensive hand held device that has the potential to be 'home-based' allowing monitoring of plaque and early demineralisation. The aim of this study was to compare the effectiveness of Toothcare™ with QLF to identify plaque and demineralisation in orthodontic patients. The comparison was with QLF and clinical photographs.

**MATERIALS AND METHOD:** Patients undergoing fixed orthodontic treatment were consecutively recruited. At each of four routine appointments, the buccal surfaces of six anterior teeth were assessed for the presence of plaque and demineralisation using white light, QLF and Toothcare™. Plaque was scored using a modification of the Sillness and Løe plaque index; demineralisation was recorded as lesions per patient.

**RESULTS:** The overall mean observed plaque scores were highest for Toothcare™ at all visits 8.35 [Standard deviation (SD) 7.23, 95% confidence interval (CI) 1.32] followed by white light 3.81 (SD 4.13, 95% CI 0.75). QLF obtained the lowest plaque scores overall 0.71 (SD 1.2 95% CI 0.22). QLF measured statistically significantly more demineralised lesions than white light 0.72 (SD 1.12, 95% CI 0.2) compared with 0.12 (SD 0.43, 95% CI 0.08).

**CONCLUSION:** Toothcare™ did not demonstrate demineralisation when compared with white light and QLF. Toothcare™ did however demonstrate plaque more readily than white light and QLF; it was simple and easy to use at the chairside with few time implications.

## SP 009

Wednesday, June 16 - at 8:30

### INFLUENCE OF THE VISUAL SYSTEM TO HORIZONTAL JAW RELATIONSHIP DETERMINATION – A PILOT STUDY

D Ohlendorf, D Desoi, A Million, S Kopp, Department of Orthodontic, Goethe-University, Frankfurt am Main, Germany

**AIM:** Documentation of changes in the visual system to horizontal jaw relationship determination by means of an intraoral, electronic needle point tracing during standing.

**SUBJECTS AND METHOD:** Measurements were carried out in 10 healthy adults with open and closed eyes during standing. For recording the horizontal jaw relationship, and particularly lateral and sagittal movements of the mandible, the IPR System (IPR Systems GmbH, Germany) was used. The following movements were registered: initial and final contact position as well as protrusion, retrusion and transverse movement. In order to record the limiting movements of the mandible, casts and a facebow transfer are necessary to produce templates of the lower and upper jaw. During registration it is necessary to use an incisal blocking of 1 mm. Subsequently, the data were statistically analysed with the software program, BIAS, using Wilcoxon's matched pairs test.

**RESULTS:** The visual system had an influence on the final contact position ( $P \leq 0.05$ ). This influence was evident regarding the sagittal

axis, as the final contact position with the eyes closed was, on average, approximately 0.7 mm further retral. Because of deprogramming through the coordination movements carried out so far, the habitual contact position changed. This is normally used as an encryption position in horizontal jaw relationship determination.

**CONCLUSION:** There is no influence of the visual system on limiting movements of the mandible. The range of motion is limited by individually predetermined anatomical structures.

## SP 010

Wednesday, June 16 - at 8:30

### THE IMPACT OF FORCED LEG LENGTH DISCREPANCIES ON BODY POSTURE AND JAW POSITION

D Ohlendorf, K Pusch, S Kopp, Department of Orthodontic, Goethe-University Frankfurt am Main, Germany

**AIMS:** To determine how orthopaedic factors can affect body posture and movement sequences in the temporomandibular joint by documentation and analysis of the interdependencies between temporary provoked leg length discrepancies on upper body posture and the position and movement path of the mandible.

**SUBJECTS AND METHOD:** A 1 cm and 3 cm thick wooden panel was consecutively placed unilaterally under the left or right foot of 30 subjects (21 females, 9 males) while simultaneously recording upper body posture and movement trajectories of the mandible. The three-dimensional imaging system MiniRotKombi (ABW / GeBioM, Germany) allowed analysis of the upper body surface curvature by applying light-line projections while the electronic registration system, Arcus Digma (KaVo, Germany) recorded the mandibular position and movement paths based on ultrasound. Statistical analysis was carried out (t-test for correlated samples).

**RESULTS:** A unilateral increase of the foot caused a change in posture in relation to the pelvis and shoulder positions ( $P \leq 0.00$ ), which may be due to a compensating movement to the increase. The range of motion correlated with the height and relevant side of the increase. Likewise, a deviation of the mandibular condyle position in centric occlusion ( $P \leq 0.03$ ) and in mouth movement ( $P \leq 0.01$ ) was observed, which also increased with the amount of the increase in size. Positional related changes were observed in the ipsi- and contralateral side of the jaw.

**CONCLUSION:** There is a statistically reliable interdependence between leg length, body posture and mandibular position. This also shows that the human body is a complex system where different parts of the musculoskeletal system are inter-linked and are able to influence each other.

## SP 011

Wednesday, June 16 - at 8:30

### THREE-DIMENSIONAL MAXILLO-MANDIBULAR DENTAL CHANGES AFTER ORTHODONTIC TREATMENT WITH PREMOLAR EXTRACTIONS

E C K Wee<sup>1,2</sup>, K W C Foong<sup>1</sup>, I K F Lim<sup>2</sup>, <sup>1</sup>National University of Singapore and <sup>2</sup>National Dental Centre, Singapore

**AIMS:** Traditionally, measurements on lateral cephalographs only determine dental changes occurring in two planes, and whilst dental models provide three-dimensional (3D) information, relating the changes is difficult due to the lack of adequate reference structures for quantifying these changes. The aims of this study were: 1) To visualize and quantify dental changes to the maxillary and mandibular teeth in patients with a Class II division 1 incisor relationship after fixed appliance orthodontic treatment, involving extraction of four first premolars, versus extraction of two upper first premolars and two lower second premolars; 2) To investigate the effect that the two different extraction patterns have on the magnitude of dental change.

**MATERIALS AND METHOD:** Pre- and post-treatment records of 60 patients who had finished fixed appliance orthodontic treatment were randomly selected from completed premolar extrac-

tion cases. Thirty had extraction of four first premolars, whilst 30 had extraction of two upper first premolars and two lower second premolars. Pre- and post-treatment study models were digitally occluded and digitally superimposed on the pre- and post-treatment lateral cephalographs, which were then superimposed on each other using stable reference structures.

**RESULTS:** Statistically greater mean retraction of the lower canines ( $3.91 \pm 2.83$  mm versus  $1.88 \pm 3.11$  mm) and lower incisors ( $3.31 \pm 3.39$  mm versus  $1.52 \pm 2.95$  mm) was noted in the group with four first premolar extractions, whilst greater mean mesialization of the lower molars ( $4.66 \pm 2.67$  mm versus  $2.50 \pm 2.76$  mm) was noted in the group with lower second premolar extractions.

**CONCLUSION:** The type of premolar extraction may influence the amount of lower incisor and lower canine retraction, and the amount of lower molar mesialization that occurs during treatment. However, a great degree of individual variability exists, and other factors, such as pre-treatment parameters, may affect the type of dental change observed.

## SP 012

Wednesday, June 16 - at 9:15

### INFLUENCE OF CORROSION OF SURFACE DEFECTS OF NICKEL TITANIUM WIRES

**C Palazón Martínez**<sup>1</sup>, C Ábalos Labruzzi<sup>1</sup>, A Paúl Escolano<sup>2</sup>, E Solano Reina<sup>1</sup>, <sup>1</sup>Department of Stomatology, Faculty of Dentistry, University of Seville and <sup>2</sup>Mechanical Engineering and Materials Engineering Department, Superior School of Engineers of Seville, Spain

**AIM:** During recent years, the role of saliva and fluoride has been extensively studied as corrosive agents of the surface of nickel titanium (NiTi) wires. These studies have shown its relationship with surface roughness, that acidic pH is more corrosive and that the effects are more severe depending on the manufacturer, etc. However, the mechanisms and factors involved in this process are largely unknown. Thus the objective of this study was to determine the influence of different surface defects in as-received NiTi wire in the degradation of its surface.

**MATERIALS AND METHOD:** In a previous study of 0.016 X 0.022 inch NiTi wires the different surface defects were established. For each of the five patterns found, five samples of each pattern were randomly selected for *in vitro* analysis following immersion in artificial saliva and hydrofluoric acid for 28 days. The surface of the samples was studied before and after using scanning electron microscopy and confocal microscopy. Quantitative analysis of the surface defects and analysis of variance of the means obtained were performed.

**RESULTS:** The patterns found were smooth, striped, with holes or globular, cracked and porous. The increase of defects ( $\mu\text{m}^2$ ) and roughness, both in saliva and saliva plus fluoride were: smooth (28; 39/-0.4; 0.3), striped (26; 47/0.31, 0.87\*), globular (-23; -15/-0.17; -0.19), cracked (182\*, 629\*/-0.1; 1.3\*) and porous (18.3, 31.2/0.3, 0.5\*). \* =  $P < 0.05$ .

**CONCLUSION:** The smooth and globular patterns do not present a significant increase in surface defects. Conversely, the cracked pattern shows an increase in defects and roughness. The increase in roughness is not related to the increase in the number of defects.

## SP 013

Wednesday, June 16 - at 9:15

### DO DIFFERENT TENSILE BOND STRENGTH STUDIES END IN THE SAME RESULTS?

**S Hemmatpour**, Orthodontic Department, Tehran Azad University Dental School, Iran

**AIM:** To compare the results of three common methods used for measuring orthodontic tensile bond strength.

**MATERIALS AND METHOD:** Thirty-three brackets were bonded to human premolars using Transbond XT. The teeth were mounted and randomly assigned to one of the three groups. In group 1, tensile force was delivered using a custom made blade, in group

2, loops of wire were engaged under the bracket tie wings, and in group 3 debonding pliers were used. The adhesive remnant index was also scored for each specimen.

**RESULTS:** A significant difference existed between the three groups ( $P < 0.001$ ). The diametral compression method showed the highest ( $16.07 \pm 3.90$  MPa) and the custom made blade the least ( $6.28 \pm 2.58$  MPa) amount of tensile bond strength. The least dispersion was observed in group 2 (Coefficient of variation = 19.50%).

**CONCLUSION:** The results of orthodontic tensile bond strength tests may vary according to the method of force application. The results of the diametral compression test were significantly different from the other two groups. There is a need for standardizing orthodontic bond strength testing.

## SP 014

Wednesday, June 16 - at 9:15

### IN VITRO CORROSION BEHAVIOUR OF LINGUAL ORTHODONTIC ARCHWIRES

**C Suárez**<sup>1</sup>, J Gil<sup>2</sup>, P Sevilla<sup>2</sup>, T Vilar<sup>3</sup>, <sup>1</sup>Division d'Orthodontie, Université de Genève, Switzerland, <sup>2</sup>Escola d'Enginyeria Tècnica Superior Industrial, Universitat Politècnica de Catalunya and <sup>3</sup>Departament d'Odontostomatologia, Facultat d'Odontologia, Universitat de Barcelona, Spain

**AIM:** To investigate the *in vitro* electrochemical corrosive behaviour of archwires used in lingual orthodontics and to determine if corrosion affects the phase transition temperatures of nickel-titanium (NiTi) and copper-nickel-titanium (CuNiTi) archwires.

**MATERIALS AND METHOD:** Six different types of archwires were used: Respond, D-Rect, stainless steel (SS) 0.016, SS 0.016 X 0.022, Titanium molybdenum (TMA) 0.016, 0.016 NiTi and 0.017 X 0.017 35°C CuNiTi (Ormco Corp., Glendora, California, USA). Corrosion tests were performed following ISO-standard 10993-15:2000. Differential scanning calorimetry (DSC) was performed on NiTi and CuNiTi archwires of as-received samples, samples immersed for 30 days, and corroded samples. Scanning electron microscopic (SEM) images were obtained to examine surface changes before and after corrosion testing.

**RESULTS:** SEM analysis showed that all as-received samples had manufacturing defects. The SS archwires showed an Ecorr around -600 mV with the exception of 0.016 SS archwire that was found to be the less resistant to corrosion with an Ecorr of -845 mV. The titanium alloy archwires showed a higher resistance with Ecorr values around 1000 mV. DSC testing detected a rhombohedral phase (R-phase) in NiTi archwires but not in CuNiTi wires. A difference of 2°C to 3.5°C to the manufacturer's claim was found in the as-received and polarized samples, respectively.

**CONCLUSION:** The archwires tested showed similar corrosion resistance as those used in conventional orthodontics. The 0.016 SS archwires were found to be the less resistant to corrosion. An R-phase was detected for the NiTi archwires but not for the CuNiTi samples. No major differences were observed between groups concerning phase transformation temperatures.

## SP 015

Wednesday, June 16 - at 9:15

### THE TEMPOROMANDIBULAR JOINTS IN PATIENTS WITH SAGITTAL MALOCCLUSIONS

**I Skrypnyk**<sup>1</sup>, V Nespryadko<sup>2</sup>, Y-A Klytynskiy<sup>2</sup>, T Nespriadko<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Prosthetic Dentistry, National O.O. Bogomolets Medical University, Kiev, Ukraine

**AIM:** To investigate the temporomandibular joints (TMJ) using computed tomography (CT) in patients with sagittal malocclusions.

**SUBJECTS AND METHOD:** Eighty-seven patients aged between 7 and 38 years. The form of the glenoid cavity, its width and depth, as well as the form of the articular head of the mandibular bone and the size of the space between the head and the cavity in the anterior, middle and posterior areas were assessed together with

the degree of displacement of the articular head prior to, during, and at various periods following treatment.

**RESULTS:** In the initial occlusion 33 per cent of cases had a mandibular shift. While compensatory mechanisms were involved, in the stable occlusion, it tended to increase changes in the joint (29%) – decompensation. Treatment was planned having regard to the functional state of the TMJ.

**CONCLUSION:** functional and aesthetic optimum appearances with a, harmonious profile, facial symmetry, multiple interdigitation, creation of balanced occlusion and adequate dentoalveolar height in lateral and anterior areas were formed.

## SP 016

Wednesday, June 16 - at 9:15

### RESVERATROL INHIBITS PERIODONTAL PATHOGENS IN VITRO

**D J O'Connor**, R W K Wong, A B M Rabie, University of Hong Kong, SAR China

**AIM:** Oral hygiene maintenance is an essential requirement for successful orthodontic treatment in all patients, particularly 'high-risk' candidates with a history of periodontal disease. The gram-negative anaerobic bacteria, *A. actinomycetemcomitans* (Aa) and *P. gingivalis* (Pg), are key components in the aetiology of periodontal disease, and associated hard tissue destruction. Resveratrol is a phytoalexin, produced naturally by several plants when under attack by bacterial or fungal pathogens. It is found in many foods including mulberries, peanuts, and the skin of labrusca and muscadine grapes. The objective of this study was to evaluate the effect of resveratrol on growth of periodontal pathogens Aa and Pg in vitro. For comparison, resveratrol's effect on a variety of other oral microorganisms was also evaluated.

**MATERIALS AND METHOD:** As resveratrol demonstrates a poor solubility in water, different concentrations of resveratrol in the solvent dimethyl sulphoxide (DMSO) were added to calibrated suspensions of the microorganisms. As a control, a parallel series of dilutions containing the vehicle DMSO alone were used to measure the effect of the solvent. Minimum inhibitory concentrations of the periodontal pathogens were calculated. All suspensions were incubated for 1, 3, 6, and 24 hours in an anaerobic chamber at 37°C. At each time interval, selected dilutions from each culture broth were plated on blood agar plates. Colonies appearing on blood agar plates were visually counted at day 3 for Aa, and day 5 for Pg.

**RESULTS:** The periodontal bacteria, Aa and Pg, showed a significant decrease ( $P < 0.05$ ) in viable counts after 1 hour, whilst no colony forming units were observed after 24 hours. There was no effect on the other microorganisms tested.

**CONCLUSION:** Resveratrol possesses significant antimicrobial properties on periodontal pathogens in vitro.

## SP 017

Wednesday, June 16 - at 9:15

### USE OF AN ER,Cr:YSGG, WATERLASE LASER IN ORTHODONTIC PATIENTS WITH GINGIVAL OVERGROWTH

**P Thongdee**, A B M Rabie, R W K Wong, Orthodontics, University of Hong Kong, SAR China

**AIM:** Gingival overgrowth is frequently encountered among patients undergoing fixed orthodontic appliance treatment causing aesthetic and functional problems. Gingivectomy using a surgical laser followed by a strict oral hygiene regimen is one recently adopted treatment measure. The aim of this study was to evaluate the application of an Er,Cr:YSGG, waterlase laser in the treatment of overgrown gingivae in patients undergoing fixed orthodontic treatment.

**SUBJECTS AND METHOD:** Twenty-two patients (10 males, 12 females, mean age 16.5 years) who had gingival overgrowth. One hundred and eighty anterior teeth with gingival overgrowth were selected for laser surgery. At baseline, each patient received initial periodontal therapy and also maintained the recall programme after surgery. Prior to surgery, the depth of the pocket was meas-

ured and the patient was instructed to rate their level of pain on a visual analogue scale (VAS) over time. Gingivectomy and gingivoplasty was performed with an Er,Cr:YSGG, waterlase laser under topical anaesthesia with Tac 20 per cent Alternate (Professional Arts Pharmacy, Baltimore, Maryland, USA) for 5 minutes before surgery. VAS score was rated during surgery and post-operatively. Pocket depth measurement was carried out immediately after surgery and repeated.

**RESULTS:** Statistically significant changes were found in pocket depth scores (mean reduction 2.8 mm, Student's t-test and ANOVA,  $P < 0.005$ ). Recurrence was observed in 30 of 180 surgical sites (16.7%) or three out of 22 patients (13.6%) at 3 months following surgery. The mean VAS score was 3.2 during surgery. No patient required analgesia after surgery.

**CONCLUSION:** The Er,Cr:YSGG, waterlase laser is effective in the treatment of gingival overgrowth associated with fixed orthodontic treatment. A significant reduction in pocket depth and evident pain control can be achieved intra- and post-operatively

## SP 018

Wednesday, June 16 - at 9:15

### EFFECT OF DIET HARDNESS ON MANDIBULAR CONDYLAR CARTILAGE METABOLISM

P Tiilikainen, A Raustia, P Pirttiemi, University of Oulu, Finland

**AIM:** To study the effect of diet hardness on mandibular condylar cartilage catabolism markers. It was hypothesized that the change of diet hardness causes significant changes in proteolytic enzyme expression and cartilage composition.

**MATERIALS AND METHOD:** Seventy-two female Sprague Dawley rats were exposed to different dietary hardness, from soft to hard. MMP-3, -8 and TIMP-1 expression, cartilage thickness, cell count and expression of type-II collagen were studied and the condyles were examined at 30, 50 and 200 days.

**RESULTS:** A change of diet from soft to hard caused a significant decrease in the number of MMP-3, MMP-8 and an increase in TIMP-1 positive cells. The ratio of MMP-3 and TIMP-1 immunopositive cartilage cells remained similar in all age groups, whereas the number of MMP-8 positive cells decreased with age. Cartilage thickness and area of type II collagen positive staining were significantly affected by diet hardness.

**CONCLUSION:** A soft diet during growth increases collagenolytic activity and may increase the vulnerability of the condylar cartilage.

## SP 019

Wednesday, June 16 - at 9:15

### COMPOSITE OCCLUSAL BITE BLOCKS BONDED WITHOUT LIQUID RESIN

**A T H Tang**, U Hägg, Orthodontics, University of Hong Kong, SAR China

**AIM:** Liquid resin in bonding composite has been identified as a major irritant in human cell culture. However, liquid resin is assumed to play a critical role in composite adhesion to enamel by forming resin tags to mechanically interlock with acid-etched enamel prisms. By bonding occlusal bite blocks to enamel surfaces without the use of liquid resin, mechanical interlocking between the liquid resin tags and acid-etched enamel prisms should not exist to enable the bite blocks to resist heavy functional occlusal forces. Survival analysis of a case series of composite bite blocks bonded without liquid resin will demonstrate if mechanical interlocking is critical in the mechanism of enamel adhesion. The aim of this study was to evaluate the role of mechanical interlocking in the mechanism of enamel adhesion of composite resin in vivo.

**SUBJECTS AND METHOD:** Twenty-three patients had bilateral occlusal bite blocks bonded on the posterior teeth to raise the bite during fixed orthodontic therapy. Depending on the individual clinical situation and enamel quality, the choice of the occlusal surfaces of teeth for the bite blocks bonded without liquid resin included 55 and 65 (1 case), 17 and 27 (1 case), 16 and 26 (14

cases), 35 and 45 (1 case) and 36 and 46 (6 cases). The service durations of the occlusal bite blocks recorded were analyzed using the Kaplan Meier product limit method.

**RESULTS:** Twenty-two of the 23 patients (95.7%) experienced no failure of the bilateral occlusal bite blocks. Forty-five of 46 (97.8%) occlusal bite blocks lasted up to the end of their intended treatment purposes. They remained on average for 27.8 months (range 1.5 to 29.0 months), and the only failure occurred at 1.5 month. The failed bite block was replaced to finish the treatment (additional 3.1 months).

**CONCLUSION:** This clinical trial demonstrates that mechanical interlocking does not seem to have a critical role in the mechanism of enamel adhesion *in vivo*.

## SP 020

Wednesday, June 16 - at 9:15

### DEVELOPMENTAL ANALYSIS OF ETHMOIDAL SINUS EMPLOYING SAGITTAL COMPUTED TOMOGRAPHIC SCANS\*\*\*

**M Numata**, H Innami, F Yamaaki, School of Dental Radiology, Meikai University, Japan

**AIM:** To investigate the use of computed tomographic (CT) image analysis to assess the relationship between sinus development stage and occlusal development.

**MATERIALS AND METHOD:** Dried human skulls served as the experimental material; five samples were selected for each stage and a total of 50 samples were examined. CT imaging of the ethmoidal sinus was performed with the Somatom Emotion 6 (Siemens). Imaging conditions were as follows: X-ray tube voltage of 130 kv, tube current of 63 mA in bone mode, scan duration of 2.7 seconds and slice interval of 2 mm. The images of the sagittal view at the midline and the axial view at the Frankfort horizontal plane were acquired at WW3000;WL480. Analyzed images were transferred to a high-speed three-dimensional analysis device, Virtual Place Advance Plus (AZE Ltd.), in order to construct images for measurement. The area and growth rate of the sagittal image and the area of the ethmoidal sinus were then measured on the images and evaluated.

**RESULTS:** 1. The area of the sagittal image at stage IA, which was about 2 cm<sup>2</sup>, increased to approximately 9 cm<sup>2</sup> by stage VA. The growth rate at stage VA was approximately 4.5-fold based on stage IA. The increase in size after stage IC, from about 5.5 to 9 cm<sup>2</sup>, was linear. 2. The growth distance in the sagittal image was identical in all sections, including the superior section, the uppermost two-thirds, and the inferior section of the ethmoidal sinus. The length at stage IA, which was about 3.5 cm, increased to approximately 6 cm by stage VA. The growth rate at stage VA was approximately 1.7-fold based on stage IA. The increase after stage IC, from about 5.04 to 5.88 cm, was linear.

## SP 021

Wednesday, June 16 - at 9:15

### STEM CELL DIFFERENTIATION INTO ANGIOGENESIS LINEAGE

**A Almoqubel**, R W K Wong, P Zuo, A B M Rabie, Prince Philip Dental Hospital, University of Hong Kong, SAR China

**AIM:** Fibroblast or fibroblast growth factors (FGFs), are a family of growth factors involved in angiogenesis, which is an important process in growth and development, as well as in wound healing. The aim of this study was to evaluate the possibility to differentiate human dental pulp mesenchymal stem cells into angiogenesis lineage. Their differentiation ability was evaluated with an inducer, human FGF1 transfection.

**MATERIALS AND METHOD:** A strain of stem cell was isolated from dental pulp, expanded and further characterized. Human FGF1 was chosen to trigger it to differentiate to angiogenesis. The gene was cloned and recombinant as pcDNA3.1/V5-His-TOPO-FGF1 vector. pcDNA3.1/V5-His-TOPO-FGF1 vector was transfected into stem cell as well as three different negative controls, blank cell without transfection, empty vector and non-related green fluorescent protein (GFP) vector. After 48 hours, the mRNA level of

both hepatocyte growth factor (HGF) and urokinase-type plasminogen activator (uPA) were measured by real-time polymerase chain reaction for confirm angiogenesis differentiation.

**RESULTS:** The transcription levels of HGF and uPA mRNA were significantly higher than all three negative controls after 48 hours transfection. The relative levels of HGF mRNA were 4.078 for blank cell, 12.58 for empty vector, 17.97 for non-related GFP vector and 48.59 for pcDNA3.1/V5-His-TOPO-FGF1 vector. The relative transcription levels of uPA were significantly higher than all three negative controls. The relative levels of uPA mRNA were 0.171 for the blank cell, 3.088 for the empty vector, 3.762 for the non-related GFP vector and 10.15 for the pcDNA3.1/V5-His-TOPO-FGF1 vector.

**CONCLUSION:** The expression of markers of angiogenesis HGF and uPA is an indication of successful differentiation of human pulp stem cells to the angiogenic lineage. The protocol presented above is effective to trigger angiogenesis from dental pulp stem cells.

## SP 022

Wednesday, June 16 - at 9:15

### ELECTROMYOGRAPHIC AGE VALUES OF STOMATOGNATHIC MUSCLES

**N Pankratova**, N Nabiev, D Lyakhov, L Persin, M Kolesov, Department of Orthodontics and Children Prosthetics, Moscow State University of Medicine and Dentistry, Russia

**AIM:** To study the bioelectric activity of the stomatognathic muscles in average rectified value (ARV) and root mean square (RMS).

**SUBJECTS AND METHOD:** Thirty-five children aged 6-9 years with a physiological occlusion. The electromyographic activity of the stomatognathic muscles were determined (BKN electromyograph, Biotronic, Italy). Bioelectric activity of the temporalis, masseter, digastric and sternocleidomastoid muscles on the right and left sides was studied in three functional tests: mandibular physiologic position; physiologic occlusal position, and maximal pressure of teeth (dental arches). Summary tables were compiled from the average measurements of the age indices. The electromyographic parameters of the stomatognathic muscles were compared.

**RESULTS:** In the physiological relaxed position the bioelectric activity of the muscles that lift the mandible were larger than those of the muscles that descend the mandible. Bioelectric activity of lower jaw descending muscles on the left side was less than on the right side. The amplitudes of maximal single biopotentials of all muscles were equal. No change in the bioelectric activity of the muscles was found when the mandible was in a physiological occlusion (compared with a previous test). At maximal tooth pressure, muscle bioelectric activity increased 10 fold. Bioelectric activity of the lower jaw descending muscles increased: digastric on both sides up to X7 and sternocleidomastoid up X3 times. Amplitudes of maximal single biopotentials increased in the lower jaw lifting muscles up to X200; in the lower jaw descending muscles up to X6 and in the sternocleidomastoid up to X4.

**CONCLUSION:** The findings show the necessity of studying different age group indices of bioelectric activity of the stomatognathic muscles by electromyography. Without exception, all bioelectric activity changes clinically observed were also tracked on the ARV and RMS indices.

## SP 023

Wednesday, June 16 - at 10:00

### SIGNIFICANCE OF AGE ON LOWER JAW MOTION – A THREE DIMENSIONAL STUDY

**N Pankratova**, L Persin, D Lyakhov, N Nabiev, T Klimova, Department of Orthodontics and Children Prosthetics, Moscow State University of Medicine and Dentistry, Russia

**AIM:** To study the characteristics of mandibular motions and to determine the age norms of patients with a physiologic occlusion.

**SUBJECTS AND METHOD:** Ninety-three patients aged 7-15 years with physiological occlusion in three age groups 7-9, 10-12 and

13-15 years. Mandibular motions were studied using a Kinesio-graph tracking device (Biotronic, Italy). The following functional tests were executed: maximal descent and lift, side (lateral) motions and maximal forward mandibular movement.

**RESULTS:** According to age, there is an increase in the kinesio-graphic index in the sagittal plane for mandibular motion trajectory diagonal length and the distance of mandibular movement to the initial position. The angle between the diagonal and vertical planes did not show any difference, while the angle index at the beginning and end of descent and lifting of the mandible did not change with age. In all groups the motion ending angle was larger than the motion beginning angle. In the frontal plane the descent of the mandible motion trajectory, and the length of descent and lift trajectories increased with age. Mandibular angle indices at the beginning and end of descent and lifting did not change with age. Graphical images of mandibular side motions were the most informative in the frontal plane. Linear and angular parameters in the age groups with this motion did not differ. Mandibular left and right motion lengths were equal. The distance of mandibular forward movement during maximal protrusion increased with age, while the descent and lift angles decreased. In the frontal plane mandibular descent during forward movement was lowest.

**CONCLUSION:** The present research indicates necessity of having age group indices of mandibular motions.

#### SP 024

Wednesday, June 16 - at 10:00

##### PULPAL REACTION TO LOADING APPLIED TO UPPER TEETH IN PATIENTS WITH CLASS II MALOCCLUSIONS

**M Szumielewicz**<sup>1</sup>, T Matthews-Brzozowska<sup>2</sup>, <sup>1</sup>Department of Maxillofacial Orthopaedics and Orthodontics, Wrocław Medical University and <sup>2</sup>Orthodontic Clinics, Poznan University of Medical Science, Poland

**AIM:** Orthodontic therapy with fixed appliances results in exposure of teeth to continuous forces for a long period of time. Orthodontic force influences the pulp condition of the teeth. The most representative indicator of pulpal vitality is estimation of the function of the pulpal vascular system. Reliable evaluation of microcirculation *in vitro* can be carried out using laser Doppler flowmetry. The aim of the study was to compare the perfusion in pulpal blood vessels of the upper incisors and canines in patients with a Class II malocclusion treated with edgewise appliances, with or without extractions.

**MATERIALS AND METHOD:** Forty-eight upper anterior teeth of young adults. Straightwire appliances were used for both group of patients, treated either with extractions or by mandibular advancement using Class II elastics. Perfusion was estimated using a laser Doppler flowmeter, MBF3D, and model P13 probe.

**RESULTS:** A statistically significant blood-flow decrease during orthodontic treatment was found in the pulp of all examined teeth. There was a significant correlation between the type of loading and blood flow for the lateral incisors in patients treated with extractions, and for the canines in patients treated with mandibular advancement.

**CONCLUSION:** During orthodontic treatment functional changes occur in the pulpal vascular system. The tooth with the most intense pulp reaction to loading depends on the treatment mechanics.

#### SP 025

Wednesday, June 16 - at 10:00

##### CLEFT LIP AND PALATE PATIENTS AND HYPODONTIA – A SINGLE NUCLEOTIDE POLYMORPHISM STUDY

**B Nemes**<sup>1</sup>, G Jobbágy-Óvári<sup>2</sup>, G Varga<sup>2</sup>, A Végh<sup>1</sup>, <sup>1</sup>Department of Orthodontics and Dentofacial Orthopedics, Heim Pál Children's Hospital and <sup>2</sup>Department of Oral Biology, Semmelweis University, Budapest, Hungary

**AIM:** There is increasing evidence that supports the assumption that genetic factors increase the risk for susceptibility of individuals to cleft lip and palate (CLP) and/or hypodontia. The purpose of

the study was to map the single nucleotide gene polymorphisms (SNP) purported to play an important role in the development of the clefts and hypodontia

**SUBJECTS AND METHOD:** One hundred and three selected CLP patients examined radiographically genetically and clinically. Sixty had congenitally tooth-germ agenesis and 43 a complete dentition. For gene polymorphism, DNA was isolated from oral mucosa scrapings. A combined and optimized polymerase chain reaction and restriction fragment length polymorphism analysis was used to identify individual SNPs. Subsequently the SNPs were genotyped by the TaqMan SNP genotyping assay kit.

**RESULTS:** Five SNP procedures were successfully optimized that are putatively involved in CLP and hypodontia (PAX9 -1031G/A, -912C/T, MSX1 3755 A/G, FGFR1 C/T rs881301, IRF6 A/G rs764093). Among the investigated five SNPs of four genes homozygotes rare alleles in all cases were found. The frequency of appearance of the minor G allele of Msx1 was significantly higher in the cleft with hypodontia group than in the cleft /complete dentition group.

**CONCLUSION:** Missing teeth as well as aplasia occurs more frequently in children with a CLP than in other configurations of the nasal floor. The results clearly show the feasibility of recognition and understanding of the genetic/genomic factors

#### SP 026

Wednesday, June 16 - at 10:00

##### ENZYMATIC ACTIVITY IN GINGIVAL CREVICULAR FLUID WITH SELF-LIGATING BRACKETS

L Bolamperti, A Macchi, L Levrini, A Tagliabue, **A Caprioglio**, University of Insubria, Italy

**AIMS:** Bone remodelling is a biological process involving an acute inflammatory response. Different gingival crevicular fluid (GCF) constituents have been shown to be diagnostic markers of active tissue destruction in periodontal disease. Lactate dehydrogenase (LDH) is an enzyme usually limited to the cytoplasm of cells and it is released extracellularly only after cell death. Its activity in GCF is significantly correlated with gingival inflammation and tissue destruction and these phenomena also take place in the periodontal ligament during orthodontics. The aim of this study was to investigate the activity of LDH in GFC during orthodontic tooth movement using self-ligating brackets. The usefulness of this enzyme as a potential biomarker of tissue response to orthodontic treatment was also examined.

**SUBJECTS AND METHOD:** Forty patients selected according to the following criteria: similar age, equal relationship between genders, FMA ~27 degrees, SN<sup>∧</sup>GoGn ~37 degrees and mandibular anterior crowding 4-8 mm with no need for extractions. They were divided into two groups, according to the type of self-ligating bracket used. Each patient received professional oral hygiene and instructions for correct home care. GCF was collected at one site for each tooth, at baseline, one hour after bonding and on days 7, 21 and 42. The test teeth were 4 1, 4.3 and 4.5 and the control teeth 1.1, 1.3 and 1.5. LDH activity was estimated using a LDH cytotoxicity assay kit. Data were analyzed using dedicated software.

**RESULTS:** Statistical analysis showed no significant differences in LDH activity among the test and control teeth, but differences were present when orthodontic forces were applied.

**CONCLUSION:** LDH activity can be successfully estimated in the GFC using dedicated assay kits. LDH can be used as a potential marker for tissue metabolism for orthodontic mechanics, even if further investigations on its activity may be necessary.

### COMPARISON OF CEPHALOMETRIC MEASUREMENTS: COMPUTERIZED VERSUS CONVENTIONAL ANALYSIS

M Iacob, I N Zetu, V Dorobat, UMF 'Gr. T. Popa', Iasi, Romania

**AIM:** To evaluate intra-examiner repeatability and reliability of cephalometric measurements, using software and the manual tracing technique.

**MATERIALS AND METHOD:** Sixty lateral cephalograms were randomly selected and 11 angular and seven linear parameters were traced and measured using the manual method and Orthalis 4.0. A Student's t-test for independent and paired samples was used to compare the mean values of the differences.

**RESULTS:** The general degree of reproducibility was high. Significant statistical differences were identified for FMIA, interincisal angle, SNOcl, SNGoGN, ANB and for I/NA, AoBo. The repeatability of the computerized measurements was greater when compared with the manual tracings. Only FMIA angle showed a difference greater than 2 degrees, which was considered to be clinically unacceptable. The overall reliability of the computerized measurements was clinically acceptable. Statistically significant differences were identified for SNA, SNB, ANB, SND, SNOCl, FMIA, IMPA, angle Z and AoBo, UL, TC, Hp and Ha. Clinically unacceptable differences were observed for SNOcl, FMIA, Z angle, Hp and Ha.

**CONCLUSION:** Measurements with the aid of Orthalis 4.0 cephalometric software presented an acceptable degree of repeatability and reliability when compared with conventional measurements. The time saving with Orthalis 4.0 can be an advantage.

### A COMPARATIVE STUDY ON THE ALIGNMENT EFFICIENCY OF A LINGUAL SYSTEM

S J Zhu, Y Q Yang, R W K Wong, University of Hong Kong, SAR China

**AIM:** The Incognito lingual system is characterized by customized brackets and wires, and thus can improve patient comfort and save chair-side time for orthodontists. However, treatment efficiency has not been studied, especially for Asian patients, whose teeth are commonly of a smaller size and lingual surface area. The objective of this study was to evaluate the alignment efficiency of the Incognito lingual system in Asian patients.

**SUBJECTS AND METHOD:** Twenty-one Asian patients treated with the Incognito lingual system were compared with a randomly selected group of 21 adult Asian patients treated with labial fixed appliances. Each arch was considered as independent, thus the sample number was 42 in both the study and control group. Two indices were used to evaluate the alignment efficiency: the treatment period (TP) was the total treatment time (months) from the date of full mouth bonding to the date when the rectangular stainless steel wire was fully engaged, and crowding release efficiency (CRE) = crowding (mm) in individual arch/corresponding TP. Statistical differences were analyzed using a t-test.

**RESULTS:** The Incognito system showed higher CRE during the alignment stage compared with the labial straightwire appliance ( $2.5 \pm 0.2$  mm versus  $1.8 \pm 0.35$  mm per month), and shorter TP of the alignment stage ( $4.6 \pm 0.5$  versus  $6.2 \pm 0.75$  months). The difference was statistically significant ( $P < 0.05$ ).

**CONCLUSION:** The Incognito lingual system is more efficient in the alignment stage compared with commonly used labial straightwire system.

### ROLE OF FLUORIDE AND CARBONATED DRINKS IN THE CORROSION OF NICKEL TITANIUM WIRES

C Palazón Martínez<sup>1</sup>, C Ábalos Labruzzi<sup>1</sup>, A Paúl Escolano<sup>2</sup>, A Jiménez Planas<sup>1</sup>, E Solano Reina<sup>1</sup>, <sup>1</sup>Department of Stomatology, Faculty of Dentistry, University of Seville and <sup>2</sup>Mechanical Engineering and Materials Engineering Department, Superior School of Engineers of Seville, Spain

**AIM:** The influence of saliva and fluoride on nickel titanium (NiTi) wires surface has been widely studied. However, the role of carbonated drinks has not been thoroughly investigated and there are no studies co-relating these corrosive agents with surface patterns based on factory defects. The objective of this research was to determine the influence of hydrofluoric acid and Coca-Cola® on the surface defects of different as received NiTi wires.

**MATERIALS AND METHOD:** A sample of 22 NiTi wires, 0.016 X 0.022 inches, previously studied to establish different wire patterns according to their main defects. Five samples of each wire pattern were randomly selected and placed in vitro with fluoride acid + artificial saliva and artificial saliva + Coca-Cola®, for 28 days. The surfaces were analysed before and after the experiment, with scanning electron and confocal microscopy. A quantitative analysis of surface defects and the medians were calculated. ANOVA was applied to evaluate the statistical significance of the means. The patterns studied were smooth, striped, with holes or globular and cracked.

**RESULTS:** The increase of defects in 10.000 X<sub>m</sub><sup>2</sup> for artificial saliva + fluoride and for saliva + Coca-Cola® were, respectively: smooth (39/1.8); striped (47/182.5\*); globular (-15/1727\*); cracked (629\*/1353\*). The increase in roughness for saliva + fluoride and saliva + Coca-Cola® was: smooth (0.3/-0.3); striped (0.87\*/0.35); globular (-0.19/0.1); cracked (1.3\*/1.2\*). \* = P 0.05.

**CONCLUSION:** The smooth pattern showed higher corrosion resistance. The cracked pattern was the most corroded. The medium fluoride produced a greater increase in roughness, and Coca-Cola® more surface defects.

### FINITE ELEMENT ANALYSIS OF ORTHODONTIC IMPLANTS WITH DIFFERENT DESIGNS

C Szuhane<sup>1</sup>, A Cernescu<sup>2</sup>, N Faur<sup>2</sup>, F Glavan<sup>1</sup>, R Jianu<sup>1</sup>, <sup>1</sup>University of Medicine and Pharmacy Timisoara and <sup>2</sup>Politechnic University of Timisoara, Romania

**AIM:** To comparatively investigate the effect of various designs on the biomechanical properties of the orthodontic implant/bone interface.

**MATERIALS AND METHOD:** Finite element models of three different types of orthodontic implants were used. The orthodontic implants had different neck designs but were of the same length and were subjected to the same orthodontic force. The first two models simulated commercially available orthodontic implants, while the third was a modification. The stress and strain at the interface between the implant and bone were evaluated using COSMOSM™ software.

**RESULTS:** Maximum stresses were always located around the neck of the implant, in the marginal bone. The modified implant design resulted in improved distribution of the stress at the neck/alveolar bone interface.

**CONCLUSION:** The design of orthodontic implants has a significant influence on primary stability and in achieving clinical success.

### EVALUATION OF CEPHALOMETRIC MEASUREMENTS OF PATIENTS AFTER ORTHOGNATHIC SURGERY

**M Janega**, A Rehacek, P Hofmanová, T Dostalova, Departments of Paediatric Stomatology, LF UK a FN Motol and 1.LF UK, Prague, Czech Republic

**AIM:** Lateral cephalometric radiography is widely used as a descriptive, analytical, and diagnostic tool in clinical orthodontics and prosthodontics. The application of numbers to the radiographic images gives the impression of mathematical accuracy. The purpose of this study was to evaluate lateral cephalometric radiographs taken before and after surgical treatment of orthognathic anomalies and to compare the post-operative radiographs with normative cephalometric data.

**SUBJECTS AND METHOD:** Twenty-nine Caucasian orthognathic patients aged from 18 to 40 years. The group consisted of 15 Class III patients, six Class II patients and eight open bite patients. Images were taken with the head in a natural position, the teeth in centric occlusion, and the lips in repose. The analysis of McLaughlin and Arnett was used. Fourteen measurements (computer analysis Dolphin Imaging 11.0) were used to determine the skeletal sagittal jaw relationship. Measurement of the soft tissues was also carried out. The cephalograms were analyzed twice by two dentists. Five randomly selected radiographs were remeasured after a period of two weeks. There were no significant differences in the measurements.

**RESULTS:** The measurements after surgical and orthodontic treatment were close to normative international standards. After orthognathic therapy there was a return to a skeletal and dental Class I relationship. The results showed that SNA, SNB, ANB angle was successfully treated in comparison with the standards. Wits appraisal remained deviated in the direction of the original defect.

**CONCLUSION:** Dental and skeletal cephalometric values after orthognathic therapy are very close to standard values. More discrepancies were found for soft tissue values.

### INFLUENCE OF ADHESIVE SURFACE ROUGHNESS ON STREPTOCOCCAL ADHESION FORCES

L Mei<sup>1</sup>, H J Busscher<sup>2</sup>, H C van der Mei<sup>2</sup>, Y Ren<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Biomedical Engineering, University Medical Centre Groningen and University of Groningen, Netherlands

**AIM:** To investigate the influence of adhesive surface roughness (Ra) on streptococcal adhesion forces.

**MATERIALS AND METHOD:** The Ra of orthodontic light-cured adhesives, Transbond XT and PADLock, were varied by abrading with aluminium and sandpaper, and divided into three groups: untreated (Ra 20 nm), aluminium (Ra 150 nm) and sandpaper (Ra 350 nm). Adhesion forces between streptococcus and adhesive surfaces, in the absence and presence of a salivary conditioning film, were measured by atomic forces microscopy at 0 and 120 seconds surface delay times. Comparisons of the adhesion forces were performed using non-parametric and Weibull analyses.

**RESULTS:** Bacterial adhesion forces increased with surface roughness in the absence of a salivary conditioning film at 120 seconds surface delay time ( $P < 0.05$ ). Roughening treatments (aluminium and sandpaper abrading) increased the adhesion forces compared with the untreated surfaces at 120 seconds surface delay time in the presence of a salivary conditioning film ( $P < 0.05$ ), and at 0 seconds surface delay time in both saliva conditions ( $P < 0.05$ ). Data at 120 seconds surface delay time [Weibull modulus (m)  $4.0 \pm 1.5$ ] were more reliable than at 0 seconds (m:  $3.0 \pm 0.7$ ) for comparison of influence of Ra on streptococcal adhesion forces, regardless of salivary conditions, according to Weibull analysis ( $P < 0.01$ ).

**CONCLUSION:** Adhesive surface roughness influences streptococcal adhesion forces.

### MATHEMATICAL APPRAISAL OF ARCH FORM IN SUBJECTS WITH CROWDING TREATED NON-EXTRACTION

**B Vande Vannet<sup>1</sup>**, P Cara<sup>2</sup>, G Ibel<sup>1</sup>, <sup>1</sup>SOPA and <sup>2</sup>DWIS, Vrije Universiteit Brussel, Belgium

**AIM:** To evaluate tooth changes, with appraisal of arch form, pre- and post-treatment of 22 patients with severe upper and lower crowding treated with the Damon® system (D3, Ormco, Orange, California, USA) without any extractions or approximal enamel reduction.

**MATERIALS AND METHOD:** The digital models obtained were analyzed with the OrthoProof® software (Nieuwegein, Netherlands) to investigate tooth movements. On each tooth at least one reference point was located before treatment. Through these points a curve was then fitted. After treatment a new curve was fitted through the new positions of the reference points. Movement of the teeth was measured by numerically computing the area between the two curves using the octave numerical package ([www.gnu.org/software/octave](http://www.gnu.org/software/octave)) in order to estimate expansion. This was done by numerically integrating the difference between the two spline interpolated curves whose origin is centred between the two anterior incisors. Statistical analysis was performed using Statistical Package for Social Sciences, version 15.0 (SPSS Inc., Chicago, Illinois, USA).

**RESULTS:** Significant increases in upper arch length and width were noted after correction of crowding. Both molar distalization and incisor protrusion contributed to the increase in arch length. A mean expansion of 1.08 (8%), 1.04 (lower) and 1.12 (upper) was calculated.

**CONCLUSION:** During correction of upper arch crowding using the non-extraction Damon® technique, arch perimeter gain is due to an increase in both arch length and posterior arch width. Arch form could be reconstructed using the above method based on the given parameters.

### RATE OF DISTALIZATION IN CLASS II SUBJECTS USING HIGH PULL HEADGEAR

**W Stós**, J Panuszka-Landowska, Department of Orthodontics, Jagiellonian University Collegium Medicum, Krakow, Poland

**AIM:** To evaluate the amount of distalization of the upper molars in patients with crowding using high pull headgear.

**SUBJECTS AND METHOD:** Ten children with an Angle Class II malocclusion aged from 11 to 13 years with insufficient space in the lateral part of upper dental arch. The facebow was adjusted so that the point of the acting force would go through the centre of resistance of the first upper molar – as close as possible to trifurcation of its roots. The level of the measured force was 450 g per side. The patients used the headgear for 12-14 hours per day. Control visits with measurement of the acting force were conducted at four to six week intervals. Active treatment was for a period of five to seven months. An untreated control group with the same malocclusion was evaluated. Dental casts were made and comparable measurements were carried out before and after treatment for the following distances: between the mesial side of the first upper molar to the incisive papilla on left and right side; between mesial side of the upper first molar to mesial angle of central incisor on left and right side and between mesial side of the first upper molar to distal side of the lateral incisor in left and right side respectively.

**RESULTS:** The average increase in the distance between the mesial of the upper first molar to the incisive papilla was 1.27 mm. For the mesial angle of the central incisor the average value was 1.73 mm. For the distal side of the lateral incisor the average distance was about 2.33 mm.

**CONCLUSION:** High pull headgear is an alternative solution to other methods of distalization.

### CERVICAL VERTEBRAL MATURATION STAGES AND CALCIFICATION OF THE MANDIBULAR SECOND MOLAR

W Stós, M Bacheleda-Księżdzularz, B Remisz, M Mazurkiewicz-Janik, B W Loster Department of Orthodontics, Jagiellonian University Collegium Medicum, Krakow, Poland

**AIMS:** To investigate the relationship between the calcification stages of the mandibular second molar and skeletal maturity based on analysis of panoramic and cephalometric radiographs, respectively. A further aim was to determine if panoramic radiographs might be valuable for assessment of mandibular growth potential.

**MATERIALS AND METHOD:** Cephalometric and panoramic radiographs of 260 subjects (158 females, 102 males) ranging in age from 10 to 14 years. In each case the cervical vertebral maturation (CVM) and dental maturation stages of the lower second molar were determined. The frequency of occurrence of each of the six cervical stages (CS1-CS6) in both the female and male group and the corresponding dental calcification stage of the mandibular second molar was recorded.

**RESULTS:** CS3 and CS4, comprising the interval of maximum mandibular growth, were observed in 51 per cent of females and 31 per cent of males. In the female sample, CS3 was correlated with stage F of molar calcification (51%), CS4 was related to F or G stage (37 and 44%, respectively). For males CS3 was related to F or G stage of molar calcification (47 and 35%) and CS4 to stage G or H (61 and 26%). Molar calcification stage F in the female group was correlated with CVM stages CS3 and CS4 (64%) and stage G to CS4 and CS5 (72%). In the male group dental stage F was related to CVM stage CS1 and CS2 (72%) and stage G to CS2, CS3 and CS4.

**CONCLUSION:** Mandibular second permanent molar calcification stages on panoramic radiographs may be a useful indicator of the pubertal growth spurt. For the majority of female subjects dental calcification stage E was observed before, and stage G after the pubertal growth spurt. For males dental stages E and F occurred mostly before the growth peak.

### MICRODAMAGE IN THE ALVEOLAR PROCESS AFTER ORTHODONTIC TOOTH MOVEMENT

T M Präger<sup>1</sup>, P Meyer<sup>1</sup>, R J Radlanski<sup>2</sup>, R Müller-Hartwich<sup>1</sup>, P-G Jost-Brinkmann<sup>1</sup>, Departments of <sup>1</sup>Orthodontics, Dentofacial Orthopedics and Pedodontics and <sup>2</sup>Experimental Dentistry, Charité - Universitätsmedizin Berlin, Germany

**AIM:** Microdamage plays a decisive role in bone remodelling and has recently been described in the pig in combination with orthodontic tooth movement (Verna and Melsen). The aim of this study was to investigate the maxillary alveolar process of the rat for microdamage after orthodontic tooth movement and its relationship with orthodontic force application.

**MATERIALS AND METHOD:** In 24 male Wistar rats (10 weeks old) all three maxillary molars on the right side were moved mesially by means of a nickel titanium coil spring that was fixed to the upper incisors and delivered a force of 25 cN. The rats were randomly allocated to four groups of six animals and forces were applied for 1, 2, 4 or 7 days. The left side served as a control. After cautious preparation, the specimens were stained *en bloc* with basic fuchsin and slices cut parallel to the mesial root of the first molar. The specimens were examined under a microscope with transmission and epifluorescence light for determination of microdamage. In addition, the number of microcracks and their length was determined and differences were examined for significance by means of the Wilcoxon test ( $P < 0.05$ ).

**RESULTS:** Microdamage could be detected as diffuse microdamage and as microcracks on both sides of the maxilla and in all animal groups. The average length of the microcracks ranged between 47 and 70  $\mu\text{m}$ , and their density between 0.7 and 1.6  $\text{mm}^2$

microcracks. Although the absolute number of microcracks was more frequent on the experimental side, there was no significant difference, either between the sides or between the groups.

**CONCLUSION:** Microdamage represented as microcracks and diffuse microdamage can be found in the maxillary alveolar process independent of orthodontic loading. A significant accumulation of microcracks on the pressure side could not be confirmed under these experimental conditions.

### EFFECT OF MINIMAL OSTEOPERFORATIONS ON BONE REMODELLING AND TOOTH MOVEMENT

I Chartres<sup>1,2,3</sup>, J Tran<sup>1,2</sup>, E Khoo<sup>1,2</sup>, Y Liu<sup>1,2,3</sup>, M Alikhani<sup>1,2</sup>, <sup>1</sup>Consortium for Translation Orthodontic Research <sup>2</sup>New York University, USA <sup>3</sup>Universitätsklinikum Freiburg, Germany,

**AIM:** It has been shown that orthodontic forces induce aseptic inflammation but the relationship between the magnitude of inflammatory response and the rate of bone remodelling and consequently tooth movement is not clear. The objective of this research was to investigate whether an increase in local inflammation affects the rate of bone remodelling and tooth movement.

**MATERIALS AND METHOD:** Forty-eight rats divided into four groups: a baseline, and three groups fitted with a 50 cN force closing coil nickel titanium spring on the maxillary first molar with implementation of a soft tissue flap, force plus flap plus three perforations of the cortical plate mesial to the first molar, or no force (controls C). After 28 days, the maxillae were analyzed by microcomputed tomography ( $\mu\text{CT}$ ) and histology. RNA isolated at different time points was used to evaluate the expression of inflammatory markers by real-time reverse transcription polymerase reaction.

**RESULTS:** Local perforation of cortical bone caused an increased inflammatory reaction as shown by both histology and expression of inflammatory cytokines. This was accompanied by an increase in the rate of tooth movement and bone remodelling as revealed by  $\mu\text{CT}$  and fluorescent microscopy. The increase in the rate of bone remodelling was not limited to the first molar region and spread to adjacent alveolar bone.

**CONCLUSION:** An increase in magnitude of inflammation is accompanied by an increase in the rate of bone remodelling and tooth movement.

### HUMAN FOETUS CADAVERS WITH A CLEFT LIP AND PALATE IN THE THIRD TRIMESTER: CASE REPORT

E Esenlik<sup>1</sup>, E H Evcil<sup>2</sup>, M A Malas<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Anatomy, Süleyman Demirel University, Isparta, Turkey

**AIM:** A cleft lip and/or palate (CLP) is one of the common forms of congenital anomalies in newborns. It can be associated with syndromes such as anencephaly. The aim of this study was to morphometrically analyze the craniofacial form and intraoral region of three human foetus cadavers in the third trimester with CLP.

**SUBJECTS AND METHOD:** Three human cadavers retrieved from the archives of Süleyman Demirel University. The first, a male, showed anencephaly and a unilateral CLP at 29 weeks gestation. The second, a female, showed anoculi and an isolated cleft lip and alveolus at 37 weeks gestation, and the last, also female, had a total bilateral CLP at 37 weeks gestation. Crown-rump length, biparietal diameter, head circumference and femur length were measured as general parameters. Vertical and transverse lengths were measured extraorally. Trays made of acrylic were prepared for intraoral evaluation. Upper and lower casts were obtained and alveolar arch width, length, and amount and shape of the cleft were evaluated.

**RESULTS:** In the anencephalic foetus, the mouth, lips, face and eyes appeared abnormal. The lack of cephalic tissue resulted in a failure in formation of the cranium. Macroglossy was one of the main anomalies of the intraoral region. In the second foetus with

anocui restricted development of the middle and upper face region was the main atypical facial feature. Deformation of the hands and feet was the other external anomaly. The last foetus with bilateral CLP resembled a normal face except for the oral and subnasal regions.

**CONCLUSION:** Frequent disturbance in closure of the palate with craniofacial malformations is identified. These cases may be important for foetopathological examination and to help diagnose craniomaxillofacial growth pathology.

### SP 039

Wednesday, June 16 - at 10:45

#### STABILITY OF IMMEDIATELY LOADED MINISCREW IMPLANTS ACCORDING TO FORCE LEVELS\*\*\*

Y-M Jo<sup>1</sup>, Y-C Park<sup>1</sup>, K-J Lee<sup>1</sup>, J-Y Cha<sup>1</sup>, A Urban<sup>2</sup>, <sup>1</sup>Yonsei University, Seoul, South Korea and <sup>2</sup>Private office, Linges, Germany

**AIM:** To evaluate the initial stability in immediate loaded miniscrew implants loaded with different force levels.

**MATERIALS AND METHOD:** Two adult mongrel dogs (12 months of age) with an initial weight of 29 kg. Fifty-four newly designed titanium self-tapping miniscrew implants were implanted in the edentulous areas. The miniscrews were loaded with high and low forces in both the maxilla and mandible.

**RESULTS:** Initial stability was decreased in the first 3 weeks due to the weakening effects of bone remodelling. The high and low force groups did not show a significant difference with regard to failure rate, force magnitude, or mechanical and histological evaluation.

**CONCLUSION:** It is assumed that force is not the primary factor for miniscrew failure and future studies should evaluate other factors or combinations of factors as reason for miniscrew failure.

### SP 040

Wednesday, June 16 - at 10:45

#### MECHANICAL PROPERTIES OF NICKEL-TITANIUM ARCHWIRES AFTER USE IN THE ORAL ENVIRONMENT

M Sarul, J Antoszewska, B Kawala, Department of Dentofacial Orthopedics and Orthodontics Wrocław Medical University, Poland

**AIM:** Knowledge of the mechanical properties of nickel-titanium (NiTi) archwires is essential for management of orthodontic therapy with fixed appliances. Loading of periodontal tissues with too heavy forces may result in either complications or delayed tooth movement. The properties of orthodontic archwires may be affected by the temperature of the oral cavity, which is usually significantly decreased in mouth-breathers; these properties may also change permanently in the oral cavity. The aim of this research was to assess the activation and deactivation forces of NiTi archwires.

**MATERIALS AND METHOD:** Thirty samples of Titanol superelastic, copper NiTi 35°C and NeoSent alloy, diameter and cross-section equal to 0.016 and 0.016 X 0.022 inch, were tested after 4-6 weeks orthodontic application *in vivo*. For laboratory evaluation, a three-point bending test at 30°C was used. The control group consisted of 90 as-received archwires.

**RESULTS:** The distribution of activation and deactivation forces showed a comparable tendency. Independent of diameter or cross-section, the values gradually increased during activation and decreased during deactivation. They were ranked in decreasing sequence: Titanol superelastic, NeoSent alloy, copper NiTi 35°C. The deactivation forces increased in the 0.016 X 0.022 NeoSent alloy and 0.016 Titanol superelastic and decreased in the 0.016 X 0.022 Titanol superelastic. The results did not allow exact definition of the mode of changes in deactivation forces evaluated after orthodontic application.

**CONCLUSION:** As the mechanical properties of *in vivo* applied archwires change unpredictably, long-term research is required to establish their clinical use in the oral environment. The results of this study showed that archwires with a round diameter release deactivation forces optimal for orthodontic displacement.

### SP 041

Wednesday, June 16 - at 10:45

#### THERMOCYCLING EFFECTS ON THE BOND STRENGTH OF BRACKETS BONDED WITH DIFFERENT LIGHT SOURCES

E Çerekja, B Çakirer, Department of Orthodontics, Marmara University, Istanbul, Turkey

**AIM:** To compare the effect of thermocycling on the shear bond strength (SBS) of brackets and to evaluate the bond failure sites after debonding.

**MATERIALS AND METHOD:** Two hundred and forty premolar teeth divided into six equal groups. The brackets (3M/Unitek) were bonded with Transbond XT. Group 1 was cured with a normal halogen curing light for 40 seconds, groups 2, 3 and 4 were cured for 2, 3 and 6 seconds, respectively, with a high-power halogen light (Swiss Master Light) and groups 5 and 6 with a 3rd generation light emitting diode Bluephase (Ivoclar) for 10 and 20 seconds, respectively. After storage for 24 hours in the dark at 37°C in water, half of the samples were thermocycled in water baths at 5°C and 55°C with a dwell time of 30 second and a transfer time between dwells of 4 seconds for 7.500 cycles.

**RESULTS:** Thermocycling significantly reduced the SBS of group 4 while the decreases in other groups were not significant. Adhesive remnant index scores were significantly different between the groups.

**CONCLUSION:** A high-power halogen light with 2 seconds curing time is not recommended.

### SP 042

Wednesday, June 16 - at 11:30

#### EFFECT OF OESTROGEN AND LOADING TO THE AMOUNT OF OESTROGEN RECEPTOR ALPHA AND MATRIX METALLOPROTEINASE-8 IN RAT CONDYLAR CARTILAGE

M Orajärvi<sup>1</sup>, O Hirvonen<sup>1</sup>, P Tiilikainen<sup>2</sup>, A Raustia<sup>2</sup>, P Pirttiniemi<sup>3</sup>, <sup>1</sup>Institute of Dentistry, and Departments of <sup>2</sup>Prosthetic Dentistry and Stomatognathic Physiology and <sup>3</sup>Oral Development and Orthodontics, University of Oulu, Finland

**AIM:** Changes in oestrogen and temporomandibular joint (TMJ) loading levels are thought to be involved in degradation of condylar cartilage. It has been suggested that matrix degradation caused by matrix metalloproteinase (MMP) is a primary event in joint diseases. The aim of this study was to examine the effect of decreased oestrogen level and altered dietary loading on condylar cartilage in the rat TMJ on the expression of oestrogen receptor alpha (ER-α) and MMP-8.

**MATERIALS AND METHOD:** Thirty-six female rats were divided into four groups: ovariectomized rats fed with normal diet, non-ovariectomized controls fed with normal diet, an ovariectomized soft diet group, and a non-ovariectomized control soft diet group. Ovariectomy was performed at the age of 60 days. Seven days after surgery the rats were sacrificed and prepared for immunohistochemical staining.

**RESULTS:** The proportional amount of MMP-8 positive cells was statistically significantly higher in the control rats fed with a normal diet than in control rats fed with a soft diet ( $P < 0.05$ ). The proportional amount of ER-α was statistically significantly higher ( $P < 0.001$ ) in the condylar cartilage of ovariectomized rats than in non-ovariectomized control rats both in the normal and soft diet groups. No statistically significant difference was found in ER-α staining when comparing the control rats fed a normal diet and the control rats fed with a soft diet.

**CONCLUSION:** Condylar cartilage is sensitive to changes in oestrogen level and loading. It seems that the expression of MMP-8 in the condylar cartilage is chiefly dependent on the level of loading. The expression of ER-α is dependent mainly on oestrogen level, not on the level of dietary loading.

**SP 043**

Wednesday, June 16 - at 12:15

**INVESTIGATION OF THE MOST PREFERRED MOUTH SHAPE DURING SMILING IN JAPANESE. PART I\*\*\***

J Funaki, S Funaki, Funaki Orthodontic Clinic, Tokyo, Japan

**AIM:** To identify the most preferred smiling mouth shape in Japan.**MATERIALS AND METHOD:** The different smiling mouth shapes were classified into 12 smile-patterns based on four positions of the corners of the mouth and three types for visibility of the teeth and gums. In classifying the positions of the corners of the mouth, the highest position was A-type, followed by B, C and D types. Regarding the classification of visibility of teeth and gums, in type 1 the upper gum and anterior teeth were visible, in type 2 the upper and lower anterior teeth were visible, and in type 3 the lower gum and lower anterior teeth were visible. The questionnaire was completed by 92 female patients and their mothers, aged from 9 to 51 years. The 12 smile-patterns were classified into four groups based on an evaluation criteria scale of three points for the most desirable mouth shape down to 0 points for the least desirable.**RESULTS:** No significant differences were found in the preference of the smiling mouth shape amongst subjects aged under 19 years and those over 20 years. For the position of the corners of the mouth, A-type was ranked highest, followed by B, C and then D types. For visibility of the teeth and gums, type 2 was ranked highest, followed by type 1 and then type 3. Significant differences ( $P < 0.01$ ) were found both in the position of the corners of the mouth and visibility of the teeth and gums. In the ranking of the 12 smile-patterns for the most desirable smiling mouth shape, the most desirable was the combination of A-type and type 2, characterized by the highest positions of the corner of the mouth with visible upper and lower anterior teeth.**CONCLUSION:** It seems Japanese people prefer a balanced smile where both upper and lower teeth are visible, showing minimal gingiva and with the corners of the mouth in the highest position.**SP 044**

Wednesday, June 16 - at 12:15

**INVESTIGATION OF THE MOST PREFERRED MOUTH SHAPE DURING SMILING IN JAPANESE. PART II CHANGES AFTER ORTHODONTIC TREATMENT\*\*\***

J Funaki, S Funaki, Funaki Orthodontic Clinic, Tokyo, Japan

**AIM:** To examine the changes in the smiling mouth shape before and after orthodontic treatment.**MATERIALS AND METHOD:** Two hundred and eighty two frontal photographs and lateral cephalograms taken before and after orthodontic treatment of 141 orthodontic patients (44 males, 97 females, average age 18 years 6 months). The smiling mouth shapes frontal photographs were classified into 12 smile-patterns (Part I of this study). Each photograph received a score (smile score) in accordance with the ranking of desirable smiling mouth shape from 12 smile-patterns (12 points for the highest ranked smile pattern and 1 point for the lowest ranked smile pattern). The increase or decrease of the smile score following treatment was examined to evaluate improvement or deterioration of the smiling mouth shape. The lateral cephalograms were used to examine the change in the occlusion represented by dental patterns (overjet and overbite) as well as skeletal patterns (AB difference) before and after treatment.**RESULTS:** After treatment the average score for the smile types increased significantly from 6.3 to 8.6. Compared with males (increase from 5.5 to 7.5), females (increase from 6.6 to 9.1) showed a higher increase in score. A significant relationship was found between the occlusion and the visibility of the gums and anterior teeth. However, the occlusion did not have a significant effect on the position of the corners of the mouth, nor on the smile pattern.**CONCLUSION:** The smiling mouth shape improved significantly after orthodontic treatment both in the male and female groups, with females expressing more interest in their smiles than males. To some extent orthodontic treatment may contribute to im-

provement in the smile. However, improvement may also come from increased confidence as a result of orthodontic treatment.

**SP 045**

Wednesday, June 16 - at 16:00

**THE EFFECTS OF A FACEMASK/BITE BLOCK THERAPY WITH OR WITHOUT RAPID PALATAL EXPANSION**

C Pavoni, F Ballanti, R Lione, Department of Orthodontics University of Rome Tor Vergata, Italy

**AIM:** There are a number of treatment approaches regarding orthopaedic treatment of Class III malocclusions. To control the sagittal discrepancy, the use of the protraction facemask (FM) with or without rapid maxillary expansion (RME) for treating Class III malocclusion has gained popularity. Control of the vertical skeletal relationships is important in orthopaedic treatment of Class III malocclusions with a FM. It has been shown that avoiding backward rotation of the mandible is crucial to prevent relapse after early orthopaedic Class III treatment. The aim of this study was to compare the effects of a FM in combination with a mandibular bite block (FM/BB) or with RME as a bonded device (RME/FM) in patients with Class III malocclusion.**SUBJECTS AND METHODS:** The FM/BB sample included 22 subjects [12 girls, 10 boys, average age before treatment (T1)  $8.7 \pm 1.2$  years, mean age after active treatment (T2)  $10.4 \pm 1.3$  years, mean treatment duration  $1.7 \pm 0.8$  years]. The RME/FM sample comprised 17 subjects (10 girls, 7 boys, average age at T1,  $7.8 \pm 1.8$  years and at T2,  $9.3 \pm 1.9$  years). The mean duration of observation was  $1.5 \pm 0.6$  years. Lateral cephalograms obtained at T1 and T2 were evaluated. T2-T1 changes in the two groups were compared with an independent sample t-test ( $P < 0.05$ ).**RESULT:** Comparison of the two treatment protocols showed that there were no significant differences for any measurements in either the sagittal or vertical planes from T1 to T2.**CONCLUSION:** FM in combination with different types of full-coverage occlusal splints results in similar dentoskeletal outcomes regardless of the presence or absence of RME. Both the BB appliance and the splinted RME limited posterior rotation of the mandible.



## Contemporary Diagnostics and Treatment Approaches

SP 046

Wednesday, June 16 - at 8:30

**WHICH FACIAL PROFILES MOSTLY DISCRIMINATE PERCEPTION OF MALE AND FEMALE ADOLESCENTS?**

**S Špalj<sup>1</sup>**, M Šlaj<sup>2</sup>, <sup>1</sup>Department of Pediatric Dentistry and Orthodontics, University of Rijeka and <sup>2</sup>Department of Orthodontics, University of Zagreb, Croatia

**AIM:** Adolescence is the age of sexual development and of seeking partners, and perception of facial attractiveness plays an important role in the process. Therefore orthodontist must be aware of facial profile preferences as therapy can alter the attractiveness of the lower facial third. The aim of this research was to establish which facial profiles mostly discriminate perception of male and female adolescents.

**SUBJECTS AND METHOD:** A random sample of 1089 adolescent (450 males, 639 females) aged 15-19 years ( $16.3 \pm 0.4$ ) from 12 secondary schools in Zagreb, Croatia. Eight profile distortions for each gender, morphed by a digital imaging technique, were assessed for attractiveness by each adolescent and scored from 1 to 8. Multivariate discriminant analysis was used for statistical evaluation.

**RESULTS:** In general male and female adolescents had similar facial profile preferences. A straight profile was considered the most attractive for both genders. Both bimaxillary alveolar protrusive and retrusive profile were equally attractive among male profiles, but for female profiles thicker rather than flat lips were significantly more attractive. An extreme convex profile was the least attractive, but a mild convex shape was more preferable than a concave shape. According to canonical discriminant function, males and females were mostly discriminated in attractiveness scores of bimaxillary alveolar retrusion and mandibular retrognathism in female profiles, and a combination of maxillary prognathism and mandibular retrognathism in both male and female profiles. Compared with males, females were less tolerant of retrusive lips in females and convex profiles in both males and females.

**CONCLUSION:** Due to facial profile preferences in borderline cases, it would be wise to treat female patients non-extraction, while for male patients the decision regarding extractions can be more easily decided.

SP 047

Wednesday, June 16 - at 8:30

**SKELETAL CHANGES INDUCED BY RAPID MAXILLARY EXPANSION ASSESSED WITH SUTURAL DENSITY ANALYSIS ON LOW-DOSE COMPUTED TOMOGRAPHY**

**L Franchi<sup>1</sup>**, R Lione<sup>2</sup>, P Cozza<sup>2</sup>, <sup>1</sup>Università degli Studi di Firenze and <sup>2</sup>Università degli Studi di Roma, Italy

**AIM:** The measurement of sutural density represents a contemporary approach to determine the outcome of orthopaedic intervention on the transverse dimensions of the maxilla and ideal timing for retention of maxillary expansion. The aim of this research was to evaluate the density of the midpalatal suture as assessed by low-dose computed tomography (CT) before rapid maxillary expansion (RME, T0), at the end of active expansion (T1) and after a retention period of 6 months (T2).

**SUBJECTS AND METHOD:** Seventeen prepubertal subjects (mean age 11.2 years) with constricted maxillary arches, and uni- or bilateral posterior crossbites. The total amount of expansion was 7 mm in all subjects. Multi-slice low-dose CT scans were taken at T0, T1, and T2. On axial CT scanned images four regions of interest (ROIs) were placed along the midpalatal suture (anterior, AS ROI, and posterior, PS ROI) and in two regions of palatal bone (anterior and posterior). Density was measured in Hounsfield units. Mann-Whitney U test and Friedman ANOVA with *post-hoc* test were used ( $P < 0.05$ ).

**RESULTS:** The densities in the AS and PS ROIs were significantly smaller than reference bone densities before RME. Both AS and PS ROIs showed a significant decrease in density from T0 to T1, a significant increase from T1 to T2, and a lack of a significant difference from T0 to T2.

**CONCLUSION:** Effective opening of the midpalatal suture by RME in prepubertal subjects was associated with a significant decrease in sutural density. Sutural density at T2 indicated reorganization of the midpalatal suture.

SP 048

Thursday, June 17 - at 8:30

**TREATMENT PLANNING FOR MAXILLARY CANINE DISPLACEMENT BASED ON INDIVIDUAL SKELETAL MATURITY**

**V Giuntini**, T Baccetti, L Franchi, Università degli Studi di Firenze, Italy

**AIM:** In treatment planning for a palatally displaced canine it is important to determine the timing of potential eruption of the canine (when interceptive procedures can be attempted) versus impaction of the tooth. Thus the aim of this study was to assess the relationship between the eruption of the permanent maxillary canine and individual skeletal maturity in subjects with different skeletal relationships in the sagittal and vertical planes.

**SUBJECTS AND METHOD:** One hundred and fifty two subjects (63 males, 89 females) with erupting permanent maxillary canines. The individual stage of cervical vertebral maturation (CVM) was assessed on lateral cephalograms. Following this analysis, the subjects were divided into pre-peak (before the pubertal growth spurt, CS1 and CS2), peak (during the pubertal growth spurt, CS3 and CS4), and post-peak (after the pubertal growth spurt, CS5 and CS6) groups. Skeletal relationships in the sagittal and vertical planes were evaluated, and relationships to timing of canine eruption were statistically tested.

**RESULTS:** The pre-peak group comprised 86 subjects, the peak group 66 subjects. No subjects were classified as post-peak. The differences in prevalence rates between the pre-peak and peak groups were statistically significant ( $P < 0.001$ ). The prevalence rate for hyperdivergent subjects showing eruption of the permanent maxillary canine in the pre-peak group (37.2%) was significantly higher than in the reference orthodontic population (21%).

**CONCLUSION:** Eruption of the permanent maxillary canine can occur at any stage during skeletal maturation before the end of the pubertal growth spurt (CS1 through to CS4), with hyperdivergent subjects showing more frequently pre-pubertal canine eruption.

**SP 049**

Thursday, June 17 - at 8:30

**A DIAGNOSTIC PERFORMANCE STUDY ON OVERJET IN CLASS II MALOCCLUSION SUBJECTS****A Vangelisti**, V Giuntini, C Masucci, Università degli Studi di Firenze, Italy**AIM:** To evaluate the associations between an increased overjet and other dentoskeletal characteristics of Class II division 1 malocclusion subjects in the mixed dentition.**SUBJECTS AND METHOD:** Nine hundred mixed dentition subjects observed by clinical inspection, analysis of dental casts, and lateral cephalograms. Overjet used as the diagnostic indicator (overjet  $\geq 7$  mm) was evaluated in relation to other Class II dentoskeletal features (Class II molar and canine relationships, and skeletal Class II relationship). Diagnostic performance was assessed by odds ratio and positive likelihood ratio within a Bayesian statistical approach.**RESULTS:** The diagnostic performance of overjet with regard to the other dentoskeletal components of Class II malocclusion was not significant.**CONCLUSION:** When used as an isolated occlusal feature, overjet is not a valid diagnostic indicator of a Class II division 1 malocclusion. Discrimination between clinical conditions showing an isolated overjet from a comprehensive Class II malocclusion during diagnosis, treatment planning, and the analysis of treatment outcomes is recommended.**SP 050**

Thursday, June 17 - at 8:30

**NON-CONVENTIONAL BRACKETS VERSUS LIGATURES FOR THE ALIGNMENT OF BUCCALLY DISPLACED TEETH****A Vangelisti**, T Baccetti, L Franchi, Università degli Studi di Firenze, Italy**AIM:** To analyze the forces released by four types of passive stainless steel self-ligating brackets and by two non-conventional elastomeric ligature-bracket systems when compared with conventional elastomeric ligatures on stainless steel brackets during alignment of buccally displaced teeth.**MATERIALS AND METHOD:** A model consisting of five brackets (from second premolar to the central incisor) was used to assess the forces released by the seven different bracket-ligature systems with 0.012 or 0.014 inch superelastic wires in the presence of various amounts of buccal canine displacement (1.5 to 6.0 mm). Comparisons between the different types of bracket/wire/ligature systems were performed with three-way ANOVA with Tukey's *post hoc* test ( $P < 0.05$ ).**RESULTS:** For buccal misalignments of 1.5 and 3.0 mm, both low-friction and conventional systems released forces for bracket alignment ranging from ~30 to ~160 g. In presence of a large amount of buccal displacement (4.5 or 6.0 mm), the low-friction systems produced significant force, while this dropped to 0 g for the conventional system.**CONCLUSION:** Non-conventional elastomeric ligature-bracket systems produce levels of force available for tooth movement that are similar to those generated in the presence of passive self-ligating brackets.**SP 051**

Thursday, June 17 - at 8:30

**THREE-DIMENSIONAL LOCALIZATION OF IMPACTED MAXILLARY CANINES AND EFFECTS ON ADJACENT INCISORS****K Mina**<sup>1</sup>, S Shahidi<sup>2</sup>, B Zamiri<sup>3</sup>, <sup>1</sup>M.O.G and Departments of <sup>1</sup>Radiology and <sup>3</sup>Surgery, Shiraz Dental School, Iran**AIM:** Traditionally, many different types of radiographs have been used to localize an impacted tooth in relation to the adjacent teeth. Although these views allowed identification as to whether the tooth was buccal or palatal (lingual), it was still not possible to measure exactly how far away they were impacted from the roots of the other teeth. Cone beam computed tomography (CBCT)

provides elements that cannot be observed during traditional radiographic analysis and is, therefore, indicated in the case of impacted teeth or craniofacial structural anomalies. The aim of this study was to evaluate CBCT for three-dimensional localization of impacted maxillary canines and the effects on adjacent incisors.

**MATERIALS AND METHOD:** Sixty impacted maxillary canines assessed regarding location, shape (with or without dilacerations), inclination, follicular size and effects on the roots of neighbouring teeth.**RESULTS:** The impaction location was 25 per cent (15 cases) buccally, 63.3 per cent (38 cases) palatally, and 11.66 per cent (7 cases) in the middle of the bone. Twenty-five canines (41.66%) had dilacerated roots. For 25 canines (41.66%) the follicle size was normal, 23 canines (38.33%) had a large follicle, and 12 canines (20%) a diminished follicle. Adjacent root resorption was mild in 25 per cent (15 cases), moderate in 5 per cent (3 cases) and severe in 11.66 per cent (7 cases). No resorption was observed in 58.33 per cent (35 cases).**SP 052**

Thursday, June 17 - at 8:30

**CRANIOFACIAL MORPHOLOGY OF CHINESE ADULTS WITH NORMAL OCCLUSION AND SKELETAL DIVERGENCE****D Xiao**<sup>1,2</sup>, **H Gao**<sup>2</sup>, **Y Ren**<sup>1</sup>, Departments of Orthodontics, <sup>1</sup>University Medical Centre Groningen and University of Groningen, Netherlands and Stomatological Hospital, NanKai University, China**AIM:** To examine the craniofacial morphologic characteristics of different vertical dysplasias in a population of Chinese adults with normal occlusion.**SUBJECTS AND METHOD:** Sixty-nine subjects with normal occlusion were selected from 800 healthy students between 18-24 years of age. Lateral cephalograms were obtained and 27 hard and 10 soft tissue measurements were analyzed. The subjects were then divided into three groups of high angle, low angle or control according to the value of FH-MP. Intraclass correlation coefficient was determined for the repeated measurements. One-way ANOVA was used to determine the differences between the groups.**RESULTS:** The low angle group had a larger cranial basis angle (N-S-Ar) and the high angle group a shorter maxilla (Ans-Ptm) ( $P < 0.01$ ). The high angle group displayed vertical hyperdivergency with increased PP-OP, OP-MP, gonial and lower gonial angles, whereas the low angle group showed significant hypodivergency with decreased values for all variables ( $P < 0.01$ ). The low angle group had a more protrusive chin and the high angle group a more retrusive chin ( $P < 0.01$ ). Differences in dentoalveolar measurements in the divergent groups were mainly in the anterior region. Moreover, the low angle group had a thicker, and the high angle group, a thinner lower dentoalveolus ( $P < 0.01$ ). For face height measurements, the main differences in the divergent groups were at the anterior lower third ( $P < 0.01$ ). Soft tissue deviations were less obvious in the high angle group and in general less significant than those of the hard tissues in both divergent groups.**CONCLUSION:** Significantly different morphological characteristics exist in Chinese adults with vertical dysplasia but normal occlusion. Major skeletal cephalometric changes existed at the lower third of the face. The soft tissues showed a well adapting mechanism of soft tissue coverage for the skeletal dysplasia.

**SP 053**

Thursday, June 17 - at 8:30

**RELATIONSHIP BETWEEN UPPER AIRWAY WIDTH AND FACIAL GROWTH CHANGES DURING ORTHODONTIC TREATMENT OF GROWING CHILDREN****Y-H Park<sup>1</sup>**, Y-J Kim<sup>1</sup>, J-S Hong<sup>1</sup>, K-H Lee<sup>2</sup>, <sup>1</sup>Kangdong Sacred Heart Hospital, Hallym University Medical Center and <sup>2</sup>Pocheon Public Health Center, Korea**AIM:** In growing patients with malocclusions, functional assessment such as respiration is vital for the restoration of normal craniofacial growth. The purpose of this study was to evaluate how upper airway changes influence facial growth during puberty in Korean children.**SUBJECTS AND METHOD:** Thirty-six patients aged 9 to 11 years (mean: 10.7 years). Cephalograms and hand-wrist radiographs were taken before treatment (T1) and after appliance removal (T2). The sample was allocated to narrow (5.2-8.6 mm), medium (8.9-11.5 mm), and wide (11.7-16.0 mm) upper airway groups according to the airway width at T1. Cephalometric measurements at T1 and T2 and growth from T1 to T2 were compared among the groups.**RESULTS:** The increase in airway size in each group was 4.55, 3.84 and 1.94 mm in the narrow, medium, and wide groups, respectively, and the differences were statistically significant. The posterior face height of the subjects in the narrow group was found to be significantly smaller than in the other two groups at T1. For growth changes from T1 to T2, the narrow group showed significantly larger values of PFH/AFH, facial axis, ANS, and Gn than the medium group. Compensatory changes in airway width and facial growth were found in the subjects with a narrow upper airway.**CONCLUSION:** In young children with a constricted upper airway, surgical intervention of adenoid and tonsillar hypertrophy should be carried out with reservations, since these airway problems could improve naturally.**SP 054**

Thursday, June 17 - at 8:30

**TWO SCORING SYSTEMS FOR TREATMENT EVALUATION OF BILATERAL CLEFT LIP AND PALATE PATIENTS****T Bartzela<sup>1</sup>**, C Leenarts<sup>1</sup>, E Bronkhorst<sup>2</sup>, C Katsaros<sup>3</sup>, A M Kuijpers-Jagtman<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and Oral Biology and <sup>2</sup>Cariology and Preventive Dentistry, Radboud University Nijmegen Medical Centre, Netherlands and <sup>3</sup>Department of Orthodontics and Dentofacial Orthopedics, University of Bern, Switzerland**AIM:** A complete bilateral cleft lip and palate (CBCLP) is an 'orphan' condition due to its low incidence. Therefore nearly all scoring systems that measure treatment outcome have been developed for unilateral clefts. However, it is essential to have reliable methods for assessment of treatment outcome in CBCLP, which involves treatment until facial growth has ceased. The aim of this study was to compare two scoring systems, i.e. the Huddart/Bodenham (HB) system and the newly developed BCLP yardstick, which classifies treatment outcome in terms of dental arch relationships in patients with CBCLP. In addition, a further aim was to evaluate the predictive value of these scoring systems for treatment outcome.**SUBJECTS AND METHOD:** Dental arch relationships of 44 CBCLP patients were evaluated at 6, 9, and 12 years using the HB-scoring system and the BCLP yardstick. In total 121 dental casts were rated. For each age group the HB-scores were correlated with the BCLP yardstick scores using Pearson's correlation coefficient. The predictive value of the two scoring systems was evaluated by backward regression analysis.**RESULTS:** Intra-observer kappa values for the BCLP-yardstick scores for the two observers were 0.506 and 0.627, respectively, and interobserver reliability ranged between 0.427 and 0.581. Intra-observer reliability for the HB-system ranged from 0.92 to 0.97 and interobserver reliability from 0.88 to 0.96. The BCLP-yardstick scores at 6 and 9 years together were predictors for the outcome

at 12 years (explained variance 41.4%). Adding the incisor and lateral HB-scores in the regression model increased the explained variance to 67 per cent.

**CONCLUSION:** The BCLP-yardstick and the HB-system are reliable scoring systems for evaluation of dental arch relationships of CBCLP patients. The HB-system categorizes treatment outcome into similar categories as the BCLP-yardstick. In case a more sensitive measure of treatment outcome is needed, selectively both scoring systems should be used.**SP 055**

Thursday, June 17 - at 8:30

**ADULT CLASS II DIVISION 1 HERBST TREATMENT – IS IT STABLE?****N Bock**, S Ruf, Department of Orthodontics, University of Giessen, Germany**AIM:** To assess the skeletal and dental changes occurring during the retention period after adult Class II division 1 Herbst treatment.**SUBJECTS AND METHOD:** Fifteen Class II division 1 subjects (overjet >6 mm) with a minimum age of 18 years who completed growth according to pre-treatment hand wrist radiographs. All subjects were treated with a Herbst appliance (mean 9 months), followed by multibracket (MB) treatment (mean 14 months). The retention period was at least 24 months (mean 36 months). Lateral cephalograms (T1 before treatment, T2 after Herbst-MB treatment, T3 after retention) were analysed using the SO-analysis described by Pancherz, and standard cephalometrics.**RESULTS:** The amount of relapse during T3-T2 was small, amounting to 1.0 mm ( $P < 0.001$ ) for overjet and 0.3 mm ( $P > 0.05$ ) for molar relationship. The final relative contribution of skeletal changes to overjet and molar relation was lower by 6 and 12 per cent, respectively, compared with the active treatment period (T2-T1). During the entire observation period (T3-T1) the overjet was, on average, reduced by 5.2 mm ( $P < 0.001$ , minimum -2.5 mm, maximum: -8.0 mm; 12 per cent skeletal changes, 88 per cent dental changes) and the molar relation improved by an average of 3.2 mm ( $P < 0.001$ , minimum:  $\pm 0$  mm, maximum: -7.0 mm; 19 per cent skeletal changes, 81 per cent dental changes). During T3-T2 the jaw base relationship deteriorated (ANB +0.3°,  $P > 0.05$ ) and the hard as well as soft tissue profile convexities increased (NAPg -0.3°,  $P > 0.05$ ; N'SnPg' +0.6°,  $P > 0.05$ ) slightly. Nevertheless, during the entire observation period (T3-T1) the jaw base relationship improved (ANB -0.6°,  $P < 0.05$ , minimum: +0.5°, maximum: -2.8°), and the hard and soft tissue profile convexities decreased (NAPg +1.2°,  $P < 0.01$ , minimum: -1.3°, Max: +5.3°; N'SnPg' +2.1°,  $P < 0.01$ , minimum: -0.5°, maximum: +5.0°).**CONCLUSION:** Although Class II correction in adult Herbst-MB treatment is achieved by mainly dental changes, the treatment results showed good overall stability.**SP 056**

Thursday, June 17 - at 8:30

**EFFECT OF AUTOTRANSPLANTATION ON BONE REGENERATION IN ORTHODONTIC PATIENTS****E Hamza**, R W KWong, A B M Rabie, Prince Philip Dental Hospital, University of Hong Kong, SAR China**AIM:** Horizontally impacted second molars pose a problem to orthodontists. The proximity of the crown of the second molar to the distal root of the first molar can denude the first molar distal root of periodontal ligament and bone. The option of uprighting the second molar extends the treatment time considerably and does not address the periodontal defect on the distal root of the first molar. The aim of this study is to present a protocol for the management of molar stacking complicated by bony defects in adjacent teeth, and to present the effect of autotransplantation on bone regeneration around adjacent teeth.

**MATERIALS AND METHOD:** Fourteen autotransplanted teeth followed-up radiographically and clinically at 1 and 4 weeks, 3 and 6 months and 1 and 2 year intervals. Examination involved the amount of bone regeneration, mobility, gingival pocket depth, and vitality test. The criteria of success were stability of the transplanted teeth, bone regeneration around periodontally affected adjacent teeth and normal lamina dura formation, and normal mobility and gingival pocket depth of the transplanted teeth as compared with control teeth. Different statistical analysis was performed, including Wilcoxon signed ranks test and repeated measures ANOVA.

**RESULTS:** The survival rate over at least 2 years (range from 2 to 7%) after autotransplantation was 92.86 per cent. Bone regeneration at the socket site and adjacent teeth was significantly improved. Mobility and gingival pocket depths of the autotransplanted teeth were normal when compared with the normal control teeth.

**CONCLUSION:** Due to the high success rate and the advantages gained, autotransplantation should be considered as a treatment option for management of molar stacking complicated by bony defects in adjacent teeth.

### SP 057

Thursday, June 17 - at 8:30

#### ANTERIOR DENTOALVEOLAR MORPHOLOGY IN RELATION TO NATURAL HEAD POSTURE

**H Gokalp**, Development of Orthodontics, Faculty of Dentistry, University of Ankara, Turkey

**AIM:** To research the relationship between natural head posture and anterior dentoalveolar morphology.

**MATERIALS AND METHOD:** Fifty-two natural head positioned lateral cephalogram of subjects with a Class I malocclusion. Six cranio-cervical and 15 dentoalveolar variables were measured. The relationship between anterior dentoalveolar morphology and head posture was analyzed by Pearson correlation analysis.

**RESULTS:** Statistically significant correlations were found. Positive correlations were observed between the upper/lower dentoalveolar structures and head posture. Negative correlations existed between overbite and head posture.

**CONCLUSION:** Extension and flexion of the head affects the anterior dentoalveolar as well as craniofacial structures.

### SP 058

Thursday, June 17 - at 9:15

#### RELATIONSHIP BETWEEN ROOT RESORPTION AND DENTAL ANOMALIES

**K Van Parys**<sup>1</sup>, I Aartman<sup>2</sup>, R Kuitert<sup>1</sup>, A Zentner<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Social Dentistry and Behavioral Sciences, Academic Centre Dentistry Amsterdam, Netherlands

**AIM:** To determine the relationship between root resorption and the presence of tooth agenesis and pipette-shaped roots.

**MATERIALS AND METHOD:** The dental anomalies (agenesis and pipette-shaped roots) and root resorption were assessed on panoramic radiographs of 88 patients [27 males, 61 females, mean age 28.7 (SD = 11.3 years)]. The random selection of patients was based on a number of inclusion criteria (e.g. 15 years or older, no previous fixed appliance treatment, good quality panoramic radiographs and visibility of the periodontal ligament of every tooth, and edgewise fixed appliance treatment with a duration of 18 months or more). A pipette-shaped root was defined as drawn by Levander and Malmgren (1988). Agenesis of a tooth was assessed on the panoramic radiograph and from the patient's dental history. Root resorption was calculated as the difference between root length before and after treatment, with and without a correction factor (crown length post-treatment/crown length pre-treatment). If one of the four upper incisors showed root resorption of 2.3 mm or more with both formulas the patient was scored as having root resorption.

**RESULTS:** Chi-square tests indicated that there was no relationship between root resorption and agenesis ( $P = 0.885$ ) or between root resorption and pipette-shaped roots ( $P = 0.800$ ).

**CONCLUSION:** It was not possible to confirm a relationship between root resorption and dental anomalies, such as agenesis and pipette-shaped roots. Panoramic radiographs might be unsuitable for the assessment of root resorption. Levander E, Malmgren O 1988 Evaluation of the risk of root resorption during orthodontic treatment: a study of upper incisors. European Journal of Orthodontics 10: 30-38.

### SP 059

Thursday, June 17 - at 9:15

#### FIXED FUNCTIONAL TREATMENT VERSUS PREMOLAR EXTRACTIONS IN YOUNG ADULTS

**I Ilhan**, T Alcan, Department of Orthodontics, Faculty of Dentistry, Marmara University, Istanbul, Turkey

**AIM:** To compare dentoskeletal and profile changes following three treatment protocols: Jasper Jumper application (J), two upper premolar extractions (2P), and four premolar extractions (4P). **MATERIALS AND METHOD:** The initial and finishing cephalograms of 91 young adults (54 males, 37 females, 12-17 years of age) with a moderate Class II division 1 malocclusion (overjet  $\leq 5$  mm, ANB  $\leq 4^\circ$ , GoMe-SN 25-39°). The distribution of the subjects was 36 per cent for the J group, 31 per cent for the 2P group and 33 per cent for the 4P group. The mean ages in three groups were 14.25, 15.08, and 14.42 years, respectively. Thirteen skeletal, 16 dental, and five soft tissue parameters were analyzed, by means of hand-traced cephalograms. The method error was calculated 1 month later.

**RESULTS:** ANB decreased most in THE J group and least in the 2P group. In the J group, correction was due to maxillary restriction rather than mandibular skeletal growth. The extraction groups showed bimaxillary retrusion, mainly in the maxilla. The mandibular retrusive effect was more evident in the 2P group. Vertical pattern and lower anterior face height increased in all groups, and dental movements were in accordance with the treatment mechanics used. Profile convexity decreased in the J and 4P groups, while the 2P group showed an insignificant increase. Intergroup comparisons revealed no significant differences in maxillary position, vertical dimension, or occlusal plane changes. Significant differences were observed mainly in mandibular sagittal position, dental movements, and soft tissue profile.

**CONCLUSION:** The decision regarding extractions in young adults should be based mainly on initial mandibular position and profile type.

### SP 060

Thursday, June 17 - at 9:15

#### A PROTOCOL FOR IMPROVED STABILITY WITH HERBST APPLIANCE TREATMENT

**D F Austin**, University of Hong Kong, SAR China

**AIM:** To present a treatment protocol that leads to increased stability in the long term by comparing the published data on the short- and long- term effects of Herbst appliance treatment with single or multiple step mandibular advancement.

**SUBJECTS AND METHOD:** Sixteen Class II division 1 adults treated with Herbst appliances for 12 months with stepwise advancement, followed by fixed appliance. Lateral cephalograms were taken pre-treatment, after Herbst appliance treatment, after fixed appliance treatment and at the 3 year follow-up after debond. The results were compared with 16 Class II division 1 adolescents treated with a Herbst appliance for 7 months with maximal single step advancement, followed by retention with an Andresen activator. Lateral cephalograms were taken pre-treatment, after Herbst appliance treatment, after 6 months retention with the Andresen activator, and at the 2 year follow up after Andresen activator retention (Phan *et al.*, 2006).

**RESULTS:** Comparing the treatment effects of the two protocols demonstrated: (1) initially greater forward positioning of the mandible in the adult group; (2) less relapse after appliance removal in the adult group; (3) long term, greater stability in the adult group; (4) greater skeletal contribution to overjet and molar correction in the adolescent group.

**CONCLUSION:** Long-term stability of Herbst appliance treatment is affected by the treatment protocol employed. Based on the increased stability of the adult group, it is recommended to treat for 12 months in a stepwise manner for mandibular advancement for both adolescent and adult patients without retention between phase I and II treatment.

## SP 061

Thursday, June 17 - at 9:15

### EDENTULOUS RIDGE ALTERATIONS FOLLOWING ORTHODONTIC TOOTH MOVEMENT

**B Lindskog Stokland**<sup>1</sup>, A Ekestubbe<sup>2</sup>, J Wennström<sup>3</sup>, Departments of <sup>1</sup>Orthodontics, <sup>2</sup>Oral and Maxillofacial Radiology and <sup>3</sup>Periodontics, Sahlgrenska Academy, University of Gothenburg, Sweden

**AIM:** In patients with edentulous areas, prosthodontic rehabilitation may be problematic because of insufficient ridge dimensions. Orthodontic movement of teeth could be a treatment option to prepare for conventional prostheses or for implant installation. The available information regarding the effect of orthodontic tooth movement on ridge dimensions is limited. The aim of this research was to study dimensional changes of the alveolar ridge and root resorption following orthodontic tooth movement into edentulous areas.

**SUBJECTS AND METHOD:** Six adult patients with a thin, resorbed edentulous ridge in the premolar/molar region. Two types of ridge areas adjacent to 10 premolars were defined for evaluation: A – the edentulous area into which the tooth was orthodontically moved and B – the newly established edentulous area from which the tooth was orthodontically moved. Computerised tomographic (CT) examinations were performed at baseline and after treatment. Intra-oral radiographic examination and three-dimensional (3D) measurements on scanned study casts were also performed 1 year after the end of treatment.

**RESULTS:** The CT radiographs revealed that the previously edentulous ridges (A) showed an increase in the bucco-lingual width ranging from 0.5 to 2.6 mm whereas the newly formed edentulous areas (B) showed a reduced width varying between 0.4-2.1 mm. 3D measurements, including soft tissue dimensions, showed an increase in the bucco-lingual width of 0-2.6 mm in A and a reduction of 0.5-1.2 mm in B. Only minor changes were noted one year after the end of treatment. Lateral root resorption was seen at the level of the bone crest on the pressure side in all teeth, varying between 0.7-7.4 mm in height and 0.2-1.4 mm in depth. Four of 10 teeth showed minor apical root resorption.

**CONCLUSION:** Edentulous ridges increased in dimension when teeth were moved into the area. Areas that became edentulous when teeth were moved decreased in dimension. Root resorption was an inevitable side-effect.

## SP 062

Thursday, June 17 - at 9:15

### THE DYNAMIC FOURTH DIMENSION IN IMAGING AND ITS RELATION TO CRANIAL INCLINATIONS

**S Usumez**<sup>1</sup>, I Ramoglu<sup>2</sup>, S Malkoc<sup>3</sup>, T Uysal<sup>2</sup>, Departments of Orthodontics, <sup>1</sup>University of Gaziantep, <sup>2</sup>Erciyes University, Kayseri and <sup>3</sup>Inonu University, Malatya, Turkey

**AIM:** Since the concept of natural head posture (NHP) was introduced, many studies have advocated its use in cephalometric investigations of craniocervical and craniovertical relationships. NHP is a dynamic concept and is successfully recorded as such in a dynamic and continuous manner using inclinometers. The aim of this study was to evaluate how these raw dynamic inclinometer

readings are related to craniovertical (SN-VER) and craniocervical (SN-CVT) measurements and whether the inclinometer readings could be used to determine these measurements without the use of cephalometric radiographs.

**SUBJECTS AND METHODS:** NHP lateral cephalograms of 16 subjects (average age  $19.1 \pm 4.9$  years) were traced and the relationship of the sella-nasion (SN) line with the inclination of the cervical column (CVT) and the true vertical (VER) were measured. The correlations of these measurements with the available dynamic inclinometer readings were calculated.

**RESULTS:** There were very weak correlations between the inclinometer readings and SN-VER ( $r = 0.261$ ) and SN-CVT ( $r = 0.382$ ) measurements.

**CONCLUSION:** Within the limitations of this study the findings suggest that raw dynamic inclinometer readings and craniovertical or craniocervical inclinations are not interchangeable and these raw readings cannot be used to determine these relationships.

## SP 063

Thursday, June 17 - at 9:15

### A NEW DIAGNOSTIC APPROACH TO ORTHODONTIC PATIENTS

**L Sonnesen**<sup>1</sup>, D Nolting<sup>1</sup>, U Engel<sup>2</sup>, KW Kjær<sup>1</sup>, I Kjær<sup>1</sup>, <sup>1</sup>University of Copenhagen and <sup>2</sup>Hvidovre University Hospital, Denmark

**AIMS:** Knowledge of aetiology and correct diagnosis of malocclusions are keys to optimal treatment and treatment outcome. It has been suggested that morphological cervical vertebral column deviations (MCVCD) are important for aetiology, diagnosis and correct treatment of orthodontic patients. The aims of this presentation are to: 1) describe the prevalence and patterns of MCVCD, 2) analyse associations between MCVCD, the craniofacial skeleton and head posture, and 3) elucidate the aetiology behind the associations between MCVCD and the craniofacial skeleton. **MATERIALS AND METHOD:** Profile radiographs of individuals with a neutral occlusion ( $n = 21$ ), severe skeletal malocclusions ( $n = 170$ ) and condylar hypoplasia ( $n = 11$ ). Material from human triploidy fetuses ( $n = 8$ ) and mouse embryos ( $n = 4$ ). The radiographic methods consisted of visual assessment of the cervical vertebral column and cephalometric analysis of the craniofacial skeleton. Histological methods consisted of histochemical and immunohistochemical analyses.

**RESULTS:** Different prevalences and patterns of MCVCD were seen in the various skeletal malocclusions that were associated with a large cranial base angle, retrognathia of the jaws, and an altered head posture. The results from the histological studies on prenatal materials further elucidated these findings.

**CONCLUSION:** These new results are important for diagnostics and aetiology and thus for optimal treatment of orthodontic patients. It is recommended that particular attention is paid to the cervical vertebral column area when analysing profile radiographs and to include any MCVCD in diagnostics and treatment considerations and in the evaluation of the aetiology behind malocclusions in orthodontic patients.

## SP 064

Thursday, June 17 - at 9:15

### SURGERY-FIRST APPROACH IN SKELETAL CLASS III MALOCCLUSION TREATED WITH TWO-JAW SURGERY

H-W Ahn<sup>1</sup>, J-Y Choi<sup>2</sup>, **S-H Baek**<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Oral and Maxillofacial Surgery, Seoul National University, South Korea

**AIM:** To evaluate surgical movement and post-operative orthodontic treatment (POT) of the surgery-first approach for the correction of a skeletal Class III malocclusion (SCIII).

**SUBJECTS AND METHOD:** Eleven SCIII patients who underwent non-extraction treatment and two-jaw surgery [Le Fort I osteotomy impaction of the posterior maxilla (IPM); bilateral sagittal split ramus osteotomy (BSSRO) setback of the mandible (BSM)]. The wafer was removed 4 weeks after surgery. The durations of POT

and total treatment time were  $8.91 \pm 3.14$  months and  $12.18 \pm 3.57$  months, respectively. Lateral cephalograms were taken during the initial examination (T0), immediately after surgery (T1), and after debonding (T2). Sixteen variables were measured. A paired t-test was used for statistical analysis.

**RESULTS:** The maxilla rotated clockwise and the nasolabial angle increased with IPM (FH-palatal plane angle, FH-occlusal plane angle,  $P < 0.01$ ; nasolabial angle,  $P < 0.05$ ) and was well maintained during POT. The mandible was repositioned backward by BSM (SNB, Pog-N perp,  $P < 0.001$ ) and relapsed forward during POT (SNB,  $P < 0.01$ ; Pog-N perp,  $P < 0.05$ ). U1-SN decreased with IPM ( $P < 0.001$ ) and relapsed labially as a result of the Class III mechanics during POT ( $P < 0.01$ ), eventually no significant difference was found between T0-T2. Although IMPA increased with POT, there was no significant difference between T0-T2. The mandible seemed to relapse forward immediately after wafer removal and prior to labioversion of the lower incisors.

**CONCLUSION:** Accurate prediction of POT is crucial in controlling dental alignment, incisor decompensation, arch coordination, and occlusal settling. Long-term wear and selective grinding of the wafer for labioversion of the lower incisors and use of miniplates/miniscrews to control the inclination of the upper incisor are necessary to prevent relapse of the mandible.

## SP 065

Thursday, June 17 - at 9:15

### CEPHALOMETRIC EVALUATION OF WITS ANALYSIS

**I Buck**, S Kopp, Department of Orthodontics, Goethe University, Frankfurt, Germany

**AIM:** The weakness of the tracing of the Wits appraisal on lateral cephalometric headfilms is that the slightest deviation caused by erroneous positioning of the head can result in significant changes in the identification of both points A and B. The alternative to classic tracing represents the direct registration of the points in a magnetic field with the noXrayCeph® device. The present study aimed to assess the diagnostic significance of the Wits analysis by means of a mobile measuring device in order to prove the suitability of this method, especially for appraisal of anteroposterior jaw disharmony. In addition, whether the measurements taken with the mobile measuring device by two different observers correlated with each other was investigated.

**SUBJECTS AND METHOD:** Thirty-six patients were selected and the double series of tracings were recorded by two observers. Dahlberg's coefficient was used to evaluate the reproducibility of the measurements.

**RESULTS:** There was no statistically significant difference in reproducibility of the Wits assessment based on a double series of tracings by either observer. Interobserver repeatability was also very good, the values varied by approximately 90 per cent.

**CONCLUSION:** The major source of error on the lateral cephalometric headfilm such as blurring, fuzziness and shifting as a result of movement during exposure was eliminated with the new mobile head cap of the noXrayCeph® device. The tracing was presented in a real size 1 to 1 scale in accordance with actual anatomy. Therefore, the main advantage of the method is the possibility to directly measure anatomical three-dimensional structures. The method reflects more accurately the extent of anteroposterior jaw dysplasia than the cephalometric lateral image.

## SP 066

Thursday, June 17 - at 9:15

### SURGICAL TREATMENT PLANNING OF PATIENTS WITH CLASS II MALOCCLUSIONS

A Drobyshev<sup>1</sup>, **K Kurakin**<sup>1</sup>, N Drobysheva<sup>2</sup>, E Merzhvinskaya<sup>2</sup>, A Kovalenko<sup>2</sup>, Departments of <sup>1</sup>Maxillofacial Surgery and <sup>2</sup>Orthodontics, Moscow State University of Medicine and Dentistry, Russia

**AIM:** Precise planning of treatment leads to the most predictive results. The purpose of this research was to define value of a di-

vergence of the planned and actual results of surgery with special attention given to soft tissue parameters and facial aesthetics.

**SUBJECTS AND METHOD:** Forty patients with a Class II malocclusion and transverse skeletal deformities. Evaluation of photographs, postero-anterior and lateral cephalographs, anthropometric measurements and computer cephalometric planning was carried out. For all stages a uniform wax record was used that provided stable position of the temporomandibular joints. Soft tissue measurements were made undertaken the method of Arnett. Dolphin Imaging was used with metal markers positioned on the cheek bone, orbital rim, subpupil, nasal base, neck-throat point and a calculation of parameters concerning the true vertical line.

**RESULTS:** According to the generated inspection and planning algorithm, a discrepancy of the changes in bone soft tissue parameters was found. The aesthetics of the face became more harmonious while bone parameters in some cases worsened. In all patients bimaxillary surgery and genioplasty were performed. After surgery all patients underwent orthodontic treatment to achieve maximal intercuspation.

**CONCLUSION:** In most cases it is not sufficient to have only orthodontic calculations and planning as they involve only the average sizes of the bone parameters without division of patients by race. Only careful inspection and planning of treatment allows achievement of an aesthetically harmonious facial profile.

## SP 067

Thursday, June 17 - at 9:15

### THE BIDIMENSIONAL TECHNIQUE EN MASSE RETRACTION USING SELF-LIGATING APPLIANCES

**L Iacopini**, M Greco, A Giancotti, Department of Orthodontics, Fatebenefratelli San Giovanni Calibita Hospital, University of Rome 'Tor Vergata', Italy

**AIM:** The bidimensional (BiDi) technique emphasizes the three-dimensional (3D) control of maxillary incisors and space closure in extraction treatment by sliding mechanics, but needs double steps of retraction phases (first the canines then the incisors). The aim of this presentation is to illustrate 3D control during en masse retraction (one single step) with sliding mechanics using the self-ligating BiDi technique in extraction treatment.

**SUBJECTS AND METHOD:** Sixteen adolescent patients (mean age  $13.9 \pm 1.42$  years) with a Class II division 1 malocclusion, normal skeletal pattern, increased overjet (7 mm) and no crowding in the lower arch, treated with two upper first premolar extractions. Orthodontic treatment was carried out using fixed orthodontic self-ligating brackets (In-Ovation-R) with a BiDi prescription. The brackets consisted of two different slot size (0.018 X 0.025 inch on the anterior teeth and 0.022 X 0.028 inch on the lateral-posterior teeth). During *en masse* retraction, a 0.018 X 0.022 inch stainless steel wire with crimpable hooks distal to the canines, molar toe-in and an accentuated curve of Spee, was inserted in order to manage torque control of the incisors. In the lateral segments the same arch was undersized allowing easy sliding mechanics during space closure with a NiTi coil spring (300 g) extended from the first molar hooks to the archwire hooks. A transpalatal bar was used for posterior anchorage. Panoramic and lateral radiographs were obtained for all patients at the beginning and end of treatment.

**RESULTS:** A Class I canine and Class II molar relationship was achieved for all patients at the end of treatment. All showed bodily movement of the upper incisors with good torque control, confirmed by cephalometric superimposition following the method of Björk, which showed a mean variation of 1-ANS/PNS of 9 degrees.

**CONCLUSION:** The modified BiDi technique with *en masse* retraction on sliding mechanics using a 0.018 X 0.022 inch stainless steel wire and self-ligating brackets could be considered an effective and less time-consuming option in extraction treatment.

**CEPHALOMETRIC DECISION: WHETHER OR NOT TO EXTRACT IN ORTHODONTIC TREATMENT PLANNING**

**P Elmi**, P Nijkamp, L Habets, Section of Orthodontics, Academic Centre for Dentistry Amsterdam (ACTA), Netherlands

**AIM:** The position of the lower incisor, cephalometrically described, is considered important in the decision as to whether or not to extract in orthodontic treatment planning. Several different measurements are known. However, Ricketts' A-Pogonion line (APg to 1) and Downs' mandibular plane to lower incisor (MPL to 1) are the most frequently mentioned, but on an individual basis. The aim of this study was to assess the interchangeability of both measurements and characterization of subjects based on distinct cephalometric differences.

**MATERIALS AND METHOD:** On pre-treatment lateral cephalographs of 115 Caucasians (64 females, 51 males, mean age of  $12.2 \pm 1.1$  years), 13 landmarks were traced twice. Two dental and nine skeletal cephalometric measurements were calculated and the mean values were used for further analysis. To compare APg to 1, mean values from the subjects were converted into Z-scores in relation to the normative means and standard deviations of the two parameters. The subjects were divided into groups based on clear differences in Z-score, group 1 (difference  $< -2$ ), group 2 ( $-2 \leq \text{difference} \leq 2$ ), group 3 (difference  $> 2$ ).

**RESULTS:** The difference between the mean Z-score of APg to 1 and MPL to 1 was statistically significant ( $P < 0.000$ , paired t-test). Statistically significant differences between groups 1, 2 and 3 were found for APg-1, MPL-1, sagittal position of the chin (Pogonion to NB), the anteroposterior relationship of the maxilla to the mandible (ANB), and the vertical inclination of the mandible (Spina Plane and Sella-Nasion to MPL respectively) ( $P < 0.007$ , one way ANOVA).

**CONCLUSION:** Cephalometric measurements, APg to 1 and MPL to 1, should not be used in isolation when irreversible decisions such as extractions are considered in orthodontic treatment planning.

**INFLUENCE OF FUNCTIONAL APPLIANCES ON POSTURAL BALANCE IN CHILDREN WITH MALOCCLUSION**

**I Rubleva**, L S Persin, A Slabkovskaya, V Yurov, Moscow State University of Medicine and Dentistry, Russia

**AIM:** To define the influence of removable functional appliances in children with malocclusions.

**SUBJECTS AND METHOD:** Twenty-six children, aged 8-12 years divided into three groups: group 1 (control) 10 children with normal occlusion; group 2, eight children with a distal occlusion (Angle Class II), orthodontically treated with the pre-orthodontic trainer, and group 3, eight children with distal occlusion, orthodontically treated with a removable functional appliance (Persin). Tests for postural balance (stabilometry) were performed during functional rest and with the teeth tightly closed in an upright stance, with/without the removable functional appliances in situ.

**RESULTS:** In group 1, no significant stabilometric changes were detected. In groups 2 and 3 there was a significant deterioration in postural balance when compared with the functional rest position; the 'passed distance' increased 1.3 fold, the 'sway area' 1.5 times and the 'surface of ellipses' 3.3 times. Stabilometric examination of groups 2 and 3 showed improvement of postural balance with functional appliances. In both groups 2 and 3 the passed distance decreased 1.3 times. The sway area decreased 1.2 times in group 2, and 1.3 times in group 3 and the surface of ellipses decreased by 1.9 times and 2.7 times, respectively.

**CONCLUSION:** Tightly closed teeth in children with normal occlusion did not significantly influence postural balance, while in children with distal occlusion (Angle Class II) this resulted in a significant deterioration in postural balance. Removable functional appliances (pre-orthodontic trainer and Persin's appliance for distal occlusion) significantly improved postural balance.

**THE EFFECT OF LIP ADHESION ON EARLY MAXILLARY GROWTH IN BILATERAL CLEFT PATIENTS**

**G De Pauw**<sup>1</sup>, S De Somere<sup>1</sup>, N Mommaerts<sup>2</sup>, <sup>1</sup>Department of Orthodontics, University of Ghent and <sup>2</sup>General Hospital St Jan Bruges Belgium

**AIM:** To investigate the effect of the lip adhesion procedure on early maxillary growth for bilateral cleft patients.

**SUBJECTS AND METHOD:** Eighteen bilateral complete cleft patients (AG) treated with the lip adhesion procedure 1 month after birth, followed by lip closure 4 months after birth. Another group of 11 bilateral complete cleft patients (PG) pre-surgically treated with a passive moulding appliance followed by lip closure at 5 months of age. Records of the AG group were made at the ages of 1 and 4 months after birth and of the PG group at the ages of 1 and 5 months after birth. Specific landmarks on casts were three-dimensionally digitised. The positional changes of the maxillary segments in both groups were analysed and statistically compared.

**RESULTS:** For the AG group there was a significant reduction of the distance between the premaxilla and lateral segments, of the width of the palatal cleft, of the protrusion of the premaxilla and of the distance between L-L'. There was a significant increase in the width of the premaxilla. For the PG group there was a significant increase in the width of the premaxilla and a significant decrease in the distance between the premaxilla and lateral segments. Between the AG and PG groups, there was a significant difference in transverse dimensions, with higher values for the PG group. There was a significantly higher reduction in the distance between the premaxilla and the lateral segments, only on the right side in the AG group.

**CONCLUSION:** The lip adhesion procedure at the age of 1 month can be advocated in bilateral complete cleft patients with severe premaxillary protrusion and a wide alveolar cleft.

**FINITE ELEMENT ANALYSIS OF STRESS AND STRAIN DURING USE OF RAPID PALATAL EXPANDERS**

**D C Bratu**<sup>1</sup>, L Culea<sup>2</sup>, E Bratu<sup>1</sup>, M Moise<sup>1</sup>, <sup>1</sup>Department of Paedodontics and Orthodontics, Faculty of Dentistry, University 'Victor Babeş' and <sup>2</sup>Department of Material Resistance of the Politechnic Institute, Timișoara, Romania

**AIM:** To analyze stress-strain of the human skull using the finite element (FE) method, simulating the loads at insertion of rapid palatal expanders with 2 or 4 arms.

**MATERIALS AND METHOD:** A digital skull model, produced with Computer Aided Design, Solid Works 2005 programs and force application with FE analysis. The forces applied to the cylindrical forms were equivalent to the teeth over which the palatal expander acts. The software package used was Cosmos Works 2005. The study was conducted first for half a skull and then for a complete (symmetric) skull. Thus two possibilities were studied; a 4 and a 2 point expander. The forces used with the 4 and 2 point expanders were  $F = 22.5$  N and  $F = 45$  N, respectively.

**RESULTS:** Maximum tension was registered at the base of the medio-palatal suture, maximum of 9.364 MPa a smaller value compared with the 133 MPa, which represents the maximum admitted tension. The maximum movement was realized at the level of the free extremity of the mesially orientated cylindrical form that corresponds to the first premolar. The force registered was 9690 for 0.019 mm displacement of the 4 arm and 0.020 mm for the 2 arm expander.

**CONCLUSION:** Regarding the four distinct FEAs, two conclusions can be drawn. The first recommends the use of a 4 arm rapid palatal expander due to the fact that it produces lower levels of stress and deformation than one with 2 arms, and thus a higher degree of comfort for the patient. The second is strictly tied to the FEA approach, the 3D geometrical model for the semi human skull being more appropriate to reality regarding the boundary conditions.

**SKELETAL MATURATION IN CHILDREN**

**M Durka-Zajac**, M Syrynska, K Wozniak, K Lubinska, Department of Orthodontics, Pomeranian Medical University, Szczecin, Poland

**AIM:** Optimal effectiveness in the use of orthodontic or orthopaedic treatment has been associated with skeletal maturation. The aim of this study was to investigate the correlation between chronological age, cervical vertebral maturation (CVM) and Björk's hand-wrist skeletal maturity indicators.

**MATERIALS AND METHOD:** A hand-wrist bone and cephalometric cervical vertebral radiographs of 30 patients (15 males, 15 females; 10-17 years of age). The hand-wrist radiographs were evaluated according Björk's index and CVM assessed using the CVM method. To define vertebral stages, the evaluation consisted of three cervical vertebrae (concavity of C2, C3 and C4 and the shape of C3 and C4). These measurements were then compared with the hand-wrist bone analysis and the results statistically analyzed using Spearman rank correlation.

**RESULTS:** The most frequent cervical vertebrae stages in females were stages CVM V (33.3%) and CVM III (26.7%). For males the most frequent stages were CVM II (46.7%) and CVM III (40.0%). For hand-wrist skeletal maturation, the most frequent stages were MP3u (33.3%) and MP3cap (26.7%) in females and MP3 (33.3%), S (26.7%) and MP3cap (26.7%) in males. Spearman rank-order correlation coefficient between chronological age and cervical vertebrae maturation was 0.53 ( $P = 0.007$ ) for the genders combined; between chronological age and hand-wrist maturation stages it was also 0.53 ( $P = 0.003$ ). Spearman rank-order correlation between hand-wrist and cervical vertebrae maturation was 0.96 ( $P = 0.00001$ ) for both genders.

**CONCLUSION:** A high correlation was found between hand-wrist skeletal maturation and vertebral maturation in Polish subjects.

**SP 073**

Thursday, June 17 - at 10:00

**EPIDEMIOLOGICAL ASPECTS OF MISSING TEETH IN A ROMANIAN POPULATION**

**I N Zetu**<sup>1</sup>, A Ogodescu<sup>2</sup>, M Pacurar<sup>3</sup>, A Petcu<sup>1</sup>, B Dragomir<sup>1</sup>, <sup>1</sup>UMF 'Gr. T. Popa', Iasi, <sup>2</sup>UMF Timisoara and <sup>3</sup>UMF Tg Mures, Romania

**AIM:** There are different opinions regard opening or closing the spaces in subjects with anodontia. The aim of this research was to study the prevalence of missing teeth and their relationship with the occlusion.

**SUBJECTS AND METHOD:** Two thousand six hundred subjects from 5 to 28 years of age (46.62% males, 53.38% females) attending for orthodontic treatment. Dental records, consisting of extra- and intraoral photographs, diagnostic casts and panoramic radiographs, were analyzed.

**RESULTS:** From the sample, 9.84 per cent presented missing teeth (49.59% males, 50.41% females), 37.82 per cent in the maxilla, 42.62 per cent in the mandible; the remaining cases were bi-maxillary. The most frequently missing tooth was the lower second premolar (71%) on one side of the arch, and then the upper lateral incisor, asymmetrical side (48.86%).

**CONCLUSION:** The prevalence of missing teeth in this population was 9.84 per cent corresponding with the literature data (3.5-10%). The sequence in this sample was lower second premolar, upper lateral incisor, upper second premolar and lower central incisor, most frequently asymmetrical. Females were more affected than males, and the maxilla more than the mandible.

**CONE BEAM COMPUTED TOMOGRAPHY FOR ROUTINE TREATMENT PLANNING IN ORTHODONTICS**

**M Nitka**<sup>1</sup>, A Bumann<sup>1</sup>, V Richter<sup>2</sup>, <sup>1</sup>Orthodontic Office, Berlin and <sup>2</sup>Hannover Medical School, Germany

The aim of this research is to demonstrate the advantages of cone beam computed tomography in daily orthodontic treatment planning to assess individual patient needs.

Based upon the findings from more than 2000 orthodontic cone beam computed tomograms (CBCT) per year, the sequence of evaluation is as follows: buccal panoramic view, lingual panoramic view, images of the skull from five angles both with and without soft tissues, images of isolated soft tissues from five angles, three-dimensional (3D) evaluation of either temporomandibular joint, assessment of buccal bone quantity prior to dental arch expansion, mapping of dental arch shape, 2D and 3D cephalometric analysis, 3D assessment of the entire paranasal sinuses, 3D analysis of upper airways and tonsils, 3D diagnosis of asymmetries, 3D imaging of impacted teeth, 3D planning of orthodontic anchorage pins, 3D planning of orthognathic surgery.

The orthodontic analysis described involves an effective radiation dose of approximately 34 to 60  $\mu$ Sv. The combined effective dose of an analogue panoramic radiograph and lateral cephalogram will total 57  $\mu$ Sv, if calculated according to current International Commission on Radiological Protection guidelines.

CBCTs are a useful alternative to conventional panoramic radiographs and lateral cephalograms for orthodontic treatment planning. Compared with conventional radiographic techniques, reasonable additional diagnostic information can be obtained without increasing radiation exposure. At the same time other dental and medical specialties can take advantage of the existing 3D data.

**SP 075**

Thursday, June 17 - at 10:00

**VALIDATION OF CONVENTIONAL RADIOGRAPHS EXAMINATIONS TO ASSESS THE MAGNITUDE OF EXTERNAL RESORPTION**

**I Marek**, M Kamínek, K Langova, Orthodontic Department, Dental Faculty, University of Palacky, Olomouc, Czech Republic

**AIM:** Assessment of radiographic validity in the evaluation of the extent of external apical root resorption (EARR) and comparison of the accuracy of the measurements.

**SUBJECTS AND METHOD:** Seventeen patients who required fixed appliance treatment together with the extraction of the maxillary first premolars. A unilateral force of 50 g was applied for six months to the premolars and a force of 150 g to the contralateral teeth. The teeth were then extracted and measured. For all patients an intraoral conventional radiographs (RVG) and dental pantomograph (DPT) was taken as well as computed tomographic (CT) scans before and after treatment. CT scans were further used to produce digital models. The lengths of the tooth and root crown were measured before and after treatment on the RTGs. Radiographic measurement accuracy was evaluated.

**RESULTS:** The most precise tooth measurements were obtained with the CT scans. The difference between 'real' tooth dimensions and the values obtained from the CT scans were statistically insignificant. Less accurate, but still reasonable, results were obtained with intraoral RVGs, and either inaccurate ones in 3D digital models. The least accurate results were found for DPT measurements – the difference between the DPT and the real dimensions was statistically significant.

**CONCLUSION:** The most accurate measurements of resorption were found with CT scans and intraoral RVGs. Digital models are beneficial for imaging of bone changes at the maxillary sinus during intrusion, and the panorex for diagnosis of severe EARR.

### A NEW CONCEPT FOR ANB ANGLE BASED ON FLOATING NORMS

**K Faltina**, H Barbosa, L Pedrosa, R Bigliuzzi, M Almeida, Department of Orthodontics, University Paulista, São Paulo, Brazil

**AIM:** To develop a new formula to determine individual ANB angle based on characteristics of different facial types.

**MATERIALS AND METHOD:** Eighty lateral telerradiographs of Brazilians of both genders aged between 11 and 17 years with a normal occlusion (skeletal Class I). The sample was divided into three groups according to the facial type: mesofacial, dolichofacial and brachyfacial. The following cephalometric variables were assessed ANB, SNA, ML-NSL, facial axis (FA), total (TFH) and lower (LFH) face height. Multiple regression analysis and t-tests (5 per cent) were used. It was found during statistical analysis that the angular cephalometric variables: FA, TFH, SNA and ML-NSL were suitable for statistical calculation.

**RESULTS:** The calculated values of ANB using the new formula ( $3.84 \pm 1.47^\circ$ ) were identical to the values of ANB measured directly on cephalograms ( $3.84 \pm 1.86^\circ$ ); the paired t-test between these two values ( $P = 0.98378$ ) also confirmed that this new formula was appropriate.

**CONCLUSION:** The established formula having SNA, ML-NSL, FA and TFH [ $ANB = 13.1865 + (-0.3855 FA) + (-0.3696 TFH) + (0.4262 SNA) + (0.3327 ML-NSL)$ ] as the independent variables proved to be adequate for determining the individual ANB angle and more appropriate and specific to the different facial types.

### EFFECTS OF RAPID MAXILLARY EXPANSION ON CRANIOFACIAL SUTURES AND SPHENO-OCCIPITAL SYNCHONDROSIS – A COMPUTERIZED TOMOGRAPHIC STUDY

**R Leonardi**<sup>1</sup>, A Cutrera<sup>1</sup>, M Migliore<sup>1</sup>, E Barbato<sup>2</sup>, <sup>1</sup>University of Catania and <sup>2</sup>University of Rome La Sapienza. Italy

**AIM:** Rapid maxillary expansion (RME) is frequently used in orthodontics. Although the major treatment effect is noticed clinically in the area of the dentition, transverse enlargement of the apical bone or the skeletal structures may be considered as additional contributions. Moreover, RME appears to involve a large portion of the craniofacial complex, with enhanced transverse growth of the circummaxillary anatomical regions. The aims of this investigation were to test the hypothesis that circummaxillary sutures and sphenoid-occipital synchondrosis show bony displacement in response to RME therapy, and to study the extent of suture opening following RME.

**MATERIALS AND METHOD:** Sixteen computed tomographic (CT) images were taken of eight growing patients (2 males, 6 females), before (T0) and after (T1) treatment with RME. All patients had been diagnosed with transverse maxillary deficiency. The mean chronological age of the patients was  $9.8 \pm 1.8$  years (range, 8 to 11.4 years). A high-resolution multislice multidetector CT was used to quantitatively study the extent of the opening of the sphenoid-occipital synchondrosis and craniofacial sutures following RME. A low-dose CT scan protocol was used (80 kV, 10 mA) and the data file of each patient was transferred to a workstation where the anteroposterior width of the sphenoid-occipital synchondrosis and craniofacial sutures were measured on axial and frontal images.

**RESULTS:** At T0, the antero-posterior mean width of the sphenoid-occipital synchondrosis was  $1.73 \pm 0.46$  mm and at T2  $2.30 \pm 0.47$  mm. The craniofacial sutures also showed a significant widening. These differences were statistically significant according to the Wilcoxon signed rank test ( $P < 0.05$ ).

**CONCLUSION:** RME leads to a small but immediate widening of the sphenoid-occipital synchondrosis and most craniofacial sutures in children.

### CEPHALOMETRIC VALUES FOR A MOROCCAN POPULATION COMPARED WITH CAUCASIAN NORMS\*\*\*

**K Arhouni**, C Gilkens, A Balon-Perin, R Glineur, Orthodontic and Dento-Facial Orthopaedic Department, Erasme Hospital, Free University of Brussels, Belgium

**AIMS:** To compare cephalometric values of the Moroccan population with the norms of Caucasians, and evaluate the need for specific norms for this population.

**MATERIALS AND METHOD:** Pre-treatment lateral cephalometric radiographs of 40 Moroccan adults (31 females, 9 males, mean age  $29.5 \pm 7.8$  years), with no history of orthodontic treatment. Eight cephalometric parameters were measured and compared with Caucasian norms (Tweed, Steiner). The parameters measured were FMA, SNA, SNB, ANB, AoBo, I/F, IMPA and facial index.

**RESULTS:** FMA was greater (mean  $30.3 \pm 7.3$ ) when compared with Caucasians. The mean SNB ( $76.5 \pm 4.6$ ) showed that this Moroccan population had mandibular retrognathism relative to the cranial base. They also had proclined upper incisors ( $113.5 \pm 7.5$ ). The other parameters, Wits appraisal ( $-1.9 \pm 5.3$ ), SNA ( $79.9 \pm 4.4$ ), ANB ( $3.4 \pm 3.3$ ), IMPA ( $90.4 \pm 8.4$ ) and facial index ( $0.66 \pm 0.10$ ) were within the normal Caucasian ranges.

**CONCLUSION:** The findings emphasize the necessity of, not only describe group-specific norms, but of individual specificities. Individual variations found within the same population are more relevant mean values.

### ANALYSIS OF OCCLUSAL PLANE POSITION IN PATIENTS TREATED WITH DISTAL BITE CORRECTORS

**M Markova**, L S Persin, L Polma, E Talalaeva, Department of Orthodontics, Moscow State University of Medicine and Dentistry, Russia

**AIM:** Correction of the functional status of dentofacial system and occlusal plane position are necessary to provide a stable outcome of orthodontic treatment. A correctly formed occlusal plane results in normal muscular function of dentofacial system and normal temporomandibular joint status. A new generation of appliances for correction of distal occlusion have recently been introduced – hybrid non-removable bite correctors such as the Forsus, SUS II, and the Twin Force Bite Corrector. However no information is available concerning their effects on morphological and functional changes in the dentofacial system after treatment. The aim of this research was to study the occlusal plane angle ( $<OP/FH: 8 \pm 4^\circ$ ) in patients with a distal occlusion before and after treatment with passive self-ligating brackets and hybrid non-removable bite correctors (SUS II, Twin Force Bite Corrector).

**MATERIALS AND METHODS:** Thirty pre- and post-treatment lateral cephalograms. The position of the occlusal plane was evaluated via  $<Ocp/FH$  angle.

**RESULTS:** In 40 per cent of patients low values for  $<Ocp/FH$  angle were determined pre-treatment that indicated a tendency for low occlusal plane position, while in 60 per cent high values were found indicating a tendency for a high position of the occlusal plane. According to the values of this angle – more or less than 8 degrees – the tendencies were for a vertical or horizontal growth type, respectively. After treatment, a clockwise rotation and an increase of the occlusal plane was noted in patients with low  $<Ocp/FH$  values. In patients with high values, anticlockwise rotation of the occlusal plane occurred and  $<OP/FH$  values decreased.

**CONCLUSION:** In orthodontic treatment with hybrid non-removable bite correctors, the direction of mandibular movement vector is similar to that of facial growth and it normalizes the position of the occlusal plane.

**DIAGNOSTIC INVESTIGATIONS OF PATIENTS WITH OLIGODONTIA AND ECTODERMAL DYSPLASIA**

**M Zadurska**, B Pietrzak-Bilińska, Orthodontic Department, Medical University of Warsaw, Poland

**AIM:** Oligodontia is defined as congenital absence of six or more permanent teeth while ectodermal dysplasia (ED) is a group of diseases characterized by the abnormal development of two or more structures derived from the embryonic ectodermal layer the skin, hair, nails, sebaceous and sweat glands and teeth. Classification of ED is based on the assessment of the occurrence of the four basic clinical symptoms proposed by Freire-Maia and Pinheiro: 1. trichodysplasia, 2. dental abnormalities, 3. onychodysplasia and 4. hypohydrosis. The aim of this investigation was to evaluate patients with oligodontia and suspected ED allowing differentiation of oligodontia and particular ED types.

**SUBJECTS AND METHOD:** A retrospective study of 25 patients (7 females, 18 males) with oligodontia and symptoms of ED, aged from 5 to 24 years. To assess trichodysplasia, a trichogram and trichoscopy were performed. Dental abnormalities were determined through clinical examination and on panoramic radiographs. To assess dyshydrosis, the pilocarpine test was performed.

**RESULTS:** Trichodysplasia was observed in 19 patients (variable shaft thickness, hair shafts without pigmentation, dominance of pilosebaceous units with a single hair, pili torti, trichoschisis and trichotiodystrophy). All 25 patients had dental abnormalities: oligodontia (from 6 to 26 missing teeth), microdontia, conical shaped crowns, taurodontism, abnormal tooth positions, and unerupted teeth. Dyshydrosis was observed in one patient. Pure oligodontia was present in six patients and ED in 19 patients.

**CONCLUSION:** Trichoscopy, trichogram and pilocarpine tests are non-invasive and inexpensive methods for differentiation of pure oligodontia and ED, and types of hydrotic and hypohydrotic ED. Although they will not replace genetic tests, they could be useful in diagnosis, differentiation and planning of comprehensive treatment.

**UNILATERAL POSTERIOR CROSSBITE CORRECTION IN THE MIXED DENTITION – A 3-YEAR FOLLOW-UP**

**S Petrén**, K Bjerklin, L Bondemark, Department of Orthodontics, Faculty of Odontology, Malmö University, Sweden

**AIM:** To compare and evaluate long-term stability of crossbite correction with a quadhelix or expansion plate in the mixed dentition.

**SUBJECTS AND METHOD:** In this randomised controlled study 35 patients with a unilateral posterior crossbite were treated with either a quadhelix or an expansion plate. The inclusion criteria were mixed dentition, unilateral posterior crossbite, no sucking habits or previous orthodontic treatment. Stability was evaluated after 3 years by study cast measurements. Twenty subjects with normal occlusion were included as controls. Success rate, maxillary and mandibular transverse dimensions, overjet, overbite and arch length were registered.

**RESULTS:** Stability was equal for the two treatment methods. Small, albeit significant, differences between the groups were assessed with reference to transverse dimensions. No significant difference was seen for overjet and overbite. The treated patients never reached the same transverse width as the normal control group.

**CONCLUSION:** There was no difference in long-term stability of posterior crossbite correction with a quadhelix and expansion plate. Three years post-treatment, the maxillary width was still greater in the control group than in the treated group.

**ESTIMATION OF DENTAL ALVEOLUS MORPHOLOGY AS A PROGNOSTIC INDICATOR OF GINGIVAL RECESSION**

**M Dominiak**<sup>1</sup>, T Gedrange<sup>2</sup>, B Kawala<sup>3</sup>, M Ziętek<sup>4</sup>, Departments of <sup>1</sup>Dental Surgery, <sup>2</sup>Orthodontics and <sup>3</sup>Periodontology Medical University of Wrocław, Poland and <sup>4</sup>Department of Orthodontics, Preventive and Pediatric Dentistry, Ernst-Moritz-Arndt University, Greifswald, Germany

**AIM:** Gingival recessions (GR) are the final effect of the action of multi-various aetiological factors, of which identification is essential in prevention and dental treatment. The aim of this study was to describe and estimate a method of assessment of labial bone width in the mandibular anterior area in a sagittal direction (API-CEJ2-B) and mental muscle (BD) width based on lateral radiographs in occlusion.

**SUBJECTS AND METHOD:** Ninety-eight randomly selected subjects with an average age 27.8 years, divided into two groups with (I) and without (II) gingival recession and two subgroups with expansion or retraction in the lower incisor region. Using a computer program, cephalometric analysis according to Björk was carried out, in which additionally some modifications were used. API-CEJ2-B [°], API-B [mm], CEJ2-B [mm], height of bone dehiscence (CEJ2-ID [mm]) and muscle (distance between point B and authors D-point, pointing at width of BD) were marked.

**RESULTS:** Significantly lower values for API-CEJ2-B and higher values for CEJ2-ID were found before treatment in group I. As a result of treatment, differences between the bone parameters in both groups were balanced. The greatest change in bone parameter values were observed in group II, and of BD in group I. Expansion treatment, regardless of grouping, significantly influenced the decrease of API-CEJ2-B and the increase of CEJ2-ID, whereas retraction in group I additionally caused a significant increase of BD width. Prediction values of parameters that considerably influence the occurrence of GR are: API-CEJ2-B <16 degrees, CEJ2-ID >1.4 mm and age >24.4. A constructed multiple regression model of parameters which significantly influence GR explained the potential recession-causing influence in 72 per cent.

**CONCLUSION:** Assessment of API-CEJ2-B is of diagnostic importance as well as for determining therapy.

**AESTHETICS BEFORE AND AFTER ORTHODONTIC TREATMENT**

**A Muntean**<sup>1</sup>, A Valceanu<sup>1</sup>, F Glavan<sup>2</sup>, E Bratu<sup>2</sup>, C Szuhaneck<sup>2</sup>, Departments of <sup>1</sup>Esthetic Dentistry, and <sup>2</sup>Orthodontics, University of Medicine and Pharmacy, Timisoara, Romania

**AIM:** An attractive smile is important for successful aesthetic and orthodontic treatment. Lip position, oral condition, personality traits, and smiling affect the aesthetics of the smile. The aim of this study was to analyse the changes of some aesthetic parameters before and after orthodontic treatment.

**SUBJECTS AND METHOD:** Ninety-two healthy subjects from the west region of Romania divided into four categories: 23 girls aged between 7 and 13 years, 23 boys between 7.5 and 12.5 years, 23 females between 19 and 35 years and 23 males between 21.5 and 32.7 years. All patients underwent a complete aesthetic analysis before and after orthodontic treatment. Aesthetic evaluation, including measurement of inferior face height, was based on photographs taken in both the frontal and profile view, and during social smiling.

**RESULTS:** At the end of orthodontic treatment the values were higher than those pre-treatment, especially for females compared with males. The profile angle showed a large change in girls compared with females. Smile analysis was used to evaluate the incisal smile line, the labial line of the smile, the buccal corridors and the midline. The changes were more obvious for females compared with males and for younger age groups compared with adults.

**CONCLUSION:** It is important to take into account aesthetic parameters while establishing an orthodontic treatment plan.

### A COMPARATIVE STUDY OF LOWER THIRD MOLAR SPACE IN SKELETAL CLASS III AND CLASS II ANOMALIES

**M Popescu**, O Popoviciu, O Eftene, I Beuran, Department of Orthodontics and Dentofacial Orthopedics, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

**AIM:** Customarily it is considered that in Class III anomalies mandibular macrognathia provides sufficient space for the eruption of the lower third molars, while in Class II anomalies the lack of space is due to micrognathia. However, the opposite situation could be encountered, as the space available for the eruption of the lower third molar depends not only on mandibular length, but also on other developmental skeletal and dental characteristics. The aim of this research was to perform a comparative study on the eruptive perspective of the third molar among subjects with skeletal Class III and Class II anomalies.

**SUBJECTS AND METHODS:** Two patient groups were selected, based on mandibular development criteria: 21 patients with mandibular prognathia (Angle Class III) and 27 patients with mandibular micrognathia (Angle Class II). Measurement of available space for the third molars (measured on profile cephalograms and dental pantomograms) was conducted between the Class II/Class III subjects, hypodivergent/hyperdivergent sub-samples in each sample and between groups of patients with the same rotational pattern in the two samples.

**RESULTS:** Space corresponding to the lower third molar did not display a statistically significant difference between the two samples as, despite mandibular macro/micrognathia, each sample contained subjects with/without adequate space available for the lower third molars.

**CONCLUSION:** Several interconnected factors influence third molar eruption and alignment, including mandibular base dimension, alveolar bone perimeter, lower arch dental perimeter and facial growth rotation pattern.

### ORTHODONTIC TREATMENT OF PATIENTS WITH CROWDING USING SELF-LIGATING BRACKETS

**O Emeliyanova**, T Yavorovkaya, Y Gioeva, N Golovinova, Moscow State Medical Dentistry University, Russia

**AIM:** To compare the results of orthodontic treatment of patients with crowding and to determine the influence of the functional condition of the orbicularis oris muscle on the results of treatment of such patients using passive and active self-ligating brackets.

**SUBJECTS AND METHOD:** Sixty patients with crowding divided into three groups: group 1, 20 patients, treated with passive self-ligating brackets, Damon-3; group 2 patients treated with active self-ligating brackets, Time-2, and group 3, patients treated with ligating brackets MBT. Dental casts were measured using electronic digital calliper before and after active treatment. A unique patented device, the Vestibulotometer (patent application number 12007115849) was used to determine the influence of the muscles surrounding the mouth on craniofacial structures.

**RESULTS:** Group 1 patients showed maximum expansion between the upper (10 mm) and lower (6 mm) premolars while a significant increase in arch length was observed in group 3. A low correlation was found that the orbicularis oris muscle tonus had an impact on treatment in group 3. A significant medium correlation was found with interincisor angle (correlation coefficient 0.5). In patients treated with active self-ligating brackets, lip tonus influence was also significant on interincisor angle alteration. In the patients with passive self-ligating brackets, there was an influence of lip tonus on the length of the dental arch and interincisor angle.

**CONCLUSION:** Passive self-ligating systems allow expansion of narrow dental arches without additional appliances. Lip tonus influences the results of treatment. Hypertonus assists in the decrease of incisor protrusion and dental arch expansion, but not in retrusion elimination.

### PREDICTION OF THE MESIODISTAL WIDTH OF PERMANENT TEETH

**T Yavorovskaya**, O Emeliyanova, Y Gioeva, N Golovinova, Moscow State Medical Dentistry University, Russia

**AIM:** During the period of the mixed dentition several methods are used to predict tooth width. They show high values of correlation and determination coefficients. The purpose of present study was to determine a formula for prediction of the width of unerupted canines and premolars based on the sum of the width of the first permanent molar and incisors.

**MATERIALS AND METHOD:** Two hundred dental casts (100 maxillary, 100 mandibular) of 26 male and 64 female Caucasian patients with an average age of 16.6 years. Mesiodistal tooth width was measured with an electronic digital calliper. Statistical and correlation analysis were used.

**RESULTS:** The results showed high correlation (r) values, varying from  $r = 0.677$  to  $r = 0.831$ . These values are among the highest described in the literature for non-radiographic prediction methods. There was no significant difference between the predicted and actual widths of the maxillary and mandibular permanent canines and premolars. The standard errors of estimation were 1.18 for males and 1.2 for females for the maxillary arch, and 1.01 for males and 1.02 for females for the mandibular arch. No statistical difference was found in tooth width between the right and left sides.

**CONCLUSION:** This method is accurate, practical, and easy to use and requires no specific equipment.

### HYOID BONE POSITION, MANDIBULAR GROWTH DIRECTION AND HEAD POSTURE IN MOUTH BREATHERS

**G Curcievea-Cuckova**<sup>1</sup>, I Kirovski<sup>2</sup>, M Pop Stefanova-Trposka<sup>1</sup>, S Popovski<sup>1</sup>, R Dimovska<sup>3</sup>, <sup>1</sup>Department of Orthodontics, P.H.O. University Dental Clinical Center, <sup>2</sup>Military Hospital and <sup>3</sup>Private dental practice-Kruna MS, Skopje, Former Yugoslav Republic of Macedonia

**AIM:** To assess the effects of impaired nasal breathing on head posture, position of the hyoid bone and the tongue, and mandibular growth direction. A further aim was to estimate the relationship between head posture and hyoid bone position.

**MATERIALS AND METHOD:** Study casts and lateral skull radiographs of 100 children from 8 to 14 years of age. After otorhinolaryngological evaluation, the patients were divided into two groups. Group 1 comprised 50 mouth breathers with nasopharyngeal obstruction and group 2, 50 nose breathers with no history of impaired nasal breathing, who served as the control. To estimate the position of the hyoid bone and the tongue, six diameters (hy-gn, hy-cv4ip, hy-MP, hy-NS, PP-hy and PP-dl), and six craniocervical angles determining the extension of the head (NS/OPT, FH/OPT, PP/OPT, NS/CVT, FH/CVT and PP/CVT) were measured. Standard lateral cephalometric radiographs were obtained to evaluate mandibular growth direction using the following measurements: inclination of the mandibular plane, gonial angle and basal plane angle.

**RESULTS:** Hyoid position in children in group 1 was found to be more inferior to normal in relation to the mandibular plane and SN reference line, and more distally positioned. The average craniocervical angles were found to be extremely large, exceeding the average values in group 2. The significantly increased gonial and basal plane angle, and the slope of the occlusal plane and higher inclination of the mandibular plane in group 1 indicated vertical growth in a clockwise rotation. A steeper PP/Mp angle, and palatal-tongue distance showed a positive correlation with an increased craniocervical angulation.

**CONCLUSION:** Determination of hyoid bone and tongue position in children with impaired nasorespiratory function can be used in therapeutic and prognostic evaluation.

### CONE BEAM COMPUTED TOMOGRAPHY OF THE TEMPOROMANDIBULAR JOINT IN THE SURGICAL TREATMENT OF CLASS III ANOMALIES

**M Popescu<sup>1</sup>**, O Dinca<sup>2</sup>, O Popoviciu<sup>1</sup>, A Bucur<sup>2</sup>, D Stanciu<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and Dentofacial Orthopedics and <sup>2</sup>Oro-Maxillary-Facial Surgery, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

**AIM:** A bilateral sagittal split osteotomy (BSSO) is one of the most frequently used surgical procedures for correcting mandibular prognathia. It has been widely accepted that maintaining the proximal mandibular segment in its pre-operative condylar position contributes substantially to the stability of the surgical result and reduces the adverse effects on the temporomandibular joint (TMJ). The aim of this research was to evaluate TMJ changes following orthognathic surgery in subjects with severe Class III anomalies.

**SUBJECTS AND METHOD:** Twenty young adults with severe Class III anomalies, who had received combined orthodontic and orthognathic surgical treatment. Cone beam computed tomograms (CBCT) of the TMJs were obtained pre- and post-surgery. Patient positioning parameters were defined in order to obtain the same axial sections on the CBCTs at the two time points, thus enabling comparative analysis. Based on the measuring method on conventional radiographs described by Pullinger and Hollinger, the distances from the centre of the condyle to the glenoid cavity border were measured at 0, 30, 60, 120, 150 and 180 degrees.

**RESULTS:** Comparative statistical analysis of the data revealed changes in the position of the condyle in the glenoid cavity 6 weeks post-surgery compared with pre-surgery, but without statistical significance (except for the left side at the 30 degree evaluation).

**CONCLUSION:** It is important for the results of surgery to be verified through a standardized CBCT method, in order to appreciate the quality of surgical intervention. The adaptive capacity of the TMJs may compensate for minor changes.

### ASSESSMENT OF CROWDING WITH A SELF-LIGATING SYSTEM IN NON-EXTRACTION THERAPY

**B Vande Vannet<sup>1</sup>**, W Jacquet<sup>2</sup>, G Ibel<sup>1</sup>, H-P Bantleon<sup>3</sup>, <sup>1</sup>Vrije Universiteit Brussel and <sup>2</sup>Visie Lab, Universiteit Antwerpen, Belgium and <sup>3</sup>Abteilung für Kieferorthopädie, Bernhard Gottlieb-Universitätszahnklinik, Medizinische Universität Wien, Austria

**AIM:** To evaluate three-dimensional tooth changes in patients with severe upper and lower crowding treated with the Damon® system (D3, Ormco, Orange, California, USA) without extractions or approximal enamel reduction.

**SUBJECTS AND METHOD:** Twenty-two patients (14 females, 8 males) were examined by two experienced orthodontists. The digital models obtained were analyzed with the OrthoProof® software (Nieuwegein, Netherlands) determining 72 parameters in the upper and 67 parameters in the lower arch. In addition lateral cephalograms of the patients were investigated with the Onyx Ceph® software (Chemnitz, Germany) analyzing 28 cephalometric parameters. With all gained data, superimpositions were made for graphical representation of the changes regarding the dental arch and the cephalometric tracing. Statistical analysis was performed using the Statistical Package for Social Sciences, version 15.0 (SPSS Inc., Chicago, Illinois, USA).

**RESULTS:** Strong interrater correlation was found between the measurements of both examiners ( $P < 0.001$  ranging between 0.83-0.96). There was no correlation regarding initial upper and lower crowding between the initial and final measurements for proclination. For protrusion, a correlation was present between arch expansion and protrusion of the upper ( $P < 0.05$ ) and lower ( $P < 0.01$ ) incisors. However, the changes concerning the position

of the upper anterior teeth showed more variability, indicating that the space needed to correct the crowding was created by other mechanisms, in particular by expansion. The soft tissues, i.e. the lip profile, mainly followed the upper anterior teeth.

**CONCLUSION:** Non-extraction therapy with the Damon® system was effective within biological limits and with positive effects on the soft tissues.

### VACUUM RETAINER TECHNIQUE IN THE TREATMENT OF ANTERIOR CROSSBITE

**I Apruzzese**, M Greco, A Giancotti, University of Rome Tor Vergata Fatebenefratelli Hospital, Italy

**AIM:** Invisalign treatment is gaining wider applications in the treatment of different malocclusions; not only for simple arches alignment. The purpose of this study was to evaluate the efficacy of the Invisalign system in the treatment of anterior crossbite.

**SUBJECTS AND METHOD:** Twelve adult patients (mean age 21.9 ± 1.42 years) with skeletal pseudo-Class III malocclusions, a Class III molar and canine relationship, normal skeletal pattern and a negative overjet (-2 mm). Orthodontic treatment was carried out avoiding extractions and by means of interproximal reduction in the mandibular arch and expansion in the maxillary arch in order to achieve anterior crossbite correction. All patients were instructed to use Class III intermaxillary elastics from the attachment on the first maxillary molars to those on the lower canines. To simplify connection of the elastics, the attachment was left empty and partially modified. Cephalometric superimposition was carried out on lateral radiographs obtained at the beginning and end of treatment.

**RESULTS:** A normal overjet relationship was achieved in all patients with a Class I molar and canine relationship. Cephalometric superimposition showed a favourable change in maxillary (1/ANS-PNS mean variation, 9°) and mandibular incisor inclination (1/GoGn mean variation: -3°).

**CONCLUSION:** The Invisalign technique proved to be an efficient option for the treatment of anterior crossbite in skeletal pseudo-Class III malocclusion subjects.

### VALIDITY OF UPPER AIRWAY ASSESSMENT IN CHILDREN

**K Pirilä-Parkkinen<sup>1</sup>**, H Löppönen<sup>2</sup>, P Nieminen<sup>3</sup>, U Tolonen<sup>4</sup>, P Pirttiniemi<sup>5</sup>, <sup>1</sup>Oral and Maxillofacial Department and <sup>4</sup>Department of Clinical Neurophysiology, Oulu University Hospital, <sup>2</sup>Department of Otorhinolaryngology, Institute of Clinical Medicine, University of Kuopio, <sup>3</sup>Department of Otorhinolaryngology, Vaasa Central Hospital <sup>5</sup>Institute of Dentistry, University of Oulu, Finland

**AIM:** Lateral cephalometric radiography is a conventional orthodontic method used for determining craniofacial morphology and possible upper airway obstruction. Assessment of pharyngeal obstruction is particularly important in children with breathing difficulties. The aim of this study was to evaluate the possibility of using two-dimensional (2D) lateral cephalograms to identify pharyngeal obstruction in relation to three-dimensional magnetic resonance imaging (MRI) and clinical observation of tonsillar size.

**SUBJECTS AND METHOD:** Thirty-six children (19 males, 17 females, mean age 7.3 years, range 4.8-9.8 years) with sleep-disordered breathing diagnosed by nocturnal polygraphy. Pharyngeal airway was imaged with a low field open configuration magnetic resonance scanner. An otorhinolaryngological clinical examination was carried out and lateral skull radiographs were taken and measured.

**RESULTS:** Nasopharyngeal cephalometric variables had a significant positive correlation with the minimal cross-sectional nasopharyngeal airway area (MRI). From oropharyngeal cephalometric variables, retropalatal measurements correlated with MRI

findings. Both the MRI and cephalometric technique showed the narrowest pharyngeal measurement to be located in the retro-palatal region. Clinical assessment of tonsillar size correlated inversely with MRI findings such as minimal retro-palatal cross-sectional airway area ( $P = 0.000$ ), minimal retro-glossal cross-sectional airway area ( $P = 0.015$ ) and intertonsillar airway width ( $P = 0.000$ ). Cephalometric soft palate and tonsillar area correlated with clinical tonsillar size ( $P = 0.001$ ). The results can be explained by anatomical and postural factors.

**CONCLUSION:** Lateral cephalometric radiography is a useful screening tool when evaluating nasopharyngeal and retro-palatal airway size. However, much of the retro-glossal oropharyngeal information is missing in the antero-posterior 2D view. Clinical inspection of oropharynx and tonsillar size is a relatively reliable method when oropharyngeal airway size is examined.

## SP 092

Thursday, June 17 – at 11:30

### AESTHETIC CHANGES OF THE PROFILE IN EXTRACTION AND NON-EXTRACTION CASES

**A Muntean**<sup>1</sup>, F Glavan<sup>2</sup>, E Bratu<sup>2</sup>, R Jianu<sup>2</sup>, C Szuhaneck<sup>2</sup>, Departments of <sup>1</sup>Esthetic Dentistry, and <sup>2</sup>Orthodontics, University of Medicine and Pharmacy, Timisoara, Romania

**AIM:** The decision concerning extraction or non-extraction in orthodontics is of interest in aesthetic dentistry, especially as it can influence the profile, smile and face of the patient. The aim of this study was to examine the effects of orthodontic treatment on the soft tissue profile treated either extraction or non-extraction with fixed appliances.

**SUBJECTS AND METHOD:** Forty healthy subjects with a Class II division 1 or division 2 malocclusion. Profile photographs were taken before treatment, after fixed appliance removal and one year post-treatment. The subjects were 20 females (10 extraction and 10 non-extraction) and 20 males (10 extraction and 10 non-extraction) aged between 19 and 31 years.

**RESULTS:** The profile angle increased in most of the non-extraction subjects by approximately 3.5 degrees and decreased in the extraction cases by 2.7 degrees. The differences between males and females were less significant. The distance between the aesthetic line and the upper lip decreased. The nasolabial angle also decreased by 2.5 degrees in the extraction cases.

**CONCLUSION:** The profiles of the treated subjects became straighter immediately after treatment and was unchanged one year later. The differences between extraction and non-extraction aesthetic profiles were minimal.

## SP 093

Thursday, June 17 – at 11:30

### ROOT DAMAGE AFTER SCREW INSERTION FOR ORTHODONTIC MINIPLATE PLACEMENT IN DOGS

**M Cornelis**<sup>1</sup>, C Behets<sup>2</sup>, <sup>1</sup>Departement of Orthodontics, Université de Genève, Switzerland and <sup>2</sup>Experimental Morphology Unit, Université catholique de Louvain, Brussels, Belgium

**AIM:** The risk of damaging roots during placement of temporary skeletal anchorage devices is not negligible. The aim of this investigation was to describe the healing reactions following root damage using a miniplate anchorage system.

**MATERIALS AND METHOD:** In four beagle dogs, 32 titanium miniplates (2 self-tapping screws/miniplate) were placed between the roots, after drilling of pilot holes (1.6 mm-Ø). During the drilling process of six screws, the root and/or periodontal ligament (PDL) of the upper canine was damaged. Two weeks later, half of the miniplates were loaded with a coil spring. The dogs were killed 7 or 29 weeks after placement. Fluorescent markers were injected at placement and at sacrifice. Jaw samples were embedded and cut into sections. The sections were microradiographed and then observed under ultraviolet light.

**RESULTS:** Four roots were directly damaged by the screws. One screw did not contact the root itself, but the hole was drilled deeper than required, damaging the root. One screw went into the PDL without damaging the root. Three screws injured the roots at the apex level, and three screws hit the mesial side of the canine root or its PDL. Among the six involved screws, two were mobile and four were stable at sacrifice. This distribution was not related to the location of the root damage or the loading status of the miniplates. Observation of the fluorescent markers displayed almost no root repair after 7 weeks. In contrast, a thick layer of mineralized material was observed on the lesions 29 weeks after root damage. The screw placed into the PDL but not in direct contact with the root, did not result in any root resorption. The presence or absence of tissue repair was not related to root damage location, screw stability or loading status.

**CONCLUSION:** Limited root damage showed signs of healing by apposition of new cementum in the defect after 29 weeks.

## SP 094

Thursday, June 17 – at 11:30

### THE ROLE OF MODERN THREE-DIMENSIONAL IMAGING IN ORTHODONTIC DIAGNOSIS AND TREATMENT PLANNING

**G Rossini**<sup>1</sup>, R Leonardi<sup>2</sup>, E Barbato<sup>1</sup>, <sup>1</sup>University of Rome and <sup>2</sup>University of Catania, Italy

**AIM:** The information obtained from conventional two-dimensional radiographs is to enable an accurate and complete orthodontic-orthopaedic treatment plan to be formulated. Three-dimensional (3D) imaging provides more detailed information, particularly of the maxillofacial area, allowing the design of a more precise and predictable treatment plan. The aim of this research is to describe the possible fields of use of 3D imaging in orthodontics and to outline the correct role, advantages and possible limitations of the new technologies of imaging and analysis.

**MATERIALS AND METHOD:** Medline, Science Direct, the Cochrane collaboration and the Blackwell DSynergy were all used to access published journals on the key words of orthodontic diagnosis, treatment plan, cone beam computed tomography, cephalometric analysis, 3D imaging. Using the available search engines, numerous studies were found relevant to the topic. The different applications of 3D imaging technologies in orthodontics were analysed, together with related clinical cases.

**RESULTS AND CONCLUSION:** The possibilities for application of the new imaging technologies are numerous and they can significantly improve the quality of treatment. The information on teeth, soft and hard tissues, upper respiratory tract and on the relationship between the different structures of the head and neck are more detailed and give many opportunities for in-depth diagnosis. The large amount of information available should be standardized and selected to be interchangeable between different operators.

## SP 095

Thursday, June 17 – at 11:30

### EFFECTS OF SURGICALLY ASSISTED RAPID MAXILLARY EXPANSION ON CRANIOFACIAL MORPHOLOGY

**A Y Gungor**<sup>1</sup>, H Turkkahraman<sup>2</sup>, T Baykul<sup>3</sup>, H Alkis<sup>1</sup>, <sup>1</sup>Department of Orthodontics, <sup>1</sup>Mustafa Kemal University, Hatay and Departments of <sup>2</sup>Orthodontics and <sup>3</sup>Oral Surgery, Suleyman Demirel University, Isparta, Turkey

**AIM:** To evaluate the effects of surgically assisted rapid maxillary expansion (SARME) on craniofacial morphology in non-growing patients.

**SUBJECTS AND METHOD:** Twenty-eight patients divided into two groups. SARME, with a banded Hyrax-type expander, was carried out in group I (4 boys, 10 girls, mean age 19.6 ± 2.73 years). Rapid maxillary expansion (RME) with an occlusal-coverage Hyrax-type expander was applied to group II (4 boys 10 girls, mean age 14.2 ± 0.74 years). All expanders were activated twice a day (0.5 mm/day). Lateral and posteroanterior cephalograms and

dental casts were obtained before and after expansion, and measurements were performed. Intra- and intergroup differences were statistically evaluated with Wilcoxon and Mann-Whitney U tests.

**RESULTS:** In both groups, significant increases were found in intercanine, interpremolar, intermolar, maxillary and nasal widths, and anterior face height ( $P < 0.05$ ). Statistically significant intergroup differences were found for all maxillary dental cast measurements ( $P < 0.05$ ) and in nasal cavity width ( $P < 0.05$ ). No significant difference was found between cephalometric measurements of the groups ( $P > 0.05$ ).

**CONCLUSION:** With both SARME and RME, successful expansion of maxillary dentoalveolar structures, nasal cavity and palatal widening were achieved. Compared with RME, SARME was found to be more effective in nasal cavity, palatal and dental arch widening.

### SP 096

Thursday, June 17 – at 11:30

#### ORTHODONTIC-PERIODONTIC INTERRELATIONSHIP: A SYSTEMATIC REVIEW

**P Christou**<sup>1</sup>, N Gkantidis<sup>2</sup>, N Topouzelis<sup>2</sup>, Departments of Orthodontics, <sup>1</sup>Athens University and <sup>2</sup>Thessaloniki University, Greece

Orthodontic treatment aims at providing an acceptable functional and aesthetic occlusion with appropriate tooth movements. These movements are strongly related to interactions of teeth with their supportive periodontal tissues. In recent years, because of the increased number of adult patients seeking orthodontic treatment, orthodontists frequently encounter patients with periodontal problems.

Aesthetic considerations, such as uneven gingival margins or functional problems resulting from inflammatory periodontal diseases should be considered in orthodontic treatment planning. Furthermore, in cases with severe periodontitis, orthodontics may improve the possibilities of saving and restoring a deteriorated dentition. In modern clinical practice, the contribution of the orthodontist, the periodontist and the general dentist is essential for optimized treatment outcomes.

The aim of the present review is to describe the existing evidence in the literature concerning certain types of orthodontic tooth movement in cases of compromised periodontium. The tooth movements considered and reported with relevant cases are: orthodontic extrusion and intrusion, molar uprighting, orthodontic movement in edentulous areas, and labial tilting.

### SP 097

Thursday, June 17 – at 11:30

#### EARLY TREATMENT OF UNILATERAL POSTERIOR CROSSBITE – AN ELECTROGNATHOGRAPHIC EVALUATION

**E Sever**, E Vehovec, I Kopač, L Marion, M Ovsenik, Departments of Orthodontics, University Medical Centre, Ljubljana, Slovenia

**AIM:** Occlusion is an important factor that affects chewing. Unilateral posterior crossbite (UPXB) patients exhibit asymmetry in motion of the mandible during mastication. The aim of this study was to investigate the characteristics of common chewing cycles in the primary dentition in children with a UPXB, when chewing gum on the UPXB side and on the non-UPXB side.

**SUBJECTS AND METHOD:** Thirty children with UPXB ( $5.42 \pm 1.2$ ) and 30 children ( $5.38 \pm 1.4$ ) with a normal primary occlusion. Chewing movements were recorded with an electrognathographic (Sirognathograph) and computer analysing system (CO-SIG II). The chewing patterns were averaged and analysed in the frontal and in the sagittal projection planes and compared using a Student's *t*-test ( $P < 0.05$ ) with respect to the relationship between chewing pattern and occlusion before (T0) and after (T1) treatment, and after retention (T2).

**RESULTS:** Children with a UPXB opened their mouths wider and had chewing cycles orientated more posteriorly at T0. They also had a shorter rest position in maximal intercuspation and more frequently reverse directed chewing cycles on the UPXB side.

At T1, children with a UPXB showed more anteriorly orientated chewing cycles in the sagittal projection plane. At T2 no statistically significant differences between the sides in UPXB children and the control group were found.

**CONCLUSION:** Early treatment is recommended to normalize the masticatory chewing cycle pattern, to ensure normal chewing function and development of the orofacial system.

### SP 098

Thursday, June 17 – at 11:30

#### ASSESSMENT OF CHEWING CYCLE MORPHOLOGY – A LONGITUDINAL STUDY IN CHILDREN FROM 5 TO 7 YEARS OF AGE

**E Vehovec**, E Sever, I Kopač, J Primožič, M Ovsenik, Faculty of Medicine, University of Ljubljana, Slovenia

**AIM:** To evaluate the relationship between chewing cycle morphology and the developmental stage of the dentition.

**SUBJECTS AND METHOD:** The chewing pattern was assessed in 31 children ( $x = 5.42 \pm 0.50$  years) with the use of the Sirognathograph electrognathographic device in the primary dentition period (T1), in the early mixed dentition ( $x = 6.02 \pm 0.50$  years) (T2) and after eruption of the first permanent molars ( $x = 6.92 \pm 0.50$  years) (T3). Chewing pattern morphology was analysed in terms of shape and duration of the chewing cycles. Statistical analysis of the data was performed.

**RESULTS:** Chewing pattern morphology was found to be statistically significantly different between T1 and T2 for seven, between T2 and T3 for two and between T1 and T3 for 10 chewing parameters. In the frontal plane, the lateral deviation of chewing cycles from the midline was enlarged from 2.51 mm (T1) to 3.28 mm (T2) to 4.52 mm (T3). The axial inclination angle and the mouth closing angle were reduced. The cycles surface was enlarged from 15.86 mm<sup>2</sup> (T1) to 33.67 mm<sup>2</sup> (T3). In the sagittal plane, an enlarged inclination angle of chewing cycles (from 88.88° to 121.70°) was measured at T1 and T2 and a decline at T3 (114.66°). The mouth closing angle (from 84.79° to 114.71°) was enlarged at T2 and reduced at T3 (106.33°). The cycles surface was enlarged (from 2.77 mm<sup>2</sup> to 8.49 mm<sup>2</sup> to 9.40 mm<sup>2</sup>). The average opening time was extended (from 0.21 to 0.24 to 0.28 seconds), resulting in a prolonged duration of the whole chewing cycle (from 0.57 to 0.65 to 0.69 seconds).

**CONCLUSION:** Eruption of the permanent teeth has a significant influence on the shape and duration of the chewing cycle. An objective assessment of chewing cycle morphology may be an important part of functional diagnostics in orthodontics.

### SP 099

Thursday, June 17 – at 11:30

#### INCOMPETENT LIP SEAL AND TONGUE POSTURE – IS THERE A CORRELATION?

**V Božnik**<sup>1</sup>, J Volk<sup>1</sup>, M Mušič<sup>2</sup>, M Ovsenik<sup>1</sup>, Department of <sup>1</sup>Orthodontics and <sup>2</sup>Institute of Oncology University of Ljubljana, Slovenia

**AIM:** An incorrect tongue posture on the floor of the mouth is an important aetiological factor for malocclusion development. It is difficult to clinically diagnose tongue posture in incompetent lip seal patients in the early dental developmental period. The aim of this study was to assess tongue posture in children with an incompetent lip seal (ILS) and in those with a competent lip seal (CLS) in the normal mixed dentition using three-dimensional (3D) ultrasonography.

**SUBJECTS AND METHOD:** Fourteen children with an ILS (mean age 7.91 years; SD =  $\pm 1.53$ ) and 15 children with a CLS (control group; mean age 7.48 years; SD =  $\pm 0.38$ ). An ultrasound system, Voluson 730 Expert, and a 3D convex transducer, RAB 2-5 MHz, were used in the study. The reference 3D ultrasound reconstructions were used for assessment of the tongue posture twice in each child. Fisher's exact test was used for statistical analysis.

**RESULTS:** Tongue posture was found to be statistically significantly different between ILS and CLS. Sixty-four per cent of children with ILS and 33 per cent of children with CLS demonstrated posture on the mouth floor. The difference was statistically significant ( $P < 0.005$ ).

**CONCLUSION:** In the majority of ILS patients the tongue is on the mouth floor. Therefore, assessment of tongue posture in ILS patients is an important aetiological factor in functional diagnosis of orthodontic patients.

## SP 100

Thursday, June 17 – at 11:30

### THREE-DIMENSIONAL CHARACTERISTICS OF FACIAL AND JAW MORPHOLOGY IN THE EARLY MIXED DENTITION

**D Malinar**<sup>1</sup>, B Krneta<sup>1</sup>, A Zhurov<sup>2</sup>, S Richmond<sup>2</sup>, M Ovsenik<sup>1</sup>,

<sup>1</sup>Department of Orthodontics, Medical Faculty, University of Ljubljana, Slovenia and <sup>2</sup>Dental Health and Biological Sciences, Dental School, Cardiff University, Wales

**AIM:** Facial diagnosis in small children performed from photography and has several drawbacks. A three-dimensional (3D) imaging system has been shown to be accurate for scanning young children and is a non-invasive and completely safe method to obtain 3D facial images. The purpose of this research was to evaluate facial and jaw morphological characteristics in children in the early mixed dentition period.

**SUBJECTS AND METHOD:** Thirty children [17 girls (mean age  $6.94 \pm 0.60$  years) and 20 boys (mean age  $6.98 \pm 0.61$  years)]. Facial images were taken using the 3D imaging system. The 3D data was imported to a reverse modelling software package, Rapidform™ 2006. Each scan of the face was processed in order to remove unwanted data, registered and merged to produce a complete facial image. A Student's t-test was used to assess differences between the facial templates of the girls and boys.

**RESULTS:** The average facial and study cast template of a Slovenian girl and boy in the early mixed dentition period was established. The difference between the male and female facial template was determined in the cheek area, and in the lower jaw and eyebrow regions. No difference between genders was found in the length of the middle and lower facial third. Study cast analysis of intermolar and intercanine width revealed statistically significant differences between the female (mandibular intercanine  $24.08 \pm 1.49$ , intermolar  $43.56 \pm 1.58$ ; maxillary intercanine  $30.73 \pm 1.75$ , intermolar  $47.79 \pm 1.78$ ) and male (mandibular intercanine  $26 \pm 2.31$ , intermolar  $43.99 \pm 3.00$ ; maxillary intercanine  $32.01 \pm 2.55$ , intermolar  $49.29 \pm 2.20$ ) templates.

**CONCLUSION:** The 3D laser scanning system is a non-invasive, safe, objective and reliable method to obtain 3D facial images of children during growth and development.

## SP 101

Thursday, June 17 – at 11:30

### ASSESSMENT OF THE OCCLUSAL CANT IN THE FRONTAL PLANE

**O Popoviciu**, M Popescu, E Ionescu, Department of Orthodontics and Dentofacial Orthopedics, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

**AIM:** While facial asymmetry is detected, to some degree, even in 'normal' subjects, its quantification is of particular interest in patients seeking orthodontic treatment. Some asymmetric features might be masked by soft tissue or postural habits and, in the absence of thorough examination, become apparent only during the finishing stages of treatment. The aim of this research was to evaluate the cant of the occlusal plane in patients seeking orthodontic treatment, and to verify the existence of significant angular differences between patients with a unilateral/bilateral crossbite compared with those with a neutral transverse occlusion.

**MATERIALS AND METHOD:** Initial posteroanterior (PA) and lateral cephalograms of 28 consecutive adult subjects were traced and angular and linear cephalometric measurements were obtained (including SNA, SNB, occlusal and mandibular angles, A0-

B0 distance on the lateral cephalogram and maxillo-mandibular/midline angle and cant of the occlusal plane on the PA cephalogram). The patients were divided into three groups, based on the lateral transverse occlusal relationship (neutral,  $n = 18$ , unilateral crossbite  $n = 7$ , bilateral crossbite  $n = 3$ ).

**RESULTS:** Three subjects in the neutral group had no occlusal cant. The angulation of the occlusal plane ranged from 2 to 9 degrees (average  $4.3^\circ$ ). The average subsample occlusal plane angle was 3 degrees for the neutral, 7 degrees for the unilateral crossbite and 4.5 degrees for the bilateral crossbite. There were no statistically significant differences between subsamples, due to the small number of subjects.

**CONCLUSION:** In order to better plan and manage orthodontic treatment, determination of the occlusal cant should be routinely performed, especially in subjects with transverse jaw discrepancies, where it might also serve as benchmark for treatment results.

## SP 102

Thursday, June 17 – at 11:30

### FACIAL MORPHOLOGY CHANGES IN THE MIXED DENTITION PERIOD FROM 8 TO 9 YEARS OF AGE – A THREE-DIMENSIONAL EVALUATION

**M Verdenik**<sup>1</sup>, D Selan<sup>1</sup>, A Zhurov<sup>2</sup>, S Richmond<sup>2</sup>, M Ovsenik<sup>1</sup>,

<sup>1</sup>Department of Orthodontics, University of Ljubljana, Slovenia and <sup>2</sup>Dental Health and Biological Sciences, Dental School, Cardiff University, Wales

**AIM:** To quantify facial morphology differences in a population of Slovenian children in the mixed dentition period, aged from 8 to 9 years, using three-dimensional (3D) laser scanning technology.

**SUBJECTS AND METHOD:** Forty-two children (19 girls, 23 boys) participated aged 8 years. One year later 30 children attended the follow-up (17 girls, 13 boys). Facial images were taken using a 3D imaging system, consisting of two high resolution laser cameras (Minolta Vivid VI910 3D). Average 3D facial shells were derived from the female and male attendees. The 9-year-old facial shell was superimposed on the 8-year-old shell. A colour map was used to highlight the differences between the two average male/female shells for the 8 and 9-year-old children.

**RESULTS:** The difference between the average facial shells after one year of growth and development was for girls (mean =  $0.32 \pm 0.25$  mm) and for boys (mean =  $0.32 \pm 0.30$  mm). The greatest soft tissue changes from 8 to 9 years of age were in the nasal and temporal area as well as in the ramus ascendens area of the lower jaw for girls, while in the boys it was in the lower jaw and premaxillary regions.

**CONCLUSION:** 3D laser soft tissue technology is suitable for monitoring growth and developmental changes in a population. Furthermore, facial morphology changes can be determined non-invasively, reliably and more objectively.

## SP 103

Thursday, June 17 – at 11:30

### BUCCALLY BLOCKED OUT CANINES – CLINICAL EVIDENCE OF THE THREE-BRACKET RELATIONSHIP

**M Svejda**, H-P Bantleon, Department of Orthodontics, Dental School of Vienna, Austria

**AIM:** Clinical observation of the levelling and aligning of buccally blocked out canines. It was hypothesized that, due to the theory of the three-bracket relationship, space opening should be a combination of protrusion of the anterior segment and distal movement of the lateral segment.

**SUBJECTS AND METHOD:** Ten patients with blocked out canines in the upper jaw were bonded with self-ligating brackets (Damon III) and a 0.014 inch Copper-NiTi wire was used for the whole levelling period. To follow the movement of the teeth, special marks were set on the archwire mesial to the bracket of the lateral incisor and distal to the bracket of the first premolar. Photographs of were taken every 4 weeks. Plaster casts were made every 8 weeks and the wire was removed and reactivated in tempered water.

Three-dimensional scans of the plaster casts were used for superimposition. Additional measurements were performed on two-dimensional occlusograms.

**RESULTS:** Distal movement of the first premolars occurred and reached values up to 4.08 mm (SD 0.47 mm), while mesial movement of the lateral incisors was within 1.96-2.76 mm (SD 1.16 mm). For unilateral blocked out canines, distal movement of the first premolar on the side of the blocked out canine showed the highest values.

**CONCLUSION:** Space opening in situations of buccally blocked out canines takes place by mesial movement of the lateral incisors as well as distal movement of the first premolars. Therefore these canines should be immediately bonded and integrated in the fixed appliance to take advantage of the special geometry and favourable vertical and horizontal forces due to the three-bracket relationship.

## SP 104

Thursday, June 17 - at 12:15

### THE PREVALENCE OF WHITE SPOT LESIONS AFTER ORTHODONTIC FIXED APPLIANCE THERAPY

T Tomažević<sup>1</sup>, M Drevenšek<sup>2</sup>, S Burnik<sup>3</sup>, R Kosem<sup>3</sup>, <sup>1</sup>Department of Pediatric and Preventive, Dentistry, University of Ljubljana and Departments of <sup>2</sup>Orthodontics and <sup>3</sup>Pediatric and Preventive Dentistry, University Medical Centre, Slovenia

**AIM:** White spot lesions (WSL), an undesirable side-effect of orthodontic treatment, are the result of prolonged plaque accumulation around the elements of fixed appliance. The prevalence of WSL after orthodontic therapy is reported to be 50-97 per cent. Laser fluorescence measurement (DIAGNOdent; DD) is an additional diagnostic measure for quantification of carious lesions. The aim of this study was to determine the prevalence of WSL after orthodontic fixed appliance therapy and to complement the findings of visual examination with DD measurements.

**SUBJECTS AND METHOD:** Forty-four patients who had finished fixed appliance therapy were consecutively included, with 1177 teeth examined. Visual examinations were first performed on wet surfaces; three surfaces per tooth were examined: mesiobuccal, buccal and distobuccal. The same procedure was repeated after air-drying with compressed air for 5 seconds. Lesions were classified according to the criteria of Ekstrand et al. (1997), modified by Aljehani et al. (2006). Lesions with enamel breakdown or greyish discolouration (code 3) were restored. DD was used to assess the lesions coded 1 and 2.

**RESULTS:** Mouth and tooth prevalence of WSL were 90.1 and 39.2 per cent, respectively. WSL were detected on 543 (15.4%) of all inspected surfaces; 124 (22.8%) were coded 1, 403 (74.2%) were coded 2 and 16 (3.0%) were coded 3. The average DD readings for WSL coded 1 and 2 were 1.5 (SD 2.5) and 4.6 (SD 7.1), respectively; the corresponding ranges were 0-20 and 0-43, respectively.

**CONCLUSION:** In the examined population, the prevalence of WSL after orthodontic treatment with fixed appliances was high, as assessed by visual examination, complemented with DD measurements.



### Class III Malocclusion

SP 105

Thursday, June 17 - at 12:15

#### THREE-DIMENSIONAL CEPHALOMETRY FOR DIAGNOSIS AND TREATMENT PLANNING OF IMPACTED CANINES

**I M Dagsuyu**<sup>1</sup>, S Dagistanli<sup>2</sup>, T Gurbuz<sup>3</sup>, M A Sumbullu<sup>2</sup>,  
Departments of <sup>1</sup>Orthodontics, <sup>2</sup>Radiology and <sup>3</sup>Pedodontics,  
Faculty of Dentistry, Ataturk University, Erzurum, Turkey

**AIM:** An impacted tooth is defined as one that fails to erupt at its appropriate site in the dental arch, within the normal period of growth. Maxillary canines are the most frequently impacted teeth, after third molars. Conventional radiographic techniques can be somewhat limited in the visualization of impacted teeth, due to problems involving superimposition and the complex nature of related dentoalveolar structures. Three-dimensional (3D) volumetric imaging with cone-beam computed tomography (CBCT) has the potential to define correct localization of impacted canines, using spatial relationships. The aim of this investigation is to propose a new 3D cephalometric analyzing system for diagnosis and treatment planning of impacted maxillary canines.

**MATERIALS AND METHOD:** Diagnostic CBCT records of 25 subjects with uni- or bilateral impacted maxillary canines were retrieved from a CBCT archive of 550 patients. All records were obtained with a NewTom 3G FP CBCT scanner. With an orthodontic image processing software (Simplant Ortho Pro v2.0), a 3D study model was reconstructed and cephalometric analyses were realized for each data set. Specialized landmarks, lines and reference planes were designed for the analysis of impacted maxillary canines to the related dentoalveolar and craniofacial structures. Axial inclination, crown rotation, bucco-palatal localization, stereoscopic position to the reference planes, proximity to the adjacent teeth and overlapping rate, root morphology and development, resorptive potential, peripheral bone density, impaction depth and distance to the ideal position, amount of available space and crown size of impacted maxillary canines were measured and analyzed. Craniofacial pattern and relative morphological and spatial relationships of other dentoalveolar structures were also evaluated.

**CONCLUSION:** Comprehensive diagnosis is the key for successful treatment planning to obtain maximum therapeutic results.

SP 106

Friday, June 18 - at 8:30

#### CORRELATION BETWEEN TOOTH SIZE AND JAW LENGTH IN CLASS III AND CLASS I MALOCCLUSION SUBJECTS

**H R Fattahi**, S Zavareian, F Nasiri, Orthodontic Department,  
Shiraz Dental School, Iran

**AIM:** To investigate the correlation between jaw length and the sum of the mesiodistal width of six and 12 teeth in the maxillary and mandibular arches of Class III and Class I patients.

**SUBJECTS AND METHOD:** This cross-sectional study included 120 patients (50 Class I, 70 Class III), 40 with mandibular excess and 29 with maxillary deficiency. The inclusion criteria were permanent dentition with no caries, proximal restorations, attrition or dental abnormalities. The greatest mesiodistal tooth width from the first molar to the first molar was measured on study models using a digital calliper with an accuracy of 0.01. The correlation between the sum of the mesiodistal size of six and 12 teeth to jaw length indices (Co-A and Co-Gn) was analyzed by Pearson test.

**RESULTS:** There was no significant correlation between arch length indices and the sum of six or 12 teeth. The correlation between Co-Gn and Co-A was significant ( $P < 0.001$ ) in both the Class III (maxillary deficiency and mandibular excess) and Class I patients. Although the correlation between the size of six teeth in the upper and lower arches was significant only in the maxillary deficiency group ( $r = 0.4$ ,  $P < 0.01$ ), the correlation was significant in all groups for 12 teeth.

**CONCLUSION:** There is almost always a high correlation between maxillary and mandibular arch length, but not between arch length and 6 or 12 teeth, in either jaw.

SP 107

Friday, June 18 - at 8:30

#### A COMPARISON OF FACEMASK TREATMENT EFFECTS ACCORDING TO CERVICAL VERTEBRAE MATURATION

**B Yang**, S Mlee, M S Yoon, J H Lee, Department of Clinical Orthodontics, Graduate School of Clinical Dentistry, Hallym University, Korea

**AIM:** Use of a maxillary protraction appliance, and the decision concerning appropriate treatment timing are important. The aim of this research was to compare the effects of facemask treatment according to the cervical vertebrae maturation (CVM) method. **SUBJECTS AND METHOD:** Thirty subjects treated with a facemask were divided into two groups according to CVM.

**RESULTS:** In CVM 1~2 group, vertical and angular measurements showed statistically significant differences, except for the palatal plane. For horizontal measurements, the CVM 3~5 group showed statistically significant differences and all angular measurements were statistically significantly different except for SNB, mandibular plane before and after treatment. The skeletal change in the CVM 1~2 group was larger than in the CVM 4~5 group, but the dental effects in the CVM 3~5 group were larger than in the CVM 1~2 group. These results show that the effect of maxillary protraction was mostly skeletal in the pre-pubertal and pubertal growth peak periods, and the dental effects were larger after the growth peak period.

**CONCLUSION:** The skeletal change in the CVM 1~2 group and the dental effects in the CVM 3~5 group contributed to overjet correction. Treatment with a facemask is more effective when it is commenced before the growth spurt. CVM is useful for evaluating treatment timing with the advantage of reducing radiation exposure.

SP 108

Friday, June 18 - at 8:30

#### EPIDEMIOLOGIC STUDY OF MANDIBULAR PROGNATHISM IN KOREAN FAMILIES

**A Lee**, Department of Orthodontics, The Institute of Oral Health and Science, Samsung Medical Center, Seoul, Korea

**AIM:** In spite of many studies the aetiology of mandibular prognathism has not been clarified. Recently due to the rapid increase in the number of Korean patients requiring surgical correction of mandibular prognathism a sufficient number of subjects is available to the genetics of this malocclusion. The aim of this study was to verify the genetic contribution to mandibular prognathism in Korean families.

**SUBJECTS AND METHOD:** The probands were 92 patients (47 males, 45 females) with mandibular prognathism who had orthognathic surgery or orthodontic treatment. Each subject completed the questionnaire of the pedigree chart comprising three generations of family members on the facial profile. Heritability ( $h^2$ ) in first degree relatives was evaluated with Falconer's formula.

**RESULTS:** The average SNA, SNB and ANB angles were:  $81.2 \pm 3.2$ ,  $84.0 \pm 4.0$  and  $-2.9 \pm 3.1$  degrees, respectively. All values, except for SNA, indicated a skeletal Class III malocclusion. A total of 3393 familial members were examined, of which 2.5 per cent had mandibular prognathism. At least one mandibular prognathic member, other than the proband, was found in 42.4 per cent of families. The affected ratio of first-degree relative was 11.1 per cent, which was much higher than that of the second- or third-degree relatives (1.6 and 1.7%, respectively). Falconer's heritability of mandibular prognathism in the first-degree relatives was 32.6 per cent.

**CONCLUSION:** There is a significant genetic influence in the aetiology of mandibular prognathism in Korean families.

### SP 109

Friday, June 18 - at 8:30

#### EARLY TREATMENT OF PSEUDO CLASS III MALOCCLUSION – A 10-YEAR FOLLOW-UP STUDY

I M Anderson, A B M Rabie, R W KWong, University of Hong Kong, SAR China

**AIM:** Pseudo Class III malocclusion subjects have anterior crossbite, and are defined as a functional forward displacement of the mandible due to retroclined maxillary incisors. About 5 per cent of the Chinese population is affected by a Class III malocclusion, and more than half are pseudo-Class III. Early treatment of pseudo Class III malocclusions has a number of advantages: it facilitates the eruption of canines and premolars into a Class I occlusion; it eliminates traumatic occlusion to the incisors; provides a normal environment for growth of the maxilla; and often improves the child's self esteem. The aim of this study was to investigate the long-term outcome of simple fixed appliance treatment of a pseudo Class III malocclusion.

**MATERIALS AND METHOD:** Lateral cephalograms of 27 consecutive young patients obtained before and after active treatment with a 2 X 4 appliance, and at the 10-year follow-up.

**RESULTS:** Eighteen patients successfully recalled for the 10-year follow-up had an average pre-treatment overjet of  $-1.7$  mm. Immediately following treatment the average overjet was 3.2 mm, which reduced to 2.2 mm at the 10-year follow-up. Only one patient presented with a reverse overjet.

**CONCLUSION:** The majority of pseudo-Class III patients who receive early intervention with a 2 X 4 appliance, will be stable in the long term and will not need further orthodontic treatment. It is recommended though that treatment of an anterior crossbite at an early age should be approached cautiously.

### SP 110

Friday, June 18 - at 8:30

#### PREDICTION OF THE OUTCOME OF TREATMENT OF CLASS III MALOCCLUSION – A SYSTEMATIC REVIEW

P Fudalej<sup>1</sup>, M Dragan<sup>2</sup>, B Wedrychowska-Szulc<sup>3</sup>, <sup>1</sup>Institute of Mother and Child, Warsaw, <sup>2</sup>Warsaw Medical University and <sup>3</sup>Private practice, Szczecin, Poland

**AIM:** To systematically review the orthodontic literature to assess the possibility of a prediction of outcome of orthodontic treatment in subjects with Class III malocclusions.

**MATERIALS AND METHOD:** Two authors, using appropriate MeSH and free search terms, independently searched the PubMed, Embase, Cochrane Central, and Lilacs databases as well as hand-searched citations from the relevant articles. Among other data, sample ethnicity, treatment method, age at the start and completion of treatment, age at follow-up, outcome measures, and identified predictors were extracted from the relevant studies. An assessment of methodological quality of the studies was performed.

**RESULTS:** Two hundred and thirty two potentially relevant citations were identified of which 14 met the inclusion/exclusion criteria. In the included studies 683 subjects were examined on average 6.3 years after treatment with various orthodontic/orthopaedic methods. The age at start of treatment was from 5.6 to 12.4 years and at follow-up from 15.8 to 22 years. The heterogeneity of the samples and treatment methods prevented the carrying out of a meta-analysis. Thirty-eight different predictors of treatment outcome were identified, 35 cephalometric and three derived from analysis of study casts. Prediction models comprising 3-4 predictors were reported in most studies. However, only two studies shared more than one predictor. Gonial angle was most frequently identified (5 publications). Study quality was low (11) or medium (3).

**CONCLUSION:** The large variety of predictors and disparities among developed prediction models indicate that the possibility of prediction of outcome of orthodontic/orthopaedic treatment of Class III malocclusion is unsubstantiated.

### SP 111

Friday, June 18 - at 8:30

#### EFFECTS OF PALATAL PLANE ROTATION ON SOFT TISSUE CHANGES IN DOUBLE JAW SURGERY

Y Kim<sup>1</sup>, C S Kim<sup>2</sup>, Y-A Kook<sup>1</sup>, J U Park<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Cranio-maxillofacial Surgery, St. Mary's Hospital, Medical College, The Catholic University of Korea, Seoul, Korea

**AIM:** For the correction of skeletal Class III malocclusions, a combination of maxillary clockwise rotation (posterior maxillary impaction) and mandibular setback is one of the most common double jaw orthognathic surgical procedures. The purpose of this study was to assess soft tissue changes in the upper lip and nose area following double jaw surgery according to the degree of rotation.

**SUBJECTS AND METHOD:** Fifty-four patients who underwent maxillary posterior impaction and mandibular setback surgery for the correction of skeletal Class III malocclusions were grouped by the degree of palatal plane rotation: group 1 ( $0-3^\circ$ ), group 2 ( $3-6^\circ$ ), and group 3 ( $6-9^\circ$ ). Lateral cephalograms taken 1 month before (T1) and 6 month after (T2) surgery were traced and measurements were performed.

**RESULTS:** With horizontal backward movement at ANS and U1E and accompanying movement of the upper lip, a progressive increase was observed from group 1 to 3, and statistically significant differences were found between groups (U1E;  $P < 0.01$ , Stms;  $P < 0.05$ ). A moderate correlation was identified between soft tissue movement of the upper lip and underlying skeletal movement (point A, U1E) in the horizontal plane ( $P < 0.01$ ) but not in the vertical plane. The ratios of soft to hard tissues changes at Stms to U1E increased from group 1 (40.3%) to group 2 (181.9%) but decreased in group 3 (106.8%).

**CONCLUSION:** The ratio of soft to hard tissue changes specific to the degree of rotation may improve the accuracy of predicting treatment results.

### SP 112

Friday, June 18 - at 8:30

#### MANAGEMENT OF SKELETAL CLASS III PATIENTS

V Tarmure<sup>1</sup>, A Serbanescu<sup>1</sup>, G Baciut<sup>2</sup>, O Almasan<sup>3</sup>, M Iancu<sup>4</sup>, Departments of <sup>1</sup>Orthodontics, <sup>2</sup>Cranio-Maxillofacial Surgery, <sup>3</sup>Prosthodontics and <sup>4</sup>Medical Informatics, University of Medicine and Pharmacy 'Iuliu Hatieganu', Cluj-Napoca, Romania

**AIM:** To assess the management of skeletal Class III malocclusion patients from 2006-2009.

**SUBJECTS AND METHOD:** Twenty-five patients (14 females, 11 males) with a Class III skeletal malocclusion, aged 6 to 23 years. Lateral cephalometry was used to establish the skeletal pattern before and after treatment. Skeletal vertical and horizontal relationships were evaluated according to Tweed, Sassouni and Hasund/Segner. The Statistica 7.0 program, applying the prin-

principal component analysis (PCA) and the Varimax method, were used to determine the correlation between different variables.

**RESULTS:** Associated anomalies were present (mandibular shift 20%, asymmetries 20%, crowding 40%, canine impaction 4%, open bite 4%, tooth transposition 4%). Nineteen patients (76%) were treated without extractions; six (24%) with extractions (premolars or first molars). Six patients (24 per cent) had orthodontic treatment and orthognathic surgery. The PCA method suggested that anterior face height, Wits appraisal, SNA, antero-posterior intermaxillary relationship (ANB), IMPA, FMIA and overjet vary according to the profile type (Sassouni) and Z angle; SNB, SNA and age according to lower face angulation (NSL-ML) and FMA, NL-ML and gonial angle (Ar-Go-Me) according to SNB angle.

**CONCLUSION:** Stepwise analysis influences the individual treatment of skeletal Class III malocclusions, which predicts the option for orthodontic treatment or combined orthodontic and orthognathic treatment. Treatment planning is more difficult with the increased degree of skeletal and dental contribution to the malocclusion. The option for a surgical or non-surgical approach is challenging, depending on the extent of the skeletal deformity, the degree of dental and soft tissue compensations, and the patient's concern.

### SP 113

Friday, June 18 - at 8:30

#### EFFECTS OF A FACEMASK COMBINED WITH A CHINCAP ON MAXILLARY GROWTH IN SKELETAL CLASS III SUBJECTS

**H N İşcan**, E Çakır, D İşcan, Department of Orthodontics, Faculty of Dentistry, Gazi University, Ankara, Turkey

**AIM:** To investigate whether posterior rotation of the mandible can be corrected with the use of a chin-cap combined with a face-mask, and if the sagittal maxillary growth will be affected.

**SUBJECTS AND METHOD:** Twenty-nine skeletal Class III patients with maxillary retrusion and normal vertical skeletal growth patterns, treated with rapid palatal expansion (RPE; acrylic cap splint type of Hyrax) and maxillary protraction with a Petit type facemask applying a force of 400 g on each side. In group 1 (16 patients) a posterior high-pull chin-cap force of 120 g on each side was applied to the chin-cap part of the facemask by means of hooks for a mean treatment period of 0.82 years, whereas in group 2 (FM) the period was 0.89 years. Pre- and post-treatment lateral cephalograms were used to study the changes of six landmarks, and maxillary rotation using Björk's maxillary superimposition, three linear and five angular measurements were made. Three-way ANOVA with repeated measurement in time was used to evaluate the data. Intragroup comparison was controlled by Duncan multiple comparison test when necessary.

**RESULTS:** No statistically significant difference was found between the groups except for the mandibular plane angle, which was increased by 1.88 degrees in group 2.

**CONCLUSION:** Maxillary growth was not affected by adding a high-pull chin-cap force to a maxillary protraction facemask to prevent mandibular posterior rotation.

### SP 114

Friday, June 18 - at 8:30

#### CEPHALOMETRIC ASSESSMENT OF CHANGES IN THE FACIAL SKELETON IN CHILDREN WITH MANDIBULAR PROGNATHISM

**M Pospieszńska**, University of Medical Sciences, Poznan, Poland

**AIM:** To evaluate changes in the morphology of the facial skeleton in children with clinically diagnosed mandibular prognathism.

**SUBJECTS AND METHOD:** Thirty-five children with mandibular prognathism ranging in age from 6 to 15 years. The cephalometric examination was carried out on lateral cephalogram of the head by means of 19 angular and 18 linear measurements. The measurements were compared with developmental norms formulated by Droschl, and statistically evaluated with a Student's t and Wilcoxon's tests.

**RESULTS:** Among the cranial angles the most significant changes were noticed on sella turcica (NSAr), which decreased on average

by 3.5 degrees and gonial angle (ArGoMe) that increased by 4.5 degrees. The angles defining the sagittal position of the mandible in relation to the base of the anterior cranial fossa (SNB) increased by 3.5 degrees, SNPog increased by 3.9 degrees, the angle defining the sagittal relationship between the mandible and the maxilla (ANB) was reduced by 3.7 degrees, and the angle illustrating the vertical relationship between them (ML-NL) increased by 3.7 degrees. The linear measurement of mandibular base length showed a mean increase of 6 mm (from 2 to 16 mm) in relation to the norm in all subjects. In 15 subjects the mandibular ramus was also lengthened. Maxillary base length showed a shortening in 11 subjects. Vertical linear measurements revealed an increase in lower face height in all subjects.

**CONCLUSION:** Children with mandibular prognathism display excessive sagittal growth of the mandibular body, often accompanied by lengthening of the mandibular ramus.

### SP 115

Friday, June 18 - at 8:30

#### TONGUE VOLUME AND POSTERIOR AIRWAY SPACE CHANGES IN CLASS III TWO JAW ORTHOGNATHIC SURGERY

**O Uslu Akçam**, H Gokalp, Department of Orthodontics, School of Dentistry, Ankara University, Turkey

**AIM:** A Class III skeletal dentofacial deformity may be the result of mandibular prognathism and/or maxillary deficiency. Surgical correction of Class III deformities can be achieved by mandibular set-back alone, maxillary advancement, or combined bimaxillary surgery. The posterior airway space is delimited by hard and soft tissues with anomalies that may produce alterations in volume. The purpose of this study was to evaluate changes in airway space after surgical correction of a Class III skeletal dentofacial deformity and to determine if orthognathic surgery alters airway space and improves obstructive sleep apnoea.

**SUBJECTS AND METHOD:** Twenty-four Class III patients (19 females, 5 males), 10 mandibular set-back or maxillary advancement patients (mean age 19.6 years) and 14 bimaxillary surgery patients (mean age 21.6 years). Lateral cephalometric radiographs were traced before and 6 months after surgery. Steiner and McNamara analyses, posterior airway (PNS-PPW1, SPT-PPW2, E-PPW3, PNS-SPT) and tongue volume measurements (TH, Tt-Eb) were performed. The data obtained was analyzed statistically by repeated measurement analysis of variance and Duncan's test.

**RESULTS:** The interaction between group and treatment time was statistically significant regarding PNS-PPW1 measurement ( $P < 0.05$ ) that demonstrated that the change in PNS-PPW1 dimension was statistically different between the groups during treatment. Before treatment, PNS-PPW1 was statistically different between the treatment groups ( $P < 0.05$ ) and it increased in the bimaxillary surgery group after surgery ( $P < 0.05$ ). The difference between the groups was statistically insignificant regarding SPT-PPW2, E-PPW3, PNS-SPT and TH measurements. However, Tt-Eb was statistically different between the groups post-surgery ( $P < 0.01$ ).

**CONCLUSION:** Surgical correction of a Class III skeletal dentofacial deformity alters posterior airway dimensions.

### SP 116

Friday, June 18 - at 8:30

#### STRUCTURAL CHANGE OF THE HYOID BONE AND UPPER AIRWAY AFTER ORTHOGNATHIC SURGERY USING THREE-DIMENSIONAL COMPUTED TOMOGRAPHY

**H-S Yu**, Y S Lee, S H Ahn, I K Paik, Department of Orthodontics, Yonsei University College of Dentistry, Seoul, Korea

**AIM:** To find structural changes of the hyoid bone and upper airway after mandibular setback osteotomy for skeletal Class III patients with anterior open bites.

**SUBJECTS AND METHOD:** Twelve patients diagnosed with a skeletal Class III malocclusion with an open bite who underwent a mandibular setback osteotomy. Skeletal measurements were analysed using three-dimensional imaging of the hyoid bone and

upper airway. Ten subjects with a 'normal' occlusion comprised the control group.

**RESULTS:** 1. In the open bite group, the hyoid bone was positioned anteriorly, compared with the controls ( $P < 0.05$ ). 2. Post-surgery the hyoid bone was positioned upward, posterior, left displacement, but this was not statistically significant. 3. The angle between the hyoid plane and mandibular plane in Class III group pre-surgery was greater than in the controls ( $P < 0.05$ ), and the difference increased significantly post-surgery ( $P < 0.01$ ). 4. In the Class III subjects, the volume of the upper airway decreased significantly post-surgery ( $P < 0.001$ ). The volume of the upper airway was significantly smaller than in the control group both pre- and post-surgery ( $P < 0.001$ ).

**CONCLUSION:** In Class III open bite subjects, the angle between the hyoid plane and mandibular plane was greater than in the normal group. Upper airway volume in the Class III patients was smaller than in the controls and volume was decreased after orthognathic surgery.

### SP 117

Friday, June 18 - at 8:30

#### LONG-TERM OBSERVATION OF MASSETER MUSCLE CHANGES IN FACIAL ASYMMETRY

**H-S Yu, S H Ahn, I K Paik C S Lee,** Department of Orthodontics, Yonsei University College of Dentistry, Seoul, Korea

**AIM:** To examine how the soft tissues are changed and adjusted by comparing and analyzing changes in the masseter muscle of patients with a skeletal Class III malocclusion with facial asymmetry 1 and approximately 4 years after bimaxillary orthognathic surgery.

**SUBJECTS AND METHOD:** Eight patients who were diagnosed with skeletal facial asymmetry and who had undergone surgery comprised the asymmetry group, and 10 subjects with a 'normal' jaw and normal occlusion were selected as the controls.

**RESULTS:** 1. Volume, maximum cross-sectional area (CSA) and thickness of the maximum CSA all increased in the asymmetry group post-surgery. The width of the masseter muscle increased, and the length of the muscle decreased, on average. 2. The most related variable to the volume of the masseter muscle 4 years after mandibular setback osteotomy was the maximum CSA of the masseter muscle and the thickness of the maximum CSA ( $P < 0.01$ ). 3. Pre-surgery there were significant differences between the asymmetry and control group in volume, masseter muscle angle, maximum cross-sectional area, thickness of the maximum CSA, width and length of the masseter muscle. However, 4 years post-surgery there were no significant differences in the masseter muscles between the groups.

**CONCLUSION:** The skeleton of patients with facial asymmetry was stable 1 year post-surgery. The shape of the masseter muscle in the asymmetry group continuously changed to be similar to that in the normal group one year after surgery.

### SP 118

Friday, June 18 - at 9:15

#### COMPARISON OF FACEMASK OR REVERSE CHIN CUP THERAPY IN TREATING CLASS III MALOCCLUSIONS

**A Jamilian<sup>1</sup>, R Showkatbakhsh<sup>2</sup>, M Toliat<sup>1</sup>, T Taban<sup>1</sup>, Z Emami<sup>1</sup>,** Department of Orthodontics, <sup>1</sup>Islamic Azad University and <sup>2</sup>Shahid Beheshti University, Tehran, Iran

**AIM:** To compare the dentoskeletal effects of facemask or and reverse chin cup therapy in treating Class III malocclusions and maxillary deficiency in growing patients.

**SUBJECTS AND METHOD:** Twenty patients with maxillary deficiency and a normal mandible. Ten patients (mean age 9.5 years) were treated with reverse chin cup. This appliance consisted of a high pull cap, a chin cup with two vertical hooks, a removable appliance on the upper jaw, and two elastics that connected the hooks of the cup to the hooks of removable appliance. The reverse chin cup was used for  $24 \pm 2$  months. The other patients

(control group) with a mean age of 9.8 years were treated with a facemask. The treatment time was  $26 \pm 3$  months. Dental pantomograms, lateral cephalograms, and photographs were taken at the beginning and end of treatment in both groups. A t-test was used for statistical evaluation of the data.

**RESULTS:** SNA was significantly increased in both groups ( $P < 0.001$ ). No statistically significant difference was seen between the groups ( $P < 0.3$ ). ANB and soft tissue profile were improved in both groups while IMPA was significantly decreased between the experimental and control groups. However, comparison of the data showed no statistically significant difference between the groups.

**CONCLUSION:** Both reverse chin cup and facemask therapy were effective in the treatment of maxillary deficiency. However, a reverse chin cup seems more favourable for patients due to its smaller size.

### SP 119

Friday, June 18 - at 9:15

#### CERVICAL VERTEBRA ANATOMY IN PATIENTS WITH NORMAL OCCLUSION AND CLASS III MALOCCLUSION

**M Pop Stefanova-Trposka, V Korunoska-Stevkovska, G Circieva-Cuckova,** Department of orthodontics, Faculty of dentistry, Dental clinical center, Skopje, Former Yugoslav Republic of Macedonia

**AIM.** To compare the dimensions of the first and second cervical vertebra, intervertebral space, head posture and anterior and posterior facial height in children with normal occlusion and class III malocclusion.

**MATERIAL AND METHOD.** We analyzed lateral cephalometric radiographs from 100 patients from both sexes, aged 10-13 years. Fifty of them were with normal occlusion, skeletal class I, and 50 with class III malocclusion and skeletal class III. On each radiograph we measured five variables for cervicovertebral anatomy, three for intervertebral space, five for head posture, seven linear and angular mandibular variables and anterior and posterior facial heights.

**RESULTS.** It was found that the children with malocclusion class III had forward mandibular rotation and horizontal type of growth. The anterior height of atlas (V), the dorsal heights of atlas (D1) and axis (D2), and the height of dens axis (DENS) were highly significantly larger ( $p < 0.001$ ) than the class I group. The intervertebral space showed weak significant differences ( $p < 0.05$ ). Children with class III malocclusion did not have extended head posture (the craniocervical angle NSL/VER and the craniocervical angle NSL/OPT were highly significantly smaller than the class I group ( $p < 0.001$ ), while the OPT/HOR was significantly increased in malocclusion class III group ( $p < 0.01$ ).

**CONCLUSION.** The children with malocclusion class III showed forward mandibular rotation with horizontal growth of the mandible, larger dimensions of the first and second cervical vertebrae, small craniocervical angle and flexion of the head.

### SP 120

Friday, June 18 - at 9:15

#### AN ETHNIC BACKGROUND-BASED CEPHALOMETRIC STUDY IN HUNGARIAN AND JAPANESE CLASS III SUBJECTS

**I Csiki<sup>1</sup>, A Végh<sup>1</sup>, M Hamada<sup>2</sup>,** Department of Orthodontics and Orofacial Orthopaedics, Heim Pal Children's Hospital, Budapest, Hungary and <sup>2</sup>Department of Orthodontics, Osaka Dental University, Japan

**AIM:** To characterize the morphological differences of craniofacial structure in two ethnic groups.

**MATERIALS AND METHOD:** Cephalometric data of Hungarian and Japanese patients with a Class III malocclusion diagnosed as requiring orthognathic surgery were compared. The study sample consisted of standardized pre-treatment cephalometric radiographs of 28 (14 males, 14 females) Hungarians (24 year 1 month) and 28 (14 males, 14 females) Japanese (22 years 11 months). Measurements were performed using the method of Ricketts

with FR Win 5.0 for Windows®. Ten linear and angular variables were analysed. Basic statistics, F-, two sample t-, Mann-Whitney and Wilcoxon tests were used.

**RESULTS:** The Japanese sample presented stronger negative values for facial convexity. Lower face height and mandibular plane angle showed increased values in the Japanese sample resulting in an increased lower facial third and a very accentuated open bite tendency. Decreased values of maxillary depth angle showed a retroposition of the upper jaw. On average, values for conical angle were smaller than normal, due to a forward position of the chin, especially in the Japanese group. Facial depth showed an increased average value, indicating a more forward mandible with reference to the cranial base in both groups. Mandibular angle (Xi) slightly differed from normal. The Hungarian group presented higher values, which can be considered as the upper limit of normal. A higher angle for the position of the mandibular ramus (ramus Xi position) was observed in both group. As expected, constant values (such as craniofacial deflection, facial axis) remained close to normal.

**CONCLUSION:** The null hypothesis that there is no difference between the data of the two groups was rejected. The results appear to suggest real differences in skeletal features between Hungarian and Japanese Class III surgical patients due to ethnicity and the nature of the samples

## SP 121

Friday, June 18 - at 9:15

### EFFECTS OF FACEMASK TREATMENT ON SAGITTAL PHARYNGEAL DIMENSIONS IN CLASS III SUBJECTS

**R Lione**, M Mucedero, M Rozzi, Department of Orthodontics University of Rome, Italy

**AIM:** The literature reports contrasting findings on the possibility of improving the sagittal airway dimension by means of maxillary protraction. The aim of this cephalometric study was to analyze the treatment and post-treatment craniofacial effects of a face-mask (FM) combined with a bite block (BB) with specific regard to the sagittal pharyngeal dimensions in Class III malocclusion subjects when compared with an untreated Class III control group.

**SUBJECTS AND METHOD:** The FM/BB group comprised 22 subjects (12 females, 10 males) with a mean age pre-treatment (T1) of  $8.9 \pm 1.5$  years, at the end of active treatment (T2) of  $10.5 \pm 1.3$  years, and post-treatment (T3) of  $12.6 \pm 1.9$  years. The treated group was compared with an untreated group of 14 subjects (6 females, 8 males) with a Class III malocclusion. The average age was  $7.6 \pm 1.4$  years at T1,  $9.8 \pm 1.9$  years at T2, and  $11.9 \pm 1.2$  years at T3. All subjects showed a prepubertal stage of skeletal growth (CS 1) at T1 and a post-pubertal stage (CS 4, CS 5, or CS 6) at T3. Comparisons of the T2-T1 and T3-T1 changes between the two groups were analyzed with the Mann-Whitney test.

**RESULTS:** During T2-T1, the treated group showed significant increments in maxillary skeletal variables; a significant increase in the vertical intermaxillary relationships was also found; no statistically significant differences were observed for any of the analyzed variables for upper and lower sagittal airway dimensions. During T3-T1 no statistically significant differences were found for the vertical skeletal relationships or for upper and lower sagittal airway dimensions.

**CONCLUSION:** The FM/BB protocol produced significantly favourable and post-treatment stable changes both in the maxilla and mandible in Class III subjects when compared with untreated controls. No significant changes in oro- and nasopharyngeal sagittal airway dimensions were induced by FM/BB therapy when compared with untreated subjects.

## SP 122

Friday, June 18 - at 9:15

### PSYCHOLOGICAL CHARACTERISTICS OF PATIENTS WITH SKELETAL CLASS III MALOCCLUSIONS

**A Kovalenko**, A Slabkovskaya, L Persin, N Drobysheva, Moscow State University of Medicine and Dentistry, Russia

**AIM:** To evaluate the psychological status of patients with skeletal Class III malocclusions with different degrees of facial deformities.

**SUBJECTS AND METHOD:** Forty-two patients aged 16-45 years with a skeletal Class III malocclusion were examined before and after orthognathic treatment. Thirty untreated subjects aged 20-30 years with harmonious faces comprised the control group. Diagnostics included photographic analysis of the face and psychological testing (EPI, Spilberger questionnaire and Q-sorting method). Based on photographic analysis, a Facial Changes Index (FCI) was developed to determine the degree of influence of the malocclusion on the facial aesthetics. A series of soft tissue parameters affecting facial changes as a result of a skeletal malocclusion was established. The normal range of these parameters was calculated analyzing photographs of a control group. The patients with a Class III malocclusion were divided into three groups according to the FCI scores minor (0-10), moderate (10-19) and severe (19 and more) changes.

**RESULTS:** Psychological testing before treatment showed statistically significant differences between the Class III malocclusion groups and the controls. The most apparent changes were found in the patients with severe facial changes. Psychological testing 6 months after orthognathic surgery showed statistically significant changes, with most of the parameters approaching normal values

**CONCLUSION:** 1. The FCI is a simple and fast way to determine the severity of facial changes due to a skeletal malocclusion. 2. The severity of facial changes in Class III skeletal malocclusions influences the psychological characteristics of patients. 3. Patients with severe facial changes experience some difficulties in social contacts tending to avoid leadership and group conflicts. 4. Orthognathic treatment can improve the quality of the social life of such patients.

## SP 123

Friday, June 18 - at 9:15

### MANDIBULAR FEATURES OF CLASS III PATIENTS WITH IMPLICATIONS ON TEMPOROMANDIBULAR DISORDERS

**M-A Bencze**<sup>1</sup>, V-M Milicescu<sup>2</sup>, E Teodorescu<sup>2</sup>, E Ionescu<sup>2</sup>, C-T Preoteasa<sup>3</sup>, Departments of <sup>1</sup>Oral Rehabilitation, <sup>2</sup>Orthodontics and <sup>3</sup>Oral Diagnosis, 'Carol Davila' University of Medicine and Pharmacy, Bucharest, Romania

**AIM:** Differences in the heights of the right and left mandible rami have been suggested as important skeletal problems associated with temporomandibular joint pathology. The aim of this research was to quantify the degree of mandibular asymmetry in Class III patients and its implications on the appearance of temporomandibular disorders (TMD).

**SUBJECTS AND METHOD:** Fifty patients (aged 11-17 years). Each patient underwent a full clinical examination and standard orthodontic records (plaster models, extra- and intraoral photographs, panoramic and lateral cephalometric radiographs) prior to orthodontic treatment. The subjects were divided into a control group of 25 Class I patients (13 females, 12 males) and a study group of 25 Class III patients (15 females, 10 males). The presence of TMD signs and symptoms was evaluated by an anamnestic questionnaire (joint pain in the previous month, joint sound and limitation of mouth opening) and a clinical examination. Mandibular asymmetry measurements (as described by Kjellberg) were performed on the panoramic radiographs.

**RESULTS:** Fifty six per cent of the control group and 48 per cent of the study group had no TMD symptoms. The asymmetry indices showed different values for the two groups, but no gender-related differences were observed. In some cases, the asymmetry value correlated with the degree of TMD.

**CONCLUSION:** Alteration of spatial intraarticular relationships due to mandibular asymmetry may contribute to the appearance of TMD, particularly in Class III patients.

### SP 124

Friday, June 18 - at 9:15

#### TOOTH-BORNE DISTRACTOR APPLICATION FOR ANTERIOR MAXILLARY POSITIONING IN CLEFT AND PALATE PATIENTS

**A Showkatbakhsh**<sup>1</sup>, F Pourdanish<sup>2</sup>, A Ghorbani<sup>2</sup>, A Jamilian<sup>3</sup>, M Behnaz<sup>1</sup>, Departments of <sup>1</sup>Orthodontics, <sup>2</sup>Maxillofacial Surgery and <sup>3</sup>Orthognathic Surgery, Shahid Beheshti Medical Science University, Tehran, Iran

**AIM:** Cleft lip and palate (CLP) patients require teamwork to correct their problem and to achieve the best result. Velopharyngeal insufficiency is a concern in maxillary advancement. In this study a new concept of tooth borne distractor application was used to avoid any side-effects of maxillary advancement for CLP patients.

**SUBJECTS AND METHOD:** Six adult CLP patients, average age 17 years, underwent pre-surgical orthodontics to provide space between the right and left upper first and second molars. An appliance constructed to be mounted on the upper first and second molars in combination with a jackscrew before surgery was tried in the patient's mouth. The surgical approach, under general anaesthesia was performed to have a transverse cut in the spaces provided between the upper molars. A buccal and labial cut with a nasal osteotomy to release the anterior part of the maxillary complex was accomplished and the distractor was then cemented in the mouth. After a period of seven days the distractor was opened 0.5 mm each day and after correct maxillary positioning, the consolidation period considered to have a good bone quantity and quality was established.

**CONCLUSION:** Since maxillary surgery might have unfavourable effect on velopharyngeal incompetence and for better muscular and soft tissue adaptation, this method might be an alternative for floating maxillary surgery in CLP patients.

### SP 125

Friday, June 18 - at 9:15

#### EFFECTIVENESS OF COMBINED SURGICAL-ORTHODONTIC TREATMENT: CEPHALOMETRIC ANALYSIS

**J Szyper-Szczurowska**<sup>1</sup>, J Zapala<sup>2</sup>, P Szczurowski<sup>2</sup>, B W Loster<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Maxillofacial Surgery, Collegium Medicum of the Jagiellonian University, Krakow, Poland

**AIM:** To analyse the facial skeleton of patients with a skeletal Class III malocclusion treated according to a combined surgical-orthodontic protocol.

**MATERIALS AND METHOD:** Thirty cephalograms of patients with a skeletal Class III malocclusion taken before treatment and 12 months post-surgery. The cephalograms were analyzed using the methods of Steiner (modified by Kaminek), Schwarz and Harvold. The results were statistically analysed using a Wilcoxon test. The level of significance was set at  $P < 0.05$ .

**RESULTS:** Statistically significance improvements regarding angles was observed: enlargement of SNA of 2.68 degrees ( $\pm 4.27$ ),  $p = 0.005$ ; reduction of SNB of 5.52 degrees ( $\pm 2.84$ ),  $p = 0.000$ ; enlargement of ANB of 8.26 degrees ( $\pm 4.25$ ),  $p = 0.000$ ; increase of Wits measurement of 10.33 mm ( $\pm 4.63$ ),  $p = 0.000$ ; increase in maxillary I-NS by approximately 5.30 degrees ( $\pm 2.91$ ),  $p = 0.005$ ; increase in I ML of about 6.36 degrees ( $\pm 4.25$ ),  $p = 0.000$ . There was also a statistically significant reduction in mandibular length gain which was evaluated according to Schwarz (average 4.72 mm;  $P = 0.000$ ) and Harvold (average 6.67 mm;  $P = 0.005$ ) analyses. In addition, there was an elongation of the maxilla estimated according to Schwartz (average 4.21 mm,  $P = 0.012$ ) and Harvold (average 3.9 mm;  $P = 0.008$ ) analyses.

**CONCLUSION:** The cephalometric measurements showed the efficiency of surgical treatment. The first stage of orthodontic treatment was adequate, because effective dentoalveolar decompensation was carried out.

### SP 126

Friday, June 18 - at 9:15

#### CORRELATION OF INCISOR INCLINATION AND THE SAGITTAL POSITION OF THE JAWS IN CLASS III SUBJECTS

**M Stošić**, P Janošević, D Stojanović, Department of Orthodontics, Medical Faculty, University of Niš, Serbia

**AIM:** To determine whether a correlation exists between inclination of the maxillary and mandibular incisors and the sagittal position of the jaws in subjects with Class III malocclusions.

**MATERIALS AND METHOD:** Fifty profile cephalograms of subjects with a Class III malocclusion. The inclination angles of the upper and lower incisors (I/SP and i/MP) were analyzed, as well as their correlation with the angles of maxillary and mandibular prognathism (SNA, SNB). The radiographs were divided into two groups, depending on the size of SNA and SNB (group I with smaller and group II with larger than standard values).

**RESULTS:** A weak, negative correlation was found between maxillary incisor inclination angle and maxillary prognathism angle in both groups (SNA  $> 82^\circ$ ,  $r = -0.53$ , SNA  $< 82^\circ$ ,  $r = -0.34$ ). A weak, negative correlation of the lower incisor inclination angle to the mandibular prognathism angle existed only when the values were less than 80 degrees ( $r = -0.45$ ).

**CONCLUSION:** Incisor inclination can be conditioned by the sagittal position of the jaws in subjects with a Class III malocclusion, which is of significance for treatment planning.

### SP 127

Friday, June 18 - at 9:15

#### FACIAL SOFT TISSUE PROFILE CHANGES FOLLOWING BILATERAL SPLIT SAGITTAL OSTEOTOMY AND SUBCONDYLAR MANDIBULAR SETBACK

S Momeni Danaei, **S Torkan**, F Khaje, M Ruinpeikar, Orthodontic Department, Shiraz School of Dental Sciences, Iran

**AIM:** Successful surgical planning and accurate prediction of orthognathic surgery outcomes include not only occlusal correction and a well-balanced skeletal relationship, but improvement of aesthetics and function. The aim of this retrospective study was to evaluate soft tissue changes following bilateral sagittal split osteotomy (BSSO) and subcondylar mandibular setback.

**MATERIALS AND METHOD:** Pre- and post-operative cephalometric variables were measured on lateral cephalometric radiographs of 35 individuals, 12 of whom had undergone BSSO surgery and 23 subcondylar surgery. Analysis of the measured variables was carried out by means of paired *t*- and independent *t*-tests.

**RESULTS:** Although only mandibular surgery was carried out, upper lip changes were significant. In BSSO cases, lip fullness was increased. Subcondylar cases showed an increase in lower face height and mentocervical angle that supports the idea of a shift toward a vertical facial pattern after this type of surgery. Overall BSSO resulted in more improvement in aesthetics compared with subcondylar surgery.

### SP 128

Friday, June 18 - at 9:15

#### CONSIDERATIONS ON THE THERAPEUTIC INTERVENTION IN MIXED DENTITION CLASS III CASES

**E Ionescu**, R Grigore, E Teodorescu, M Popa, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

**AIM:** The timing of orthodontic intervention in Class III anomalies is controversial. The aim of this study was to investigate orthodontic intervention in Class III subjects (especially pseudo Class III) by taking into consideration several advantages related to the response to orthodontic treatment, orthodontic means, duration and stability over time.

**SUBJECTS AND METHOD:** Thirty patients in the early mixed dentition (approximately 7-8 years of age), divided into two equal groups. The first group included patients with true mandibular

prognathism and the second patients with pseudo-prognathism, orthodontically treated during 2002–2005. The current clinical assessment conducted in 2009, a minimum of 5 years after the end of treatment, focused on the stability of the treatment result while maintaining the overbite.

**RESULTS:** For the 15 patients with pseudo prognathism, the results were maintained. Five of the skeletal Class III patients required further treatment, due to the association with other dentofacial anomalies, and four needed comprehensive orthodontic treatment for Class III anomalies.

**CONCLUSION:** Early treatment of Class III anomalies is beneficial to ensure harmonious growth of both jaws, to create conditions for maxillary development along with bite jumping, set the correct occlusal relationships and improve dentofacial functions, including aesthetics. Early orthodontic treatment, although limited, is indicated in skeletal Class III subjects as an element of interception for severe occlusal disturbances, or as palliative factor leading to a less severe clinical outcome in complex orthodontic interventions.

## SP 129

Friday, June 18 - at 9:15

### CORRELATION BETWEEN MANDIBULAR AND CRANIAL BASE FEATURES IN SKELETAL CLASS III ANOMALIES

**R Grigore**, E Ionescu, O Popoviciu, E Teodorescu, A Bencze, Departments of 1 Orthodontics and Dentofacial Orthopedics and 2 Oral Rehabilitation, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

**AIM:** In Class III anomalies, the mandible is located in a forward position in relation to the maxilla, causing a characteristic facial appearance, with major implications for aesthetics. The aetiology and expression of a Class III malocclusion must be understood before it can be clinically corrected. The aim of this study was to determine whether any correlations exist between mandibular and cranial base alterations.

**MATERIALS AND METHOD:** Lateral cephalograms of 20 skeletal Class III patients from the southern region of Romania, aged 16–24 years, and a control group of 20 skeletal Class I subjects. The Class III patients were selected according to the following criteria: Class III anomaly, age (after the pubertal growth peak), no evident genetic syndromes or other dental and/or dentomaxillary anomalies. The control group was selected according to similar criteria in terms of age and gender.

**RESULTS:** For all skeletal Class III subjects, mandibular length was significantly increased, the cranial base was affected not only in terms of dimension, but also morphology (N-S-Ba), leading to a more forward position of the mandible by modification of condylion angle (Se-S-Cd).

**CONCLUSION:** The increase of mandibular sagittal length is associated with a normal/shortened maxilla. An overall reduction was also observed in the cranial base. S-Ba distance reduction was documented in most cases, while N-S distance reduction was variable.

## SP 130

Friday, June 18 – at 10:00

### DENTAL EFFECTS OF MAXILLARY PROTRACTION WITH A FACEMASK, WITH OR WITHOUT PALATAL EXPANSION

**A Cesta**, C Berto, T Avventi, F A Miotti, Department of Orthodontics, University of Padova, Italy

**AIM:** To compare the dental effects of maxillary facemask protraction (FM) and rapid palatal expansion (RPE) followed by facemask protraction (RPE+FM), in subjects with a skeletal Class III malocclusion, compared with an untreated control group.

**SUBJECTS AND METHOD:** Forty-seven subjects divided in to three groups: group 1 (RPE+FM), 15 subjects; group 2 (FM), 18 subjects; group 3 (control), 14 subjects. Cephalometric radiographs were taken before and after treatment and, in 33 cases after a follow-up period (1.9 years). Changes between the three groups were analyzed with analysis of variance.

**RESULTS:** Cephalometric analysis showed statistically significant changes in the dental variables. The short-term evaluation showed an increase in overjet in both treated groups (group 1 +3.7 mm \*\*\*, group 2 +4 mm \*\*\*), and a significant increase in PTV-6 distance in group 1 (+4 mm \*\*\*). The other dental values examined did not show significant modifications between the three groups, although a tendency towards an increased vestibular inclination of the upper incisors (I/SN, I/Mx) was observed, with a decreased inclination of the lower incisors (i/Md). In the long-term the results obtained were maintained.

**CONCLUSION:** Maxillary protraction induced a marked improvement in the dental relationships both in the short- and long-term, although treatment with the FM mainly produced orthopaedic effects.

## SP 131

Friday, June 18 – at 10:00

### MAXILLARY PROTRACTION WITH A FACEMASK: EFFECTS ON SAGITTAL AIRWAY DIMENSIONS

**T Avventi**, A Cesta, C Berto, F A Miotti, Department of Orthodontics, University of Padova, Italy

**AIM:** A skeletal Class III malocclusion is one of the most severe maxillofacial deformities, causing many functional, phonetic, aesthetic and social problems. Maxillary protraction with a facemask is one of the most frequently used therapies in growing patients. The aim of this study was to analyze the effects of maxillary facemask protraction (FM), or rapid maxillary expansion followed by maxillary facemask protraction (RME+FM), on the sagittal airway dimensions in subjects with a skeletal Class III malocclusion, compared with an untreated control group.

**SUBJECTS AND METHOD:** Forty-seven subjects divided into three groups: group 1 (RME+FM), 15 subjects (T1-T2 period 1.5 years); group 2 (FM), 18 subjects (T1-T2 period 1.5 years); group 3 (control), 14 subjects (T1-T2 period 1.6 years). Cephalometric radiographs were taken immediately before (T1) and after (T2) treatment and, in 33 subjects, after a follow-up period of 1.9 years (T3). Linear and angular measurements were evaluated: PNS-AD1, PNS-AD2, SPS, MPS, IPS, SN/CVT. Differences between the three groups were analyzed with analysis of variance.

**RESULTS:** Favourable changes induced by maxillary protraction, with or without palatal expansion, were associated with only slight changes in the antero-posterior oro- and rhino-pharyngeal dimensions. A general increase in airway dimensions was observed in the treated subjects, though none of the changes was statistically significant (PNS-AD1 group 1, +0.73 mm, group 2, +1.46 mm, group 3, -0.5 mm; IPS group, 1 +1.2 mm, group 2, +1.6 mm, group 3 -1.87 mm; SN/CVT group 1 +6.16 degrees, group 2 +4.9 degrees, group 3 +1.41 degrees).

**CONCLUSION:** Orthopaedic treatment of a Class III malocclusion did not result in a statistically relevant increase in airway dimensions in the short or long term.

## SP 132

Friday, June 18 – at 10:00

### MAXILLARY PROTRACTION WITH OR WITHOUT PALATAL EXPANSION: CEPHALOMETRIC SAGITTAL EFFECTS

**C Berto**, A Cesta, Tavventi, F A Miotti, Department of Orthodontics, University of Padova, Italy

**AIMS:** Growing patients with a Class III skeletal pattern often exhibit a hypoplastic maxilla with a maxillary transverse deficiency. Maxillary protraction has become an established paradigm in the early management of skeletal Class III malocclusions, and rapid palatal expansion (RPE) is often used to increase the transverse width of the upper arch. The aims of this study were to analyze the sagittal effects of treatment with a facemask, with or without palatal expansion, and to compare the results with an untreated control group.

**MATERIALS AND METHOD:** Lateral cephalograms of 47 patients with a Class III malocclusion were examined before and after treatment and, in 33 cases, approximately two years post-treatment.

The sample was divided into three groups 1) 15 subjects RPE and facemask; 2) 18 subjects facemask only; 3) 14 subjects, control with untreated Class III malocclusions. Differences between the groups were determined with analysis of variance (ANOVA).

**RESULTS:** Sagittal skeletal parameters were significantly affected by facemask therapy. No significant differences were found between the expansion and non-expansion groups for any measured variable. Sagittal treatment effects were characterized by skeletal maxillary advancement and a restraining effect on mandibular growth rate. Cephalometric variables showed improvements in ANB (group 1, +3.06°<sup>\*\*\*</sup>, group 2, +3.36°<sup>\*\*\*</sup>) and Wits appraisal (group 1, +2.9 mm<sup>\*\*\*</sup>, group 2, +3.9 mm<sup>\*\*\*</sup>). Comparisons of treated and control subjects showed significant treatment effects beyond normal Class III growth. Long-term, the treated patients tended to resume a Class III growth pattern (Harvold  $\Delta$  group 1, +2.42 mm<sup>\*</sup>, group 2, +3.81 ns, group 3, +3.71<sup>\*\*</sup>), though the correction of the malocclusion was often maintained.

**CONCLUSION:** Facemask protraction produces similar changes, characterized by a maxillary forward displacement, whether associated or not with RPE.

### SP 133

Friday, June 18 – at 10:00

#### VERTICAL SKELETAL CHANGES WITH FACE MASK THERAPY, WITH OR WITHOUT PALATAL EXPANSION

**C Berto**, A Cesta, Tavventi, F A Miotti, Department of Orthodontics, University of Padova, Italy

**AIMS:** A Class III malocclusion is important in orthodontic practice, representing a difficult challenge due to its functional, aesthetic and psychological repercussions. Recent investigations have focused on three-dimensional changes due to facemask therapy, with accurate analysis of sagittal, transverse and vertical dento-facial parameters. The purposes of this research were to evaluate the vertical skeletal effects of treatment with the facemask, with or without rapid palatal expansion (RPE), and to compare these results with an untreated control group.

**SUBJECTS AND METHOD:** Forty-seven patients with a Class III malocclusion, divided into three groups 1) 15 subjects RPE and facemask; 2) 18 subjects facemask only; 3) 14 subjects, control, with untreated Class III malocclusion. Cephalometric radiographs were taken before and after treatment and, in 33 cases, about two years post-treatment. Differences between the groups were determined with the analysis of variance test.

**RESULTS:** There was a significant increase in face height but only in group 1 (Harvold Hf + 4.26 mm<sup>\*</sup>). All other vertical parameters increased, but their differences were not significant. At the long-term observation, gonial and Schwarz angles decreased in group 1 (ArGoMe -1.53°<sup>\*</sup>, Schwarz -3.7°<sup>\*\*</sup>) while face height increased in group 3 (Harvold Hf +4.46mm<sup>\*\*\*</sup>).

**CONCLUSION:** Cephalometric analysis showed an increase in intermaxillary vertical relationship in both treated groups, but statistically significant only in group 1. Further analysis of 33 patients, 21 months after the end of treatment, showed a normalization of the cephalometric vertical variables during the post-treatment period. This could be explained by the reduction in clockwise rotation, due to the tendency of both maxillary and mandibular planes to return to their previous inclination.

### SP 134

Friday, June 18 – at 10:00

#### A NOVEL METHOD OF ANALYSIS OF CLASS III PATIENTS USING THREE-DIMENSIONAL SOFT TISSUE ANALYSIS

**M Božič<sup>1</sup>**, C H Kau<sup>2</sup>, S Richmond<sup>3</sup>, M Ovsenik<sup>4</sup>, N Ihan Hren<sup>1</sup>,  
<sup>1</sup>Clinical Department of Maxillofacial and Oral Surgery and  
<sup>4</sup>Orthodontics, University Medical Centre Ljubljana, Slovenia,  
<sup>2</sup>University of Texas Health Science Centre, Houston, Texas, USA  
and <sup>3</sup>Cardiff University, Dental Health and Biological Sciences, Wales

**AIM:** To determine differences between Class III patients and a normative dataset using three-dimensional analysis.

**SUBJECTS AND METHOD:** Fourteen Class III males (MCIII) and 15 Class III females (FCIII). Forty-three males and 44 females with a Class I occlusion allowed construction of an average male (AvM) and female (AvF) face. Coordinates of three points on the facial templates of groups MCIII, FCIII and the templates AvM and AvF were compared. MCIII-AvM and FCIII-AvF superimpositions were evaluated for differences.

**RESULTS:** Vertical distances [sellion (S)-soft tissue pogonion ([P])] were statistically significantly higher for the AvM (9.1%) and MCIII (10.1%) than for AvF and FCIII ( $P < 0.05$ ). The distance of point P in the horizontal (x) axis was positive in 80 per cent for the FCIII group and in 85.7 per cent for the MCIII group. The Class III subjects differed from the average face in the lower two-thirds but in 50 (MCIII) and 60 (FCIII) per cent they also differed in the upper facial third.

**CONCLUSION:** (1) The average and Class III Slovenian male morphological face height is statistically significantly higher than the females; (2) The Slovenian Class III males and females tend towards having a left-sided chin deviation; (3) Differences between Class III patients and a normative dataset were determined.

### SP 135

Friday, June 18 – at 10:00

#### PRIMARY VERSUS PERMANENT DENTITION ANCHORAGE IN EARLY CLASS III ORTHOPAEDIC TREATMENT

**M Rosa<sup>1</sup>**, **P Lucchi<sup>2</sup>**, **E Frumusa<sup>1</sup>**, **A Caprioglio<sup>1</sup>**, Departments of Orthodontics, <sup>1</sup>University of Insubria, Varese and <sup>2</sup>University of Cagliari, Italy

**AIM:** The most appropriate timing for maxillary orthopaedic treatment in a Class III malocclusion subject is the first phase of the mixed dentition. A facemask (FM) and rapid palatal expander (RPE) are reported to be the most effective therapy for maxillary sagittal and transverse expansion. Some authors report that primary teeth can support the orthopaedic load. The purpose of this preliminary study was to evaluate the skeletal effects in two groups of young Class III young patients, treated using RPE and FM anchored on first permanent molars or on primary teeth in the mixed dentition group.

**MATERIALS AND METHOD:** Lateral cephalograms taken before (T0) and after (T1) treatment. Cephalometric analysis was based on 11 measurements. Two statistical analyses were undertaken, a paired t-test was performed to evaluate changes during T0 to T1 in each group and an unpaired t-test to study intergroup comparison.

**RESULTS:** No statistical significance was found at T0 between the two groups in distribution and treatment time. SNA, ANB and AoBo significantly increased in both groups during the treatment period, while only the primary group showed a significant improvement in facial convexity (ANPog). The mandibular plane (SN<sup>^</sup>GoGn) did not change significantly.

**CONCLUSION:** Following RPE and FM therapy in early mixed dentition, maxillary advancement could be more effective using primary teeth as anchorage. A further advantage could be the prevention of any risk of damaging already erupted permanent teeth or dentoalveolar compensation.

### SP 136

Friday, June 18 – at 10:00

#### EFFECTS OF CHIN CUP THERAPY ON SYMPHYSIS MORPHOLOGY

**H Gokalp**, **G Görgülü**, **P Demir**, Department of Orthodontics, Faculty of Dentistry, University of Ankara, Turkey

**AIM:** The chin cup has been used for many years to treat skeletal mandibular prognathism in growing patients. The pull on the orthopaedic-force chin cup is orientated along a line from the mandibular symphysis to the mandibular condyle. The purpose of this study was to evaluate the changes in symphysis morphology after chin cup therapy.

**MATERIALS AND METHOD:** Pre- and post-treatment lateral cephalometric radiographs of 36 subjects who had Class III malocclusion or anterior crossbite and seven subjects without a Class II malocclusion (control group). SNA, SNB, ANB, Na-Ba/Ptm-Gn, SN-MP, PP-MP, Ar-Go-Na, N-S-Ar, N-Ar-Go, ANS-Me/N-Me, S-Go/N-Me, symphysis height, (SY), symphysis depth (SD), SY/SD and NB/MP measurements were performed. The data obtained were statistically analyzed using a Student's t- and paired t-tests.

**RESULTS:** Pre- and post-treatment SNA, ANB, Ar-Go-Na and NB/MP measurements ( $P < 0.001$ ) were statistically significant. Pre- and post-treatment SNB, SY, SY/SD measurements were also statistically significant ( $P < 0.01$ ). Pre- and post-treatment S-Ar-Go measurements were significant ( $P < 0.05$ ). There was no statistical significance for any of the other measurements. The differences between the treatment and control group was statistically significant for all measurements ( $P < 0.001$ ).

**CONCLUSION:** Chin cup therapy has an effect on symphysis morphology; it increases symphysis height. It also affects SNA, ANB angles and lower anterior face length.

### SP 137

Friday, June 18 – at 10:00

#### CONDYLAR TRACINGS IN CLASS III MALOCCLUSION SUBJECTS BEFORE AND AFTER ORTHOGNATHIC SURGERY

M G Piancino, G Frongia, A A Bracco, C Debernardi, P Bracco, Department of Orthodontics and Gnathology, University of Turin, Italy

**AIM:** To longitudinally evaluate the condylar tracing characteristics (length, morphology, superimposition, deviations, sagittal condylar inclination and Bennet angle) recorded with computerized axiography of patients with a severe Class III malocclusion before and after orthognathic surgery and orthodontic therapy.

**SUBJECTS AND METHOD:** Eleven subjects with mandibular prognathism (6 males, 5 females, mean age  $\pm$  standard deviation  $23.10 \pm 3.13$  years) scheduled for combined orthodontic-surgical treatment. Computerized axiography was recorded with a Cadiax® Diagnostic axiograph, connected to a condylograph facebow, and Gamma Dental Software for data storage and analysis before (T0) and after (T1) therapy. Statistical analysis was performed with the chi square test to evaluate T0-T1 differences of length, morphology, superimposition, deviations, sagittal condylar inclination and Bennet angle.

**RESULTS:** There was a significant improvement at T1 of all parameters: length ( $P < 0.05$ ), normal morphology ( $P < 0.05$ ), superimposition ( $P < 0.01$ ), deviations ( $P < 0.01$ ), sagittal condylar inclination ( $P < 0.001$ ), and Bennet angle ( $P < 0.01$ ).

**CONCLUSION:** Temporomandibular joint (TMJ) border movements after surgery not only improved morphology, but also showed a significant change ( $P < 0.001$ ) in sagittal condylar inclination during protrusion (T0:  $27.5 \pm 6.9^\circ$ , T1:  $46.9 \pm 6.5^\circ$ , normal values  $52 \pm 7^\circ$ ) meaning that functional movements of the TMJ are influenced by surgery and orthodontics in severe Class III patients. Further controls and long-term evaluation are necessary to confirm the results.

### SP 138

Friday, June 18 – at 10:00

#### REVERSE-SEQUENCING CHEWING PATTERNS BEFORE AND AFTER CLASS III ORTHOGNATHIC SURGERY

M G Piancino, G Frongia, F Talpone, T Vallelonga, P Bracco, Department of Orthodontics and Gnathology, University of Turin, Italy

**AIM:** Reverse-sequencing chewing patterns are highly dyskinetic cycles characterized by asymmetric neuromuscular function. They occur with high frequency in patients with a posterior crossbite. This longitudinal study analyzed the prevalence of reverse-sequencing chewing cycles before and after orthognathic surgery and orthodontic therapy of severe Class III patients.

**SUBJECTS AND METHOD:** Eleven subjects with mandibular prognathism (6 males, 5 females, mean age  $\pm$  standard deviation  $23.10 \pm 3.13$  years) scheduled for combined orthodontic-surgical treatment. The chewing cycles were recorded with a kinesigraph (K7-I, Myotronics Inc. Tukwila, Washington, USA) and analyzed with custom-made software for data storage and subsequent analysis before (T0) and after (T1) treatment, and at a three-year follow-up (T2). Statistical analysis was performed with the chi square test to evaluate the differences between T0-T1 and T0-T2.

**RESULTS:** The percentage of reverse-sequencing chewing cycles significantly decreased with both the soft and hard boluses after therapy (T0-T1  $P < 0.01$ ) and at the 3 year follow up (T0-T2  $P < 0.05$ ). The total number of chewing cycles increased after therapy with both the soft and hard bolus (T0-T1  $P < 0.01$ ; T0-T2  $P < 0.01$ ).

**CONCLUSION:** Combined orthodontic-surgical treatment not only corrects the malocclusion, but also improves masticatory function, decreasing the reverse sequencing chewing cycles and enhancing the total number of chewing cycles. Masticatory function improvement was stable at T2.

### SP 139

Friday, June 18 – at 10:00

#### A COMPARATIVE STUDY OF CLASS III TREATMENT WITH A TONGUE APPLIANCE AND REVERSE PULL CHIN CAP

A Showkatbakhsh<sup>1</sup> A Jamilian<sup>2</sup> M Nori<sup>1</sup> F Flahiardekani<sup>3</sup> N Aghaeipour, <sup>1</sup>School of Dentistry, Shahid Beheshti Medical Science University, <sup>2</sup>School of Dentistry, Islamic Azad University and <sup>3</sup>Dental Clinic, Iran Medical Science University, Iran

**AIM:** Correction of maxillary deficiency has many treatment approaches. One of these is the tongue appliance. In this study this method was compared with a reverse pull chin cap to position the retruded maxilla in a forward position.

**SUBJECTS AND METHOD:** Two groups of growing patients. Group 1 comprised eight girls and six boys (average age 10.45 years) treated with an upper removable tongue appliance, and group 2, seven girls and seven boys (average age of 10.78 years) treated with an upper reverse pull chin cap with hook mounted in the buccal segment of the canine area to attach a rubber band from the hook of the mandibular chin cap to pull the retruded maxilla to a forward position. The records included models, radiographs and photographs taken before treatment and after correction of the skeletal malocclusion.

**RESULTS:** The following results were obtained: tongue appliance, Ar-Pog (0.001) ANS-Pog (0.05) S-Go (0.001) SNA (0.05) ANB (0.001); chin cap, Ar-Pog (NS) ANS-Pog (0.001) S-Go (0.05) SNA (NS) ANB (0.05).

**CONCLUSION:** Mandibular growth in group 1 continued to be normal and better coordination between the PFH and the PFH was observed. Growth of the retruded maxilla was improved and the increase of ANB resulted in correction of the Class III malocclusion.

### SP 140

Friday, June 18 – at 10:00

#### NATURAL HEAD POSITION IN CLASS III MALOCCLUSION SUBJECTS

Z Hedayati, Faculty of Dentistry, Shiraz University of Medical Sciences, Iran

**AIM:** Head position is very important in diagnosis based on lateral cephalometrics and soft tissue profile evaluation. It may be also different in an individual in different health situations. The aim of this study was to investigate natural head posture (NHP) in adults with a Class III malocclusion

**MATERIALS AND METHOD:** Lateral cephalograms of 45 adult Class III subjects (23 females, 22 males with a mean age of  $19.7 \pm 1.2$  years). All cephalograms were taken in NHP. The inclusion criteria were: no visual, hearing, breathing or swallowing disorders, complete permanent dentition (except for third molars), no history of orthodontic treatment or orthognathic surgery, and no

burns, injuries, or scars on the head or neck. The lateral cephalograms of 35 Class I subjects in NHP comprised the control group. Statistical analysis using paired t-tests was conducted on the obtained data.

**RESULTS:** The mean value of SN/TH was +6.5 and +11.6, and for FH/TH +0.75 and +3.5 in the Class III and Class I groups, respectively. There were significant differences between the two measurements in both groups ( $P = 0.05$ ).

**CONCLUSION:** Head posture in Class III adult subjects is rotated more downward in comparison with Class I adults.

### SP 141

Friday, June 18 – at 10:45

#### FACIAL MORPHOLOGY OF CLASS III PATIENTS IN THE MIXED DENTITION USING THREE-DIMENSIONAL IMAGING

**B Krneta**<sup>1</sup>, M Darija<sup>1</sup>, A Zhurov<sup>2</sup>, S Richmond<sup>2</sup>, M Ovsenik<sup>1</sup>,  
<sup>1</sup>Department of Orthodontics, Medical Faculty, University of Ljubljana, Slovenia and <sup>2</sup>Dental Health and Biological Sciences, Dental School, Cardiff University, Wales

**AIM:** To compare facial shape in children with Class I and Class III malocclusions in the early mixed dentition period.

**SUBJECTS AND METHOD:** Nineteen Class III subjects [9 males, 6-7 years old (mean age  $6.96 \pm 0.51$ ) and 10 females, 5-7 years old (mean age  $7.12 \pm 0.66$ )]. Thirty-six Class I subjects [20 males, 6-7 years old (mean age  $6.98 \pm 0.61$ ) and 16 females, 5-7 years old (mean age  $6.94 \pm 0.60$ )]. Three-dimensional (3D) data were acquired using a laser scanning system and associated software. Average faces were constructed for Class I and III male and female groups and superimposed to quantify differences in the facial shells. Differences in facial landmarks between the groups of children were statistically analysed using t-tests.

**RESULTS:** Absolute facial shell differences between the subgroups ranged from 2.1 to 3.00 mm. The average difference in facial shells for male Class I/III was 0.90 and for females 0.53 mm. Similarities in the facial shells between subgroups ranged from 37.40 per cent (males CI I/III) to 57.41 per cent (females CI I/III). The vertical distances N-SN and SN-P showed no statistically significant differences between the average and the Class III group.

**CONCLUSION:** Patients with a Class III facial morphology expressed more prominent mandibles when compared with the Class I facial template. Using 3D imaging technology, differences in facial morphology can be accurately quantified and visualized.

### SP 142

Friday, June 18 – at 10:45

#### ASSESSMENT OF THE PHARYNGEAL AIRWAY IN SURGICALLY CORRECTED CLASS III PATIENTS

**U Toygar Memikoğlu**<sup>1</sup>, H Aydemir<sup>1</sup>, H A Karasu<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Maxillofacial Surgery, Faculty of Dentistry, Ankara University, Turkey

**AIMS:** Surgical procedures that reposition the facial skeleton alter the soft tissues that are attached to the bone in order to effect facial changes. One aspect of this surgery is the effect of skeletal movements on posterior airway space. Most pharyngeal airway studies have been evaluated in mandibular set back procedures. The purpose of this research was to determine the effect of orthognathic surgery on pharyngeal airway in Class III patients, to compare the results of different surgical techniques, and to determine the change at the position of hyoid bone.

**SUBJECTS AND METHOD:** Forty-eight Class III adult patients, nine of whom were treated with maxillary advancement (MA), seven with mandibular set-back (MS), and 32 with bimaxillary (BM) surgery. Cephalometric records were taken before treatment, after surgery and about 1 year after surgery (end of treatment). The pharyngeal airway was assessed using planimetric and linear measurements. For statistical analysis, a repeated analysis of variance Bonferroni test were used.

**RESULTS:** No differences were found at the position of the hyoid bone. The nasopharyngeal area was significantly increased in all

groups. Oropharyngeal area, SPSS and IPS parameters were significantly decreased after MS. In the BM and MA groups, PPS was significantly increased, IPS was significantly decreased, and differences were detected in the oropharyngeal and hypopharyngeal areas. Maximum contraction was seen in the MS group.

**CONCLUSION:** Different surgical procedures result in different effects on pharyngeal airway space.

### SP 143

Friday, June 18 – at 10:45

#### THE PREDICTION OF FUTURE ORTHOGNATHIC SURGERY NEED IN CHILDREN

**G Schuster**, M Linke, Department of Orthodontics, University of Frankfurt and Private Office, Germany

**AIM:** In a previous multicentre study a total 88 patients with a Class III malocclusion were grouped into orthopaedic/orthodontic ( $n = 65$ ) and surgery patients ( $n = 23$ ) according to their records after puberty. Discriminant analysis (DA) and logistic regression (LogR) were applied to 20 landmarks of the patients cephalograms before puberty to identify dentoskeletal variables, which provided the best group separation, respectively best predictability of group membership. Both models were highly significant ( $P < 0.001$ ), classifying over 93 per cent of the patients correctly. The extracted variables were identical for both procedures Wits appraisal, palatal plane angle (NLNSL) and individualized inclination of the lower incisors (M1MLind). Equation for DA  $2.277 \pm 0.492$  Wits  $+0.116$  NLNSL  $-0.056$  M1MLind; score =  $-0.653$ . Equation for LogR  $-7.968 - 1.323$  Wits  $- 0.363$  NLNSL  $+ 0.153$  M1MLind; score =  $0.5$ .

To prove the formulas and their predictive value cephalograms of 94 newly recruited patients with a Class III from different private orthodontic offices were evaluated according to the same criteria as above. Sixty-two were non-surgery, 32 surgery cases.

**RESULTS:** The new group showed a significantly worse overall classification (65% compared with the formula for the DA, and 84% for LogR). In DA analysis the non-surgery cases were classified correct in 67 per cent correct whereas for LogR correction classification was found for 89 per cent (former study 95% for DA, 98% for LogR). Surgery cases reached 63 per cent for DA and 75 per cent for LogR (former study 87% DA, 82% LogR). The reasons for the reduction in the predictive value might be due to the independent new patient group, not following the inclusion criteria as strictly and different treatment procedures in the various private offices.

### SP 144

Friday, June 18 – at 10:45

#### HYPERDIVERGENT CLASS III AND MAXILLARY HYPOPLASIA: IS FACEMASK THERAPY NECESSARY?

**M Rosa**<sup>1</sup>, **P Lucchi**<sup>2</sup>, Departments of Orthodontics, <sup>1</sup>University of Insubria, Varese and <sup>2</sup>University of Cagliari, Italy

**AIM:** Approximately one-third of Class III malocclusions involve a hypoplastic maxilla, often combined with a hyperdivergent growth pattern. The purpose of this study was to compare the skeletal effects in two groups of young Class III hyperdivergent patients treated using a rapid palatal expander (RPE) and a Delaire facemask, or only RPE.

**MATERIALS AND METHOD:** Each appliance was anchored on the second primary molars and two lateral cephalograms were taken before (T0) and after (T1) the selected treatments options. Twelve cephalometric measurements were made at T0 and T1 and two statistical analyses were performed. A paired t-test to evaluate changes during T0 and T1 in each group and an unpaired t-test to study intergroup comparison.

**RESULTS:** There was no statistical difference between the two groups at T0. At T1, SNA, ANB, AoBo and NAPog showed a different improvement in group A compared with group B

**CONCLUSION:** Both treatment options are effective. Clinically the main difference in choosing RPE and a Delaire facemask or only RPE, is the evaluation of facial convexity.

**SP 145**

Friday, June 18 – at 10:45

**COMPARISON OF CRANIAL BASE ANGLE BETWEEN CLASS II AND CLASS III MALOCCLUSION SUBJECTS****Z Hedayati**, Faculty of Dentistry, Shiraz University of Medical Sciences, Iran**AIM:** To retrospectively determine differences in cranial base flexion that can differentiate Class II and Class III malocclusion subjects.**MATERIALS AND METHOD:** Fifty-six cephalometric radiographs (26 Class II, 26 Class III). The age range of the subjects was from 8 to 18 years. The cephalograms were traced and digitally analyzed using Dolphin 9 software. The measured angles were SNA, SNB, N-S-Ar. The linear measurements consisted of the anterior and posterior cranial bases, Co-ANS and Co-Pog. The data were statically analyzed by paired *t*-test.**RESULTS:** Cranial base angle was smaller and cranial base length shorter in the Class III group ( $P = 0.05$ ). There was a significant difference in maxillary and mandibular length between the two groups. A longer maxillary length was observed in the Class II group and a longer mandible was found in the Class III patients.**CONCLUSION:** As well as the length of jaw bases, the cranial base angle may be used as an indicator of Class II and III malocclusions.**SP 146**

Friday, June 18 – at 10:45

**THE CRANIAL BASE AND CRANIOFACIAL COMPLEX IN CLASS III MALOCCLUSION: A CEPHALOMETRIC STUDY\*\*\*****J Blaga**, Department of Orthodontics, University of Oradea, Romania**AIMS:** To evaluate the cephalometric characteristics of the cranial base, the middle cranial fossa and the nasomaxillary complex, to identify the proportions of the soft tissue profile, and to establish differences in the parameters of the craniofacial complex in patients with Class III malocclusions.**MATERIALS AND METHOD:** Lateral pre-treatment cephalograms of 100 patients (40 females, 60 males) aged between 6 and 30 years diagnosed with a Class III malocclusion. The patients presented mandibular prognathism or maxillary retrusion, or a retrusive maxilla with a protrusive mandible and an open bite facial type. Jarabak, Ricketts, Tweed and Downs methods were used. The data were statistically analysed.**RESULTS:** Significant differences between males and females were found for anterior cranial base angle. Statistically, a growth deficiency at the mid cranial fossa was evident.**CONCLUSION:** A posterior position of the maxilla in Class III patients is not only caused by a short maxilla, but can significantly be influenced by a growth deficiency in the mid cranial fossa. The mandibular prognathism was related to a small saddle angle. Different cranial growth base patterns influence the characteristics of the craniofacial complex.**SP 147**

Friday, June 18 – at 10:45

**PREVALENCE OF DIFFERENT CLASS III MALOCCLUSIONS AND THE TIMING OF TREATMENT****A Toth Pirnovar**<sup>1</sup>, M Ovsenik<sup>2</sup>, F Farčnik<sup>3</sup>, <sup>1</sup>Community Health Centre, <sup>2</sup>Department of Orthodontics, University of Ljubljana and <sup>3</sup>Orthos, Ljubljana, Slovenia**AIM:** To determine the prevalence of different types of Class III malocclusions in Slovenia in the primary and mixed dentitions.**SUBJECTS AND METHOD:** Seventy-two patients with Class III malocclusions referred between 2004–2005 were examined at the beginning of treatment. For each patient an anamnesis, a clinical examination, facial photographs, study casts and radiographs (panoramic and lateral cephalographs) were obtained and analyzed.**RESULTS:** Most of the Class III malocclusions (74%) were a combination of dentoalveolar and skeletal; in only 4 per cent was the malocclusion dentoalveolar. Twenty-nine per cent of the subjects had a retrognathic maxilla and orthognathic mandible. In 32 per cent treatment was started before 10 years of age, in 54 per cent of patients between 10 and 18 years, and in 14 per cent after 19 years. The patients in the last group needed a combination of orthodontic treatment and jaw surgery to correct the malocclusion.**CONCLUSION:** Most of the patients with Class III malocclusions had a combination of a dentoalveolar and skeletal Class III malocclusion. The majority were not directed on time for orthodontic treatment and therefore treatment planning was more demanding and expensive, treatment time was longer, and the stability of achieved result could be questionable.**SP 148**

Friday, June 18 – at 10:45

**LONG-TERM STABILITY OF SKELETAL CLASS III PATIENTS TREATED WITH SPLINTS AND CLASS III ELASTICS****F Ferro, D Apicella, V Assumma, S Nucci, L Perillo**, Department of Orthodontics, Second University of Naples, Italy**AIM:** To evaluate stability after facial growth and treatment with splints, Class III elastics, and chin cup (SEC III) and to investigate the main determinants of relapse.**MATERIALS AND METHOD:** Data from pre-, post-treatment, and long-term (at least 3 years after retention) serial cephalograms of 52 patients who received SEC III treatment were studied only for those whose growth had ceased (females over 18 years, males over 21 years).**RESULTS:** At the end of the follow-up period (an average of 9 years), clinical relapse was observed in six subjects (overjet  $\leq 0$ ). A low Wits appraisal, ANB angle, and overbite, and high SNB angle were the best predictors of relapse at the end of treatment. Significantly greater decreases of Wits appraisal and increases of ramus length during follow-up were further associated with relapse.**CONCLUSION:** SEC III is a reliable treatment for skeletal Class III malocclusions. Long-term stability can be enhanced by a deeper overbite and the best possible skeletal correction. The results suggest that the correction occurs with no mandibular rotation but might be better with forward rotation.**SP 149**

Friday, June 18 – at 10:45

**LINEAR VARIABLES AND THE PRESENCE OF THIRD MOLARS IN CLASS III AND CLASS II DIVISION 2 PATIENTS****B Mady Maričić**<sup>1</sup>, M Legović<sup>1</sup>, R Ružman Knežević<sup>2</sup>, S Špalj<sup>1</sup>, <sup>1</sup>Department of Paediatric Dentistry and Orthodontics, School of Medicine, University of Rijeka and <sup>2</sup>Dental Polichlinic Zagreb, Croatia**AIM:** To determine the presence of third molar germs in patients with Class II division 2 and Class III malocclusions regarding linear craniofacial characteristics.**SUBJECTS AND METHOD:** One hundred and forty six children [Class II division 2 ( $n = 77$ ), Class III ( $n = 69$ )]. Assessments were made on panoramic radiographs and lateral cephalograms. Two types of measurement of the linear cephalometric variables were undertaken on the lateral cephalograms: the length of the maxilla (CoA) and the length of the mandible (CoGn). The Pearson chi squared, Fisher's exact test and non-linear regression equation was used to determine differences that were statistically significant.**RESULTS:** Third molar germs were present significantly more often in the upper jaw in Class II division 2 subjects (58 versus 44%) and in the lower jaw in Class III subjects (83 versus 69%). A correlation between the presence of upper third molar germs was found for CoA, and a correlation between the presence of lower third molar germs for CoGn.

**CONCLUSION:** There is a correlation between the presence of third molar germs and sagittal maxillomandibular relationship and jaw length of the jaw. Investigations of the differences in calcifications of all permanent teeth in such malocclusion subjects are necessary.

**SP 150**

*Friday, June 18 – at 10:45*

**PREVALENCE OF CLASS III MALOCCLUSIONS – IS THERE SUFFICIENT EVIDENCE?**

**M Orellana**, D Hary, University of California, San Francisco, USA

**AIM:** An Angle Class III malocclusion, commonly characterized by a negative anterior overjet, can be handicapping both functionally and socially. There is ambiguous information regarding its prevalence; most studies report Class III malocclusions as the smallest proportion of malocclusion types in any given population. Nonetheless, Class III malocclusion patients present some of the greatest need for orthodontic treatment. Multiple epidemiological studies have documented the prevalence of Class III malocclusions in many different populations; however, and in spite of the continually increasing data on Class III malocclusion prevalence, little has been done to consolidate this information in a comprehensive and critical way. Thus, this study seeks to review the relevant literature on Class III malocclusion prevalence.

**MATERIALS AND METHOD:** The electronic databases, PubMed and Embase, were searched using specific inclusion criteria to obtain applicable articles; all pertinent references were also examined for acceptability.

**RESULTS:** A total of 21 articles were identified using the inclusion criteria. The prevalence of reported malocclusion ranged from 0 to 26.7 per cent in different populations examined. A mean prevalence of 7.8 per cent was calculated across all populations. The highest prevalence of 15.6 per cent was found among individuals of Eastern Asian descent, while Indian populations had the lowest rate at 1.1 per cent.

**CONCLUSION:** The prevalence of Class III malocclusions varies greatly among racial and ethnic groups. However, these findings should be interpreted with caution, as the reported prevalence data depends greatly on several factors, including sampling, population age range, and methodology for recording Class III malocclusions.



## SCIENTIFIC POSTERS

### Free Topics

#### SP 151

Friday, June 18 – at 10:45

##### PREVALENCE OF FUNCTIONAL DISORDERS IN PATIENTS WITH CLASS III MALOCCLUSIONS

**M Mikac Cankar**<sup>1</sup>, M Ovsenik<sup>2</sup>, <sup>1</sup>Community Health Centre and <sup>2</sup>Department of Orthodontics, Faculty of Medicine, University of Ljubljana, Slovenia

**AIM:** Causal factors for the development of a Class III malocclusion include heredity and functional disorders of the orofacial system. The aim of this study was to determine the prevalence of the latter in patients with a Class III malocclusion.

**SUBJECTS AND METHOD:** Seventy-two patients with a Class III malocclusion were evaluated for the presence of an incorrect breathing mode, swallowing and chewing, common deleterious habits and speech impairment with the use of functional testing, history and speech assessment. The prevalence of functional disorders was also determined from analysis of sagittal values on lateral cephalometric radiographs.

**RESULTS:** Functional abnormalities of the orofacial system were identified in all 72 patients. The prevalence did not decrease with age. Some were more common in patients requiring combined orthodontic and surgical treatment than in the group as a whole. The prevalence of functional irregularities was related to the position of the jaws in the sagittal plane.

**CONCLUSION:** There is a close association and interdependence of morphological and functional disorders of the orofacial system.

#### SP 152

Wednesday, June 16 – at 10:00

##### LONG-TERM OUTCOME AND STABILITY IN TREATED CLASS I EXTRACTION AND NON-EXTRACTION PATIENTS

**M R Gjethammer**, M Midtbø, O E Bøe, Department of Clinical Dentistry, Division of Orthodontics, University of Bergen, Norway

**AIM:** When evaluating successful treatment result, both the effects of growth and the outcome of orthodontic treatment should be assessed. The purpose of this longitudinal cephalometric study was to evaluate changes in hard and soft tissue variables after treatment, post-retention and 10 years post-retention, and to compare extraction and non-extraction patients.

**MATERIALS AND METHOD:** Cephalograms of 37 patients, (extraction and non-extraction) with Angle Class I malocclusions were digitized and traced according to hard and soft tissue landmarks.

**RESULTS:** No significant differences in cephalometric values were identified between the extraction and non-extraction group. Significant changes over time, from were found for SNB, ANB, SNPog, Gonial angle, ML/NSL, ML/NL, interincisal angle, Ili/NB, Holdaway ratio, Ls/EL and Li/EL.

**CONCLUSION:** Soft tissue variables seemed to change during all of the intervals in the study, while the hard tissue measurements changed mostly during treatment and after the retention phase.

#### SP 153

Wednesday, June 16 – at 10:45

##### THE QUERCETIN EFFECT ON SOX9 AND COLLAGEN II EXPRESSIONS IN SPHENO-OCCIPITAL SYNCHONDROSES

A Veerawattanagul, R Wong, B Rabie, The University of Hong Kong, Prince Philip Dental Hospital, 34 Hospital Road, Hong Kong SAR, China

**BACKGROUND:** Quercetin is a flavonoid phytoestrogen that stimulates osteoblasts. Spheno-occipital synchondrosis is an important growth centre of the craniofacial skeleton and its growth affects positions of both maxilla and mandible, sagittally and vertically. It is important to identify chemicals that affect its growth.

**OBJECTIVE:** To investigate the effect of quercetin on the growth of spheno-occipital synchondrosis by measuring the expressions of Sox9 and type II collagen in spheno-occipital synchondrosis, using an in vitro mouse model.

**MATERIAL AND METHODS:** A total of 50 spheno-occipital synchondroses and surrounding tissue dissected from 1-day-old male BALB/c mice were randomly assigned to control and experimental groups. Each group was subdivided into 5 different time points (6, 24, 48, 72 and 168 hours), each subgroup contained 5 synchondroses (n=5). In the experimental group, the spheno-occipital synchondroses were immersed in the BGJB medium and 1 &#61549;M quercetin solution. In the control group, the spheno-occipital synchondroses were immersed in the BGJB medium. Tissue sections were subjected to immunohistochemical staining for quantitatively analysis of Sox9 and type II collagen expression, using image analyzer.

**RESULTS:** There was a statistical significant increase of 32.3% (P<0.001) in the expression of Sox9 in the experimental group compared to the control group at 24 hours. Furthermore, there was a statistical significant increase of 23.0% (P<0.001) in the expression of type II collagen in the experimental group compared to the control group at 72 hours.

**CONCLUSION:** Quercetin increased the growth response of spheno-occipital synchondrosis through increasing expressions of Sox9 and type II collagen.

#### SP 154

Wednesday, June 16 – at 10:45

##### EFFECT OF QUERCETIN ON EXPRESSION OF BMP2 AND PTHRP IN MICE SPHENO-OCCIPITAL SYNCHONDROSES

K Wangwarunyo, R Wong, B Rabie, The University of Hong Kong, China

**BACKGROUND:** Spheno-occipital synchondrosis is an important growth center that affects the position of maxilla and mandible. Due to its inaccessible position, modification of its growth by mechanical means using appliance is not possible. To explore the biological possibility of its growth modification with chemicals could be an important breakthrough. Quercetin, one of the flavonoids found in onions, apples and grapes, has been reported to affect new bone formation locally. It may be used for growth modification.

**OBJECTIVE:** To evaluate the expressions of BMP2 and PTHrP during growth in spheno-occipital synchondroses in response to quercetin, using a mouse model in vitro.

**MATERIALS AND METHODS:** Spheno-occipital synchondroses together with adjacent structures were aseptically excised from fifty 1-day-old male BALB/c mice that were randomly assigned to 5 control and 5 experimental groups in 5 experimental periods. In the control groups, the specimens were immersed in BGJb medium. In the experimental groups, 1 µM quercetin was added into BGJb medium. Both groups were then cultured for 6, 24, 48, 72 h and 7 days. Alcian blue-PAS staining was used to study growth of the synchondroses; immunohistochemical staining to identify BMP2 and PTHrP expressions. The areas of BMP2 and PTHrP expressions were measured.

**RESULTS:** Quantitative analysis showed that both BMP2 and PTHrP expressions increased significantly at hour 48 of quercetin application in the experimental group ( $p < 0.001$ ). Compared with the control groups, both BMP2 and PTHrP were expressed consistently higher in the experimental groups at all time points.

**CONCLUSIONS:** Quercetin enhances an increase in BMP2 and PTHrP expressions in spheno-occipital synchondroses. It is a new discovery that the growth of spheno-occipital synchondrosis can be modified by chemical mean.

### SP 155

Wednesday, June 16 – at 10:45

#### OCCLUSAL FEATURES OF CHILDREN WITH ADENOID HYPERTROPHY IN COMPARISON WITH A CONTROL GROUP

**K Lubinska**, M Syrynska, K Wozniak, M Durka-Zajac, Department of Orthodontics, Pomeranian Medical University, Szczecin, Poland

**AIM:** There are various reports concerning the possible relationship between adenoid hypertrophy and occlusion. Adenoid hypertrophy is the most common cause of upper airway obstruction in children. The consequence of this disorder is mouth breathing. The purpose of this study was to investigate the association between overbite, overjet, distal occlusion, crossbite and adenoid hypertrophy.

**SUBJECTS AND METHOD:** Fifty-six Polish children aged between 2 and 13 years (medium age 5.95 years) with adenoid hypertrophy identified for surgical intervention were compared with a control group matched for age, gender and number. None of the children in the control group had adenoid hypertrophy or had undergone adenoidectomy or orthodontic treatment. Evaluation of anamnestic history, clinical examination, and dental casts measurements were undertaken and the group means were compared. Statistic analysis was performed.

**RESULTS:** Children with enlarged adenoids had a larger overjet ( $P = 0.00104$ ) and a deeper overbite ( $P = 0.03664$ ) compared with the control group. A crossbite occurred significantly more often in the study group than in the control group ( $P = 0.02346$ ). Children with adenoid hypertrophy were 4 times more likely to develop distal occlusion [odds ratio (OR) 4.18,  $P = 0.003$ ], and 3 times more likely to develop a crossbite (OR 3.37,  $P = 0.046$ ) than children in the control group.

**CONCLUSION:** The occlusal features of children with adenoid hypertrophy differed from those of the control group, suggesting that adenoid hypertrophy influences occlusion. This influence is probably not direct but caused by mouth breathing.

### SP 156

Wednesday, June 16 – at 10:45

#### ADVANCED MORPHOMETRICS: REGISTRATION OF THREE-DIMENSIONAL FACIAL IMAGES

**A Zhurov**, A Toma, S Richmond, School of Dentistry, Cardiff University, Wales

**AIM:** Appropriate registration (superimposition) of three-dimensional (3D) facial images is the most important prerequisite for evaluating and comparing facial variations. Traditional registration techniques, based on a small number of landmarks, are not accurate enough for the assessment of facial growth in children, facial asymmetries, and other aspects. The aim of this research was

to develop an advanced registration techniques that take into account all data points available.

**SUBJECTS AND METHOD:** Two Konica Minolta VIVID 900/910 cameras were used for data acquisition. Rapidform® software and in-house developed subroutines were employed for data processing. A sample of 350 scanned faces of 15-year-old British Caucasians was examined and 21 facial landmarks were placed on each image. Generalised Procrustes analysis was used to register the sets of landmarks.

**RESULTS:** Mid-endocanthion was found to be the least variable point in the eye region, which is known to undergo the least variations as the face grows. A method for identifying a common reference frame for each face was developed as a set of subroutines for Rapidform®. The reference frame has mid-endocanthion as its origin and involves three reference planes: (i) mid-sagittal as the symmetry plane of the symmetrised structure consisting of maxilla and its mirror counterpart, (ii) transverse as the plane normal to the axis of the cylinder fitting the symmetrised structure and (iii) coronal as the plane normal to the first two.

**CONCLUSIONS:** The proposed common reference frame is automatically identified for each landmarked face and can be used to superimpose 3D facial images. This approach is based on all facial data available, is more accurate than the traditional approach based on landmarks only, and is especially suitable for studying facial growth, evaluating facial asymmetries and constructing average faces.

### SP 157

Wednesday, June 16 – at 11:30

#### RADIOGRAPHIC STUDY OF SUPERNUMERARY TEETH CAUSING NON-ERUPTION OF PERMANENT INCISORS

**M T Chew**, National Dental Centre Singapore

**AIM:** To investigate the radiographic characteristics of anterior maxillary supernumerary teeth associated with impacted maxillary permanent central incisors in an Asian population.

**SUBJECTS AND METHOD:** Sixty-five patients aged between 7 to 16 years (mean = 9.1 years, s.d. = 2.2 years) who presented with impacted permanent central incisors associated with the presence of supernumerary teeth. Radiographs of these patients were evaluated to investigate the features of the supernumerary teeth.

**RESULTS:** The majority of the cases ( $n = 63$ ) involved unilaterally impacted permanent central incisors (28 left, 35 right) while only two involved both permanent central incisors. There were more males than females, a ratio of 2:1. The supernumerary teeth in 31 cases (48%) were odontomes, 18 (28%) were conical shaped, while the remaining 16 (24%) were tuberculate shaped. All but two of the supernumerary teeth were unerupted with the majority positioned palatal to the impacted permanent central incisors (60%) while the rest were either occlusal to the impacted incisors (32%) or at the midline (18%).

**CONCLUSION:** Males in this study were twice as likely to present with unerupted permanent central incisors associated with submerged supernumerary teeth. Odontomes were found to be most commonly associated with unerupted permanent central incisors.

### SP 158

Wednesday, June 16 – at 11:30

#### ACCURACY OF MECHANICAL TORQUE LIMITING GAUGES FOR MINI-SCREW PLACEMENT

**M Schätzle**<sup>1,2</sup>, D Golland<sup>1</sup>, M Roos<sup>3</sup>, B Stawarczyk<sup>4</sup>, <sup>1</sup>Orthodontics and Pediatric Dentistry, <sup>2</sup>Institute of Social and Preventive Medicine and <sup>4</sup>Clinic for Fixed and Removable Prosthodontics and Dental Material Science, University of Zürich, Switzerland and <sup>2</sup>Orthodontics, Prince Philip Dental Hospital, Hong Kong University, SAR China

**AIM:** To determine and compare the accuracy of four available mechanical torque limiting gauges (MTLGs) for mini-screw placement.

**MATERIALS AND METHOD:** The torque output of six randomly obtained MTLGs, either of the screwdriver or torque ratchet type of four miniscrew manufacturers (AbsoAnchor, Spider Screw, Tomas pin and Lomas orthodontic mini Anchor System). Mounted on a joint, a universal testing machine applied a perpendicular force to a lever arm with a crosshead speed 1.0 mm/minute. For each device, 10 repetitions of the corresponding target torque level were recorded after initial sterilization and after 5, 10, 20, 50 and 100 times to evaluate their potential influence on MTLGs. The breakpoints (Ncm) were calculated for comparison of the groups. Descriptive statistics and mean breakpoints values for each MTLG were computed and compared with the reference values indicated on the respective torque gauges provided by the manufacturer.

**RESULTS:** The mean torque values for the AbsoAnchor MTLG device was significantly below torque levels, but provided consistent torque values. At higher torque levels, all but one obtained values for the Spider Screw. The ratchet type of MTLGs (Tomas pin, Lomas orthodontic mini Anchor System) yielded statistically significant different mean breakpoints than the indicated limits. Each individual MTLG produced, independently, constant breakpoint torque values, but differed significantly from each other. For all but the Spider Screw MTLG, the sterilisation process had a statistically significant different influence at the various breakpoint torque levels.

**CONCLUSION:** Compared with the manufacturers' preset torque levels, significant variations were observed between individual devices. The torque output of each individual device deviated in varying degrees from target torque values and was influenced to various degrees by the sterilization process over time.

### SP 159

Wednesday, June 16 – at 11:30

#### CHANGES IN SKULL PARAMETERS IN CLASS II DIVISION 2 SUBJECTS

**M Dushenkova**, A Anikiyenko, Moscow State Medical Dentistry University, Russia

**AIM:** To study age-related change intensity in the cerebral and facial parts of skull of Angle Class II division 2 children to detect the skull morphological structures differing from their normal growth intensity during the primary dentition period.

**MATERIALS AND METHOD:** Sixty-two teleroentgenograms of the head in lateral projection. Thirteen longitudinal and 14 vertical parameters of the facial and cerebral parts of skull of 30 children (7-12 years of age) during the primary dentition period and of 32 children (12-15 years of age) were measured.

**RESULTS:** The intensity of age-related changes of many longitudinal and vertical parameters of the skull significantly differed from normal.

**CONCLUSION:** Children with an Angle Class II division 2 malocclusion should be treated as early as possible.

### SP 160

Wednesday, June 16 – at 11:30

#### CORRELATION BETWEEN ANTEGONIAL NOTCH DEPTH WITH SAGITTAL AND VERTICAL PARAMETERS

**H R Fattahi**, E Yosefipoor, S Zandi, A A Karamifar, Faculty of Dentistry, Shiraz University of Medical Sciences, Iran

**AIM:** Facial growth, especially of the mandible, is an important factor in diagnosis and treatment planning. It has been suggested that the depth of the mandibular antegonial notch can be used as a predictor of facial growth pattern. The purpose of this study was to investigate the correlation between antegonial notch depth with sagittal and vertical indices.

**MATERIALS AND METHOD:** This cross-sectional study was carried out on 150 cephalometric radiographs of 72 males and 78 females aged 13 to 26 years (mean 19 years) selected using the non-probability sampling method. The vertical (Y-axis, SN/MP angle, Björk angle and Jarabak index) and sagittal (ANB and Wits) parameters in three subgroups of antegonial notch (shallow, me-

dium and deep) were measured. Finally the correlation was determined between the antegonial notch with these parameters using ANOVA and Pearson's tests.

**RESULTS:** There was a high correlation between vertical parameters and antegonial notch depth. The mean of the antegonial notch was 2.4 mm in normal growth patterns and 1.8 and 3.9 mm in horizontal and vertical growth patterns, respectively. Although the Student-Neumann-Keuls test showed significant differences between the groups ( $P = 0.001$ ), the antegonial notch in the sagittal groups was different only between Class III to Class I and Class II malocclusions. Pearson correlation test showed a significant relationship between antegonial notch depth and all parameters, except Y-axis in males and ANB in females.

**CONCLUSION:** There is a close relationship between antegonial notch depth and vertical determining parameters.

### SP 161

Wednesday, June 16 – at 11:30

#### EXTRAORAL JUDGEMENT OF SKELETAL CLASS

**M Miguez Contreras**<sup>1</sup>, I Nieto Sanchez<sup>2</sup>, R Ruano Gomez<sup>1</sup>, F Rodriguez Arnal<sup>1</sup>, M Romero Maroto<sup>1</sup>, <sup>1</sup>Stomatology, Anatomy and Embryology Department, Universidad Rey Juan Carlos, Madrid and <sup>2</sup>Private Practice, Spain

**AIM:** To determine whether it is possible to distinguish the cephalometric skeletal Class from a digital lateral photograph.

**MATERIALS AND METHOD:** One hundred lateral pre-treatment photographs of orthodontic patients (which had cephalograms assessed with Nemoceph® software) were shown to two groups of participants (7 males, 21 females): nine orthodontists and 19 Master of Orthodontic students. All the right profile photographs were taken by the same photographer under the same light conditions using a digital camera (Reflex Olympus E330). The participants were asked to indicate on a questionnaire if the patient was a skeletal Class I, II or III according to Ricketts, with no limit on the time allowed. Data were analysed using one-way repeated measures ANOVA with Bonferroni *post hoc* adjustment for multiple testing and an unpaired *t*-test for gender using the program, Prism 3.0.  $P < 0.05$  results were considered statistically significant.

**RESULTS:** Only 47.96 per cent of the participants correctly identified the skeletal Class of the patients. A Class III was the most difficult to identify (36.76%) versus Class I (39.46%). A Class II malocclusion was correctly identified most often (60.71%). Differences were significant only between Class II and the other two Classes ( $P < 0.001$ ) but not between Class I and Class III ( $P > 0.05$ ). There were no differences between genders ( $P > 0.05$ ).

**CONCLUSION:** Class II patients are easier to distinguish from a lateral photograph than Class III or Class I. Lateral photographs do not allow correct identification of the skeletal Class – cephalometry is also required.

### SP 162

Wednesday, June 16 – at 11:30

#### LONG-TERM SKELETAL AND SOFT TISSUE OUTCOMES OF BIONATOR THERAPY IN CLASS III MALOCCLUSION SUBJECTS

**V Giuntini**<sup>1</sup>, K Faltin<sup>2</sup>, L Franchi<sup>1</sup>, <sup>1</sup>Università degli Studi di Firenze, Italy and <sup>2</sup>University Paulista, São Paulo, Brazil

**AIM:** The literature lacks long-term controlled data concerning the outcomes of functional jaw orthopaedics in Class II patients. The aim of this research was to evaluate the long-term skeletal and soft tissue changes induced by the bionator in Class II subjects at the completion of active growth.

**SUBJECTS AND METHOD:** Twenty Class II patients (6 males, 14 females) treated consecutively with the bionator. The sample was evaluated at T1, start of treatment; T2, end of Bionator therapy; and T3, long-term observation (including fixed appliances). Their mean age at T1 was approximately 10 years (T1), at T2 approximately 12 years and at T3 approximately 19 years (CS 6). The control group consisted of 20 subjects (8 males, 12 females) with untreated Class II malocclusions. Lateral cephalograms were ana-

lyzed at the three time points for all groups. Student's t-tests were used for comparisons of starting skeletal facial forms, and of the T1-T2 and T1-T3 changes between the groups.

**RESULTS:** The bionator group showed significant favourable T1-T2 changes both at the skeletal and dentoalveolar levels. The vertical dimension was increased. Significant modifications were also found for the soft tissues. The treated group showed a final improvement in soft tissue Pogonion of about 2.5 mm. Significant mandibular changes were noted in the treated group, with a net average 3.3 mm long-term increase in mandibular length in comparison with the untreated Class II controls.

**CONCLUSION:** Bionator treatment of Class II malocclusions maintains favourable results over the long-term with a combination of skeletal, dentoalveolar, and soft tissue changes.

## SP 163

Wednesday, June 16 – at 11:30

### A SYSTEMATIC REVIEW OF INTERCEPTIVE TREATMENT OF PALATALLY DISPLACED MAXILLARY CANINES

**J Naoumova**, J Kuroi, H Kjellberg, Department of Orthodontics, Institute of Odontology at the Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden

**AIM:** To assess whether interceptive treatment in the mixed dentition prevents impaction of palatally displaced maxillary canines (PDC) by systematically reviewing the literature.

**MATERIALS AND METHOD:** A literature search of PubMed, Cochrane Library electronic databases and Scopus, was performed covering the period from January 1966 to May 2009. The inclusion criteria were mixed dentition with uni- or bilateral PDC, randomized controlled trials (RCT), prospective and retrospective studies with untreated controls, and clinical trials comparing at least two treatment strategies. Three reviewers selected and extracted the data independently and evaluated the quality of the studies. Inter-examiner reliability was measured using Intraclass Correlation Coefficient (ICC).

**RESULTS:** The search strategy resulted in 686 articles, of which two met the inclusion criteria. Because of the unequivocal results and heterogeneity in the study methods, the scientific evidence was too weak to fully evaluate the effect that interceptive treatment might have on PDC and which treatment modalities were most effective. The quality of the studies was rated as low because of inadequate sample selection and deficient description of sample size, confounding factors, uncertainty of randomization and no blinding in measurements. The ICC value for total scores was >0.80, e.g. perfect agreement.

**CONCLUSION:** To obtain reliable scientific evidence as to whether interceptive treatment prevents impaction of PDC and which treatment modalities are the most effective, better controlled and well-designed RCTs are needed. Future studies should also include assessment of patient satisfaction and pain experience as well as analysis of cost and side-effects of the treatments.

## SP 164

Wednesday, June 16 – at 11:30

### VEGETABLE ALTERNATIVES TO FIGHT AGAINST DENTAL MICROBIALS

**I Ionescu**, M Burlibasa, G Tanase, A Vereanu, D Mocuta, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

**AIM:** Plant extracts used as anti-infectious ecological alternatives to conventional treatments have many advantages, the most important being the very low risk of side-effects. Many antiseptic substances similar to antibiotics (usnic acid), which curb or destroy the development of microorganisms, have been found. The aim of this research was to investigate *in vitro* the influence of usnic acid on microbial development in orthodontic devices.

**MATERIALS AND METHOD:** Twenty dental plates, obtained from 20 different subjects obtained under sterile conditions were placed in physiological sterile water for 1-15 minutes with a suspension of usnic acid at a concentration level of 200 µg/ml. A sam-

ple for each schema was taken and put placed a brain heart infusion environment in order to set the increase curb by determining the total number of germs developed at different time intervals since incubation (3-24 hours). Determination of the total number of microorganisms was achieved using the method of decimal microdilution and spot sampling on blood gelose.

**RESULTS:** Usnic acid selectively curbed the development of biofilms formed by gram-positive bacteria and haemolytic expression of the microorganisms isolated on the dental plates. The rate of multiplication of the isolated microorganisms on the dental plate was influenced by the preceding contact with usnic acid; their increase curb being modified. These effects demonstrate the interference of usnic acid with the mechanisms of intra and inter-signalling based 'quorum sensing and response'.

**CONCLUSION:** Use of usnic acid prevents dental plate microorganism formation and is a new ecological anti-infective alternative to conventional antibiotherapy.

## SP 165

Wednesday, June 16 – at 11:30

### TESTING THE FIDELITY OF IMPRESSION MATERIALS USED IN ORTHODONTICS

**G Tanase**, I Ionescu, M Burlibasa, R Sfeatcu, C Cristache, University of Medicine and Pharmacy 'Carol Davila', Bucharest, Romania

**AIM:** Impression materials used in orthodontic practice may suffer multiple modifications of physical and dimensional parameters. These modifications, not obvious to the naked eye, may lead to significant defects in removable orthodontic devices. This study aimed to compare *in vitro* the dimensional fidelity and stability of impression materials either silicone or polymeric materials used in orthodontic practice.

**MATERIALS AND METHOD:** Testing was carried out in accordance with the standard SR EN 4823:2002, using a fidelity testing device constructed on a stainless steel test block. The fidelity of elastomeric impression materials were determined both qualitatively and quantitatively using an electronic microscopy (ESEM XL30) at X 300000 magnification. Qualitative analysis of the material fidelity of the elastomeric sample refers to the reproduction test of details (measures of width on the block test, and statistical analysis, ANOVA), at the same standard, and qualitative analysis, at the linear dimensional variation.

**RESULTS:** Different results were recorded for the qualitative and quantitative fidelity of elastomeric impression materials, as well as for the models (rough and extra rough gypsum). Thus, for quantitative and qualitative analysis of fidelity of recent generations of elastomeric, silicone and polieteric impression materials, measurements were obtained by electronic microscope using the environmental scanning electron microscopy technique. The device used proved not to be precise which was confirmed by the electronic microscope findings.

**CONCLUSION:** The degree of fidelity of the detail reproduction is different from an impression material to another one used in this study. Thus, the most suitable samples were identified to be using condensation silicones.

## SP 166

Wednesday, June 16 – at 11:30

### ERRORLESS PHOTOMETRY – IS IT POSSIBLE?

**Z Lendengolts**, E Karton, Z Vagapov, A Eghiazaryan, Department of Orthodontics, Moscow State University of Medicine and Dentistry, Russia

**AIM:** It is important to achieve a natural head position (NHP) when undertaking a photometric study. NHP has been recognized as the most suitable position for cephalometric and photometric studies. When photographing a person's face, to obtain identical images, it is advisable to use a special device that would allow fixation of the NHP. The purpose of this study was to clinically test the development of such a device, the Photostat, for head stabilization.

**SUBJECTS AND METHOD:** Thirty patients with a normal occlusion. The Photostat consisted of a light emitting diode headlamp shell, a bulb from a laser pointer, and a dual-axis level for up/down and left/right control for tilting the head. The Photostat is fastened on the head with a soft rubber strap, and has the opportunity to change the size, depending on the size of the head of the individual. The patient is asked to look straight ahead, as if he/she was looking into their own eyes (mirror technique). The laser pointer, mounted in front of the flashlight, is then turned on. Side sloping, as well as the antero-posterior of the head, was regulated by a dual-axis level, so that the beam of the laser pointer was parallel to the floor. This was visible on the surface in front of the patient (wall, hanging fabric, etc.) allowing the patient to independently control the slope of the head. The patient sits freely and relaxed.

**RESULTS:** Clinical examination showed that the Photostat can be successfully used for positioning the head in a natural position in photographic and radiographic studies. It also provides reliable fixation and standardization to enable comparison of results.

### SP 167

Wednesday, June 16 – at 11:30

#### HUMAN MASTICATORY MUSCLE DIMENSIONS AND VERTICAL FACE HEIGHT - A MRI STUDY

**P Y E Sham**<sup>1,2</sup>, **W C K Foong**<sup>1</sup>, **H P Ng**<sup>3</sup>, **S H Ong**<sup>1</sup>, **Y H Chan**<sup>1</sup>,  
<sup>1</sup>National University of Singapore, <sup>2</sup>National Dental Centre and  
<sup>3</sup>Singapore Bioimaging Consortium, Singapore

**AIM:** The vertical facial dimension in humans varies considerably between individuals. The masticatory muscles have been hypothesized to play a major role in vertical facial development. Published studies show a positive correlation ( $P \leq 0.05$ ) between masticatory muscle volume and cross-sectional area with anterior face height, while some authors found otherwise. These studies used radiographs and computer tomography that exposed subjects to unnecessary radiation. The aim of this research was to explore the relationship between volume and area of masticatory muscles (masseter, medial pterygoid and lateral pterygoid) and anterior face height using magnetic resonance images (MRI).

**MATERIALS AND METHOD:** MRIs of 40 healthy subjects (13 females, 27 males, age range 20-38 years). Computer algorithms for accurate automated image segmentation were used to create patient-specific masticatory muscle models. The volume and maximum cross-sectional area of each masticatory muscle were calculated. Linear measurements of the soft tissue of anterior and lower face height were also measured. The relationship between the maximum cross-sectional area and volume of the masticatory muscles and anterior and lower face height was analyzed using Spearman's rank correlation coefficient. All measurements for the first 20 subjects were repeated and intra-examiner reliability was determined using a Bland-Altman plot.

**RESULTS:** No statistically significant correlation was found between masticatory muscle volume and area with lower or anterior face height. The Bland-Altman plot showed good intra-examiner reliability.

### SP 168

Wednesday, June 16 – at 12:15

#### DEGREE OF MANDIBULAR MOLAR ROOT DEVELOPMENT IN RELATION TO CRANIOFACIAL GROWTH TYPE

**J Andjelic**, Health Centre, Tivat, Montenegro

**AIM:** As there is a direct relationship between growth of the body and face, it is possible to indirectly determine the stage of facial growth. The position and development of the mandibular molars are important for normal growth and development. The aim of this study was to determine the degree of mandibular molar root development in relation to craniofacial growth type.

**SUBJECTS AND METHOD:** Five hundred and thirty eight (238 males, 300 females) aged 13 to 18 years of age who were orthodontically treated. The children had all teeth as expected with normal growth and development for their age. Development of the

mandibular molars was studied on dental pantomographs. The following 10 stage development classification (Demirjian) was used.

**RESULTS:** For all males examined, 48 had horizontal growth, 58 neutral growth and 132 vertical growth. For females, 74 had horizontal growth, 64 neutral growth and 162 vertical growth.

**CONCLUSION:** There was a statistically significant difference between subjects with neutral and vertical growth type. There was no statistically significant difference in mandibular third molar development between subjects with different growth types.

### SP 169

Wednesday, June 16 – at 12:15

#### ELECTRON PROBE MICROANALYSIS OF TEN REPRESENTATIVE NICKEL TITANIUM ARCHWIRES

**Y Bolender**<sup>1</sup>, **L Aranda**<sup>2</sup>, **A Vernière**<sup>3</sup>, **C Rapin**<sup>2</sup>, **M-P Filleul**<sup>1</sup>,  
Departments of <sup>1</sup>Dento-facial Orthopedics <sup>2</sup>Solid-state and  
Surface Chemistry and Physics and <sup>3</sup>Condensed-matter and  
Material Physics, Nancy University, France

**AIM:** To determine the composition of 10 representative nickel titanium (NiTi) archwires by electron probe microanalysis.

**MATERIALS AND METHOD:** Ten upper 0.017 × 0.025 inch NiTi archwires: Nitinol®, Nitinol SE®, Nitinol HA® (3M-Unitek), Neosentalloy® 100 and 200 g (GAC), Copper NiTi® 27°C, 35°C and 40°C (Ormco), Rematitan Lite®, Rematitan Lite White® (coated version) (Dentaurum). The archwires were cut into segments and embedded in cross-section in an epoxy resin. The samples were polished initially with 120 to 4000 grid SiC paper under water, and finally with a 1 µm diamond suspension and lubricant. Wavelength dispersive spectrometry measurements were then taken at 10 different locations of the same cross-section with a SX100 Cameca® microprobe and average values were calculated.

**RESULTS:** Nitinol®, Nitinol SE®, Nitinol HA®, Neosentalloy 100 and 200g® and Rematitan Lite were exclusively made of Ni (~54 to 55 ± 1.0% weight composition) and Ti (~45 to 46 ± 0.6% weight composition). These alloys showed only minute differences in composition. Copper Niti® showed a different content, in as much that Ni weight composition dropped to 48.4 per cent whereas Cu represented 5.7 per cent and Cr 0.2 per cent of their weight. NiTi alloys were nearly equiatomic whereas Cu and Cr represented respectively 4.8 per cent and 0.2 per cent of the atomic composition of Copper NiTi® at the expense of the Ni fraction.

**CONCLUSION:** Two categories of wires can be distinguished by their composition, the standard NiTi alloys and the Copper NiTi®. Since no difference in composition can be detected within these two categories, the distinct mechanical properties of these wires within their own group may be due to their specific thermo-mechanical treatment during the manufacturing process.

### SP 170

Wednesday, June 16 – at 12:15

#### DENTOSKELETAL AND SOFT TISSUE PROFILE CHANGES IN CLASS II HIGH MANDIBULAR ANGLE PREADOLESCENTS

**T Boskovic-Brkanovic**, Department of Orthodontics, Kotor Dental Centre, Montenegro

**AIM:** To show the skeletal, dental, and soft tissue changes in pre-adolescent patients with a Class II malocclusion and high-angle growth pattern treated with a headgear-activator.

**SUBJECTS AND METHOD:** Two groups of 25 subjects aged between 8 and 12 years, sufficient with respect to age, gender, observation time, and dentofacial characteristics. The first group were treated with an activator-headgear for an average period of 22 months until a Class I molar and canine relationship and reduction of overjet was established. The second group (control) did not receive any form of orthodontic treatment. Pre- and post-treatment cephalograms were analyzed and paired t-tests were used for comparison of the differences between the control and treatment group.

**RESULTS:** Compared with the control group, the treatment group showed a decrease of 3.41 mm in forward growth of the maxilla, an increase of 4.06 mm in mandibular protrusion, a decrease of 1.63 degrees in ANB angle, and a decrease in overjet and overbite. There were no significant differences in craniofacial growth between the two groups. Posterior face height showed a similar increase in both groups. The activator appliance caused lingual inclination of the upper incisors, whereas inclination of the lower incisors remained unchanged. The soft tissue profile in the treatment group was significantly improved: a decrease of 3.33 degrees in soft tissue convexity, a decrease of 4.8 mm in protrusion of the upper lip, an increase of 5.4 mm in lower lip protrusion and an increase of 6.29 mm in chin prominence.

**CONCLUSION:** The activator headgear appliance had a positive influence on the anteroposterior aspect of the facial profile. With regard to vertical facial dimensions, whilst the appliance did not significantly change the existing growth pattern, it did not cause additional mandibular inclination or undesirable posterior mandibular rotation.

### SP 171

Wednesday, June 16 – at 12:15

#### COMPARATIVE STUDY OF BOND STRENGTH OF NEW AND RECONDITIONED BRACKETS AND ASSESSMENT OF RESIDUAL ADHESIVE

**B Wendl<sup>1</sup>**, P Muchitsch<sup>1</sup>, M Pichelmayr<sup>1</sup>, H Droschl<sup>1</sup>, W Kern<sup>2</sup>,  
<sup>1</sup>Department of Clinical Orthodontics, University Dental Clinic, Graz and <sup>2</sup>Institute of Chemical Technology of Organic Materials, University of Technology, Graz, Austria

**AIM:** An average rate of bracket loss of approximately 6 per cent is to be expected in daily clinical orthodontic practice. For reasons of economy, detached brackets are commonly reattached after sandblasting to remove adhesive, or replaced with used brackets reconditioned by specialist companies.

**MATERIALS AND METHOD:** Sandblasting and specialist bracket reconditioning procedures were systematically compared by comparative shear testing of re-bonded, reconditioned and new brackets using light and chemically cured adhesives.

**RESULTS:** The mean bond strength of reconditioned brackets was in each case less than that of new brackets, with the lowest value obtained with sandblasted brackets. This nevertheless exceeded the minimum value of 5-8 MPa required for orthodontic use. Bond strength was generally higher with a chemically curing adhesive than with a light curing adhesive; the chemically curing adhesive provided bond strength on previously bonded enamel higher than the light curing adhesive on intact teeth. Consistent with this, the results of an Adhesive Residue Index study demonstrated better bonding of the chemically curing adhesive than the light curing adhesive to the bracket base.

**CONCLUSION:** Despite the weaker bond strength than new brackets, sandblasting brackets accidentally detached during orthodontic treatment will generally allow effective reattachment to be achieved. Bond strength can moreover be improved with the use of a chemically curing adhesive. Used brackets reconditioned by specialist companies provide a second alternative to new brackets and higher bond strengths than sandblasted brackets.

### SP 172

Wednesday, June 16 – at 12:15

#### HOW WELL DO ORTHODONTISTS DIAGNOSE AND RECORD TEMPOROMANDIBULAR DISORDERS?

**S Visram**, D Waring, K D O'Brien, University Manchester Dental Hospital, England

**AIM:** To compare the recording of signs and symptoms temporomandibular disorders (TMD) in a sample of patients examined by consultant orthodontists (uncalibrated and calibrated) and a calibrated TMD examiner. The sample was taken from consecutive patients attending new patient clinics at Manchester Dental Hospital. A sample size of 22 patients was required in the uncalibrated and calibrated phase.

**SUBJECTS AND METHOD:** The study was carried out in two phases. In phase 1 the orthodontic consultants were not calibrated in TMD diagnosis. Twenty-two patients were examined by the orthodontists and the calibrated TMD examiner. Assessment was recorded on data collection sheets. In phase 2, the orthodontic consultants were calibrated in TMD diagnosis and the patients were assessed by the calibrated consultants and calibrated TMD examiner. The level of agreement between the orthodontist and the TMD examiner was assessed.

**RESULTS:** Recording of TMD examination in patients' notes was low in the uncalibrated phase (45.8%) and increased in the calibrated phase (88.5%). For all outcome measures, the level of agreement in diagnosing signs and symptoms of TMD was greater in the calibrated phase compared with the uncalibrated phase. The sensitivity of the TMD examination was higher in the calibrated phase.

**CONCLUSION:** Without calibration, clinicians show low reliability compared with a calibrated TMD examiner for the recording and diagnosis of TMD. The findings highlight the importance of reliable clinical standards for examination and diagnosis of TMD.

### SP 173

Wednesday, June 16 – at 12:15

#### SKELETAL MATURATION IN CHILDREN WITH A CLEFT LIP AND PALATE

**M O Akcam**, M E Emral, Department of Orthodontics, Faculty of Dentistry, Ankara University, Turkey

**AIMS:** To investigate the correlations between chronological age and skeletal age in subjects with a cleft lip and palate (CLP) and to compare it with a Class I control group, according to hand-wrist skeletal maturity indicators.

**MATERIALS AND METHOD:** Hand wrist films of 45 subjects with a CLP (16 males, 29 females, age range 6-17 years) and 45 Class I subjects (17 males, 28 females, age range 8-16 years). In each case skeletal age was determined on the basis of the Greulich-Pyle atlas method using gender specific hand wrist radiographs standardised by age, by one experienced examiner. Statistical evaluation was performed with the Statistical Package for Social Sciences (SPSS Inc., Chicago, Illinois, USA) using analysis of variance (ANOVA), Duncan's test and Pearson correlation analysis.

**RESULTS:** In the overall study group (n = 90) a high correlation was observed between skeletal and chronological age (0.94,  $P < 0.01$ ). In the CLP subjects, skeletal age was retarded (11.4 years) when compared with chronological age (12.3 years), although it was statistically insignificant. However, in the control group skeletal age was similar to chronological age (13.6 versus 13.1 years). In both groups, skeletal age was ahead of chronological age in female subjects when compared with males, while in the male group, skeletal age was behind chronological age (11.7 versus 12.2 years). Skeletal age was retarded in the CLP group when compared with chronological age, both in male (n = 41, 10.8 versus 13.2 years) and female (n = 49, 12.2 versus 13.9 years) subjects.

**CONCLUSION:** As CLP subjects may have a skeletal age behind chronological age, hand wrist evaluation should be considered rather than the chronological age for the timing of orthodontic/orthopaedic treatment.

### SP 174

Wednesday, June 16 – at 12:15

#### USE OF TEMPORARY ANCHORAGE DEVICES IN INTRUSIVE MECHANICS

**M Maddalone**, M Ferrari, P Arrigoni, S Barrilà, A Stanizzi, Università degli Studi di Milano, Bicocca, Italy

**AIM:** To compare intrusive mechanics that use miniscrews with a traditional mechanics.

**SUBJECTS AND METHOD:** Fifty patients aged between 12 and 54 years who required intrusion of the upper or lower incisors from 3 to 7 mm. The patients were randomly assigned to two groups: 25 treated with utility arches and 25 treated with continuous arches

connected with a temporary anchorage device (TAD) for intrusion. The intrusive force was delivered by an elastic chain. The miniscrew was positioned 4-5 mm below the gingival margin between the upper central or lower incisors. The direction of insertion had a variable inclination to avoid contact with the incisor roots. The head of the miniscrews served as a reference point to evaluate the speed of intrusion both for patients that used miniscrews as anchorage and for the control group that used utility arches in the first 4 months of treatment.

**RESULTS:** Intrusion speed with miniscrews was on average higher than with utility arches. In eight of 25 subjects it was impossible to reach the intrusion target with utility arches in the 4 month period. For all 25 patients with a utility arch, molar dislocation was noted in the posterior segments. Two miniscrews were lost in the TAD group at the end of the experimental period, but without effects on intrusion.

**CONCLUSION:** Intrusive mechanic were faster with the use of TADs without distal tipping of the anchorage molars. The miniscrews were well tolerated by the patients during therapy.

### SP 175

Wednesday, June 16 – at 12:15

#### EVALUATION OF ORAL HYGIENE IN PATIENTS TREATED WITH REMOVABLE APPLIANCES

**P Pietrzak**, G Smiech-Slomkowska, Department of Orthodontics Medical University of Lodz, Poland

**AIM:** To evaluate oral hygiene and oral hygiene index (OHI) values in 72 patients aged 7 to 13 years, treated with removable appliances.

**MATERIALS AND METHOD:** Dental instruments were to assess oral hygiene during two control visits. Three teeth in the maxilla (labial surfaces) and three teeth in the mandible (lingual surfaces) on the opposite side were examined. OHI values were measured and the results were summed and divided by the number of surfaces. The correct tooth brushing technique was shown using models and a toothbrush. Each patient was provided with toothpaste (Elmex) for home use. Oral hygiene and tooth brushing technique were evaluated again using the OHI after 2 months.

**RESULTS:** The average OHI was 0.83 (0.93 in boys; 0.75 in girls). The average OHI after 2 months was 0.63 (0.74 in boys; 0.53 in girls). According to Warsz *et al.*, the OHI value in children without proper dental care was 0.32 (0.25 in girls, 0.4 in boys). These OHI values were lower than those found in the present study.

**CONCLUSION:** The OHI was higher in boys than in girls. The OHI value after 2 months was lower both in boys and in girls. No correlation between the OHI and patient age was found.

### SP 176

Wednesday, June 16 – at 12:15

#### CEPHALOMETRIC EVALUATION OF PATIENTS WITH OBSTRUCTIVE SLEEP APNOEA

S H Tuna<sup>1</sup>, O Ozturk<sup>2</sup>, **H Turkkahraman**<sup>3</sup>, H Alkis<sup>3</sup>, A H Balcioglu<sup>4</sup>, Departments of <sup>1</sup>Prosthodontics, <sup>2</sup>Respiratory Disease, Sleep Center and <sup>3</sup>Orthodontics, Suleyman Demirel University, Isparta and <sup>4</sup>Department of Anatomy, Istanbul University, Turkey

**AIM:** To evaluate the cephalometric features of patients with obstructive sleep apnoea (OSA), and to elucidate the relationship between cephalometric variables and severity of the apnoea-hypopnoea index (AHI).

**SUBJECTS AND METHOD:** Ninety-three patients with OSA, diagnosed by overnight polysomnography, were classified into three subgroups according to the AHI: mild (5-15, n = 24), moderate (16-30, n = 35) and severe (>30, n = 34) OSA. Twelve cephalometric measurements were carried out for all patients to clarify the relationship between cephalometric variables and the severity of AHI. One-way analysis of variance, Mann Whitney U, Pearson correlation test and stepwise regression analysis were performed with the Statistical Package for Social Sciences, version 16.0 (SPSS Inc., Chicago, Illinois, USA)

**RESULTS:** Most male patients (38%) were classified with severe OSA and most females as moderate OSA (40.9%). IAS, MaxSoftPalate, SNGoGn, HyoidMPPerp variables were found to be higher in subjects of both genders with severe OSA. IAS, MAS, MaxSoftPalate and HyoidMPPerp variables were positively correlated with AHI ( $P < 0.001$  and  $P < 0.05$ ), however SPAS was negatively correlated ( $P < 0.05$ ). Stepwise regression analysis; mean SaO<sub>2</sub>, arousal index, Body Mass Index and MAS were included in the model to estimate AHI properties with R square 67 per cent ( $P < 0.01$ ).

**CONCLUSION:** The values of IAS, MaxSoftPalate, SNGoGn and HyoidMPPerp increased with the severity of OSA. However, the cephalometric variables used to estimate AHI severity had no role in males, but PNStoPA had an influence in females. Overall, the cephalometric measurements used in this study had limited accuracy in estimating AHI severity in both genders.

### SP 177

Wednesday, June 16 – at 12:15

#### USING ULTRAVIOLET LIGHT TO ARTIFICIALLY AGE ELASTOMERIC CHAIN

S Hj Abd Wahab<sup>1</sup>, **D Bister**<sup>1</sup>, M Sherriff<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Biomaterials, King's College London Dental Institute, England

**AIMS:** To investigate the effect of ultraviolet type A light (UVA) exposure on the tensile properties of elastomeric chain and to compare its effect to the 'as received' product. UVA light exposure was used as a model for artificial ageing simulating prolonged storage of elastomeric chain.

**MATERIALS AND METHOD:** The three products tested were from Ormco, Forestadent and 3M. Tensile strength (n = 60) was measured after exposing the chains to UVA light for 0, 2, 3 and 4 weeks. Force decay was measured (n = 60) using chains that were artificially aged for 5, 10 and 14 days. The chains were subsequently stretched at a constant distance and the resulting forces were measured at 0, 1, 24 hours, 7, 14, 21 and 28 days. This test was used to simulate a clinical scenario of pre-stretching and subsequent shortening of elastomeric chain during orthodontic space closure.

**RESULTS:** In the tensile strength test, statistically significant differences were found that were related to the duration of UVA light exposure. The Forestadent chain, which had the second highest value for the as received product, showed the most consistent values over time with the lowest degradation. The Ormco product showed the lowest as received values as well as after UVA exposure; the 3M product had the highest loss of tensile strength. Significant differences were also found among chains in the force decay test. UVA light exposure of 10 days or more appears to mark a 'watershed' between the products – the 3M product had the most intact chains, while the Forestadent chain had some survivors, depending on the time the chain was stretched for. None of the Ormco product survived UVA light exposure for more than 5 days in the force decay test.

**CONCLUSION:** UVA light exposure may be used as a model for artificial ageing as it reduces force delivery and tensile strength of exposed chains.

### SP 178

Wednesday, June 16 – at 13:00

#### IN VITRO PEAK INSERTION TORQUE VALUES OF SIX COMMERCIALY AVAILABLE MINI-IMPLANTS

**C Z-Y Whang**<sup>1</sup>, D Bister<sup>1</sup>, M Sherriff<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Biomaterials, Biomimetics and Biophotonics, King's College London Dental Institute, England

**AIM:** Mini-implants are a popular adjunct of orthodontic treatment. There are a number of unwanted potential side-effects associated with the use of mini-implants, such as fracture during insertion. The aim of this study was to compare peak insertion torque values.

**MATERIALS AND METHOD:** Six commercially available self-drilling mini-implants [Mini Spider Screw®, Health Development Company (1.5 X 8 mm), Infinitas®, DB Orthodontics (1.5 X 9 mm), Vector TAS®, Ormco (1.4 X 8 mm), Dual Top®, Jeil (1.6 X 8 mm), Tomas Pin®, Dentaurum (1.6 X 8 mm) and Ortho-Easy®, Forestadent (1.7 X 8 mm)]. Twenty implants each were drilled into acrylic rods at eight rotations per minute using a motorized torque measurement stand. An additional 20 Ortho-Easy® implants per group were tested at a slower speed of four rotations per minute and with 6 and 10 mm lengths, to assess the impact of speed and length.

**RESULTS:** There were significant differences for some of the maximum torque values for the different mini-implants with the same length. The Mini Spider Screw and Infinitas showed the lowest average torque values (in Ncm) 6.5 and 12.4, compared with Vector TAS, Dual Top, Tomas Pin and Ortho-Easy implants (30.9, 29.4, 25.4 and 24.8 respectively). The Tomas Pin showed the largest standard deviation (7.7); the Dual Top was the most consistent (0.6). Different insertion speeds did not result in significant differences in peak torque values for Ortho-Easy® implants but the 6 mm implant was significantly stronger than those measuring 8 and 10 mm.

**CONCLUSION:** The different torque values could be due to the variation of core diameter, implant shape and/or composition of the Titanium alloy. Differences in production technique and alloy could account for variability of standard deviations. Although clinical torque values rarely exceed 12 Ncm, pre-drilling cortical bone to reduce insertion torque values appears justified for some implants to reduce the risk of fracture during insertion.

## SP 179

Wednesday, June 16 – at 13:00

### CLEFT LIP/PALATE IN MAYO, IRELAND 1999–2007

J M Waldron<sup>1</sup>, J R Sandy<sup>2</sup>, A R Hewson<sup>1</sup>, C M McNamara<sup>1</sup>, <sup>1</sup>Dental & Orthodontic Services, Ireland and <sup>2</sup>University of Bristol, England

**AIM:** A cleft lip and palate (CL/P) is the most common congenital abnormality in the craniofacial region. The incidence varies according to race/ethnicity, gender, family history and cleft type. The population of County Mayo, Ireland was 123,839 in 2006. The aims of this study were to identify and classify all cleft-affected children born in Mayo, Ireland between January 1999 and December 2007, and to assess their dental status.

**MATERIALS AND METHOD:** Dental records centralised at Mayo General Hospital were reviewed. Two clinicians carried out a clinical examination, along with completing a standardised questionnaire regarding the subject's medical history.

**RESULTS:** Thirteen CL/P subjects were identified, all Caucasian, nine males (69%), four females (31%) with a M:F of 2.25:1. The mean age was 5 years 10 months (range of 2 years 1 month, to 9 years 4 months). Four (31%) had CL/P, seven (54%) an isolated CP, and two (15%) an isolated cleft lip. Males predominated the CLP group and females the CP group. Twenty-three per cent were syndromic. A family history existed in 31 per cent. Fifty-four per cent required ear, nose and throat assessments. Sixty-two per cent had received speech therapy. The mean age at lip repair was 4.16 months and 11 months at palatal repair. The age at the first dental visit ranged from 3 months to 7 years 1 month (mean of 21 months). The mean number of decayed missing and filled teeth was 0.037. Fifty-four per cent had received fissure sealants. Fifteen per cent presented with hypodontia. Eight-five per cent had an orthodontic malocclusion.

**CONCLUSION:** From 1999 to 2007, the most common cleft anomaly in Mayo, Ireland was an isolated CP. The findings, including gender, cleft type, side and associated dental findings, followed international trends. Despite dental access being available from infancy, further investigation is necessary to determine the wide age range for the first dental visit found in this study.

## SP 180

Wednesday, June 16 – at 13:00

### META-ANALYSIS OF STUDIES ON PAEDIATRIC CRANIOMANDIBULAR DISORDERS

A Pilotto, G Galluccio, E Barbato, Department of Orthodontics, University of Rome Sapienza, Italy

**AIM:** The signs and symptoms of temporomandibular disorders (TMD) in children have been intensively studied. A mainly high, but variable, prevalence of TMD is reported in the literature, but it remains unclear whether TMD in children and adolescents should be considered a major health problem. The aim of this meta-analysis was to evaluate the incidence of paediatric craniomandibular disorders.

**MATERIALS AND METHOD:** Four hundred and ninety four articles identified from PubMed were collected and 97 were selected by applying the following criteria: subject's age 0–14 years; condition not treated; report published after 1990; more than 100 subjects surveyed; and TMD evaluated with auscultation, muscle palpation, and mandibular movement capacity.

**RESULTS:** The reported incidence of paediatric TMD was very high, at 27.3 per cent. Children with TMD can present with bruxism, onychophagy, speech disorders, partial inability to open the mouth, mastication difficulties, and malocclusions.

**CONCLUSION:** The identification and recognition of factors such as malocclusions and parafunctions are considered fundamental to early diagnosis of problems of the temporomandibular joint, which is the most effective way to avoid dysfunction of the stomatognathic system.

## SP 181

Wednesday, June 16 – at 13:00

### EFFECTS OF CONTINUOUS AND INTERMITTENT CONTROLLED FORCES ON CANINE DISTALIZATION

S Usumez<sup>1</sup>, Z M Sari<sup>2</sup>, S Hakki<sup>3</sup>, T Ataoglu<sup>3</sup>, <sup>1</sup>Department of Orthodontics, University of Gaziantep and Departments of <sup>2</sup>Orthodontics and <sup>3</sup>Periodontology, Selcuk University, Konya, Turkey

**AIM:** The duration of force application is an important factor, together with the magnitude of force, in stimulating osteoclastic activity. There is still ambiguity about whether continuous or intermittent orthodontic forces produce more efficient canine distalization. The aim of this study was to evaluate the effects of continuous and intermittent orthodontic forces on canine distalization and to compare the effectiveness of laceback ligatures with that of superelastic nickel titanium (NiTi) closed coil springs.

**SUBJECTS AND METHOD:** Ten subjects for whom fixed orthodontic treatment was planned with the extraction of first premolars to resolve crowding in the upper and/or lower arches. Stainless steel direct-bonded Roth brackets (0.018 inch) were used. For canine distalization, superelastic NiTi closed coil springs generating a force of 125 g were used on one side while lacebacks made of 0.010 inch wire were used on the contralateral side. Lateral cephalometric and panoramic radiographs and dental casts were used to evaluate the dental changes. Wilcoxon test was applied to determine the differences between pre- and post-distalization mean values and the mean differences between the groups.

**RESULTS:** The duration of distalization was significantly shorter ( $P < 0.001$ ) and mesiopalatal rotation of the anchorage molar was significantly greater in the coil spring group ( $P < 0.05$ ). No significant differences were observed in the amount of canine distalization, amount of molar mesialization, distal tipping of the canine, or mesial tipping of the molar.

**CONCLUSION:** The intermittent forces created using laceback ligatures proved to be effective for canine distalization and resulted in less mesiopalatal rotation of the anchorage molars. This, however, required double the time for the same amount of canine distalization compared with continuous forces using NiTi closed coil springs.

**SP 182**

Wednesday, June 16 – at 13:00

**STRUCTURAL ADAPTATION OF RABBIT JAW MUSCLES TO ALTERED MASTICATORY FUNCTIONAL LOAD****M Vreeke**, G E J Langenbach, J A M Korfage, A Zentner, T Grünheid, Academic Centre for Dentistry Amsterdam (ACTA), Netherlands

**AIM:** Skeletal muscle fibres can change their myosin heavy chain (MyHC) isoform and cross-sectional area (CSA) to adapt to altered environmental conditions. In general, reduced muscle activity induces transition towards faster fibres (MyHC IIX > IIA > cardiac alpha > I) with smaller CSA. The aim of this study was to investigate the effect of a reduction in masticatory functional load on MyHC composition and fibre CSA of three functionally different jaw muscles in a rabbit model.

**MATERIALS AND METHOD:** MyHC composition and the corresponding CSA of fibres were determined in the masseter, temporalis and digastric muscles of male rabbits aged 20 weeks. Between 8 and 20 weeks of age the experimental animals (n = 8) had been fed pellets requiring a significantly lower force to break the pellet (10 N) in comparison with the standard pellets (120 N) fed to the control animals (n = 8).

**RESULTS:** The proportion and CSA of the slow-type fibres co-expressing MyHC-I and MyHC-cardiac alpha were significantly lower in the masseter muscles of the experimental animals than in those of the control animals ( $P < 0.05$ ). In contrast, the proportions and CSAs of the various fibre types in the temporalis and digastric muscles did not differ significantly between the groups of animals ( $P > 0.05$ ).

**CONCLUSION:** Long-term reduction in masticatory functional load contributes to selective disuse of jaw muscles. The muscles affected adapt structurally to the reduced functional load with decreases in the proportion and CSA of their slow fibres. Supported by a European Orthodontic Society research grant.

**SP 183**

Wednesday, June 16 – at 13:00

**CHARACTERIZATION OF AS-RECEIVED AND IN VIVO ORTHODONTIC STAINLESS STEEL BRACKETS****S Hoste**<sup>1</sup>, J-P Celis<sup>2</sup>, G Willems<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Metallurgy and Material Engineering, Catholic University of Leuven, Belgium

**AIM:** Contemporary orthodontic treatment mostly involves the use of bonded brackets, which can be manufactured from different materials. Most commonly used are stainless steel brackets. These brackets are predominantly made of austenitic stainless steel due to its superior corrosion resistance, excellent formability, and low cost. Because of the manufacturing process, variable amounts and different types of surface defects can be included. These surface defects provide predilection spots for the onset of corrosion. Additionally, galvanic corrosion can occur not only between the wire and the bracket, but also within the bracket if the wing and base material are not identical. The aim of this research was a topographical analysis of the bracket slot of as-received and used brackets and characterization of the alloys of two types of brackets.

**MATERIALS AND METHOD:** Standard edgewise (GAC Dentsply, New York, USA) and mini twin (Ormco Corporation, Glendora, California, USA) brackets. The brackets were collected, cleaned and coated with a thin gold layer for scanning electron microscopic analysis. Magnification went up to X20000. Elemental analysis of the bracket base, wing and brazing material was performed to determine the alloys of each bracket.

**RESULTS:** As-received standard edgewise brackets showed more plastic deformation than the mini twin brackets. Pits, crevices and grain boundaries were observed on all brackets. There was an increase of plastic deformation and scratches on both types of brackets. Elemental analysis revealed that the base and wing material were not the same either in the GAC or Ormco brackets. A different brazing material was used by both manufacturers.

**CONCLUSION:** As-received brackets show different degrees of surface defects. These defects tend to increase after intra-oral use. As a consequence of the different alloys in each bracket, galvanic corrosion may occur.

**SP 184**

Wednesday, June 16 – at 13:00

**EVALUATION OF SKELETAL CHANGES INDUCED BY SURGICALLY ASSISTED RAPID PALATAL EXPANSION****G Iodice**, A Laino, T Bocchino, R Del Giglio, R Martina, University of Naples Federico II, Italy

**AIM:** A Class II anterior open bite and/or a steep mandibular plane angle are frequently considered a contraindication to surgically assisted rapid palatal expansion (SARPE). Nevertheless, few studies have investigated the maxillary and mandibular effects after SARPE on the sagittal and vertical planes. The aim of the present research was to evaluate the sagittal and vertical effects after SARPE.

**SUBJECTS AND METHOD:** Twenty-one consecutive adult patients (7 males, 14 females; mean age  $25.6 \pm 6.3$  years). A subtotal Le Fort I osteotomy with pterygomaxillary disjunction was carried out in all patients. Lateral cephalometric radiographs were taken during pre-operative assessment (T0) and 6 months after the end of expansion (T1). The cephalometric measurements on the radiographs were: SNA, SN-Sna; Frankfort horizontal plane^NA, SN^palatal plane, CF-A, Frankfort horizontal plane^palatal plane, SNB, SnPg, SN^GoGn, mandibular plane angle (FMA), upper anterior face height/total anterior face height, lower anterior face height/total anterior face height, upper posterior face height/total posterior face height, total anterior face height/total posterior face height, upper incisor^NA, upper incisor^SN, upper incisor^, Frankfort horizontal plane. Independent sample t-tests and ANOVA analyses were used.

**RESULTS:** Statistically significant changes were observed only for upper incisor^NA ( $P = 0.04$ ). No skeletal sagittal or vertical variation was found after SARPE.

**CONCLUSION:** No sagittal and/or vertical skeletal displacement could be linked to SARPE treatment. Consequently, Class II, anterior open bite and/or a steep mandibular plane angle cannot be considered an outright contraindication to its use. Upper incisors retroclination could result after SARPE.

**SP 185**

Wednesday, June 16 – at 13:00

**ANTERIOR MAXILLARY AND MANDIBULAR INTERRADICULAR DIMENSIONS AND CORTICAL BONE THICKNESS****M Fayed**<sup>1,2</sup>, P Pazera<sup>1</sup>, C Katsaros<sup>1</sup>, <sup>1</sup>Department Of Orthodontics and Dentofacial Orthopedics, Bern University, Switzerland and <sup>2</sup>Department of Orthodontics, Cairo University, Egypt

**AIMS:** There are many factors that affect mini-implant success, including implant location related factors such as interradicular distance, buccolingual bone depth, and cortical bone thickness. Although several studies have assessed these factors in the posterior region of the jaws, only limited information is available as regards the anterior region. The aim was to study the interradicular distance, buccolingual bone depth, and cortical bone thickness in the anterior region of the maxilla and mandible using cone beam computed tomography (CBCT), and to detect possible differences associated with age and gender.

**MATERIALS AND METHOD:** CBCT images of 100 subjects (46 males, 54 females) divided into two age groups (13-18 years) and (19-27 years). The following interradicular measurements were performed: buccolingual (BL) bone thickness, mesiodistal interdicular spaces both buccally (MD-B) and palatally/lingually (MD-P/L), and buccal (BC) and palatal/lingual (P/LC) cortical thickness.

**RESULTS:** In the maxilla the highest BL ( $9.6 \pm 2.16$  mm) was found between the central and lateral incisors. The highest MD-B ( $4.27 \pm 1.23$  mm) was between the central incisors. The highest MD-P ( $3.73 \pm 1.28$  mm), and BC ( $1.24 \pm 0.53$  m) were between the lateral

incisor and canine. The greatest PC ( $1.85 \pm 0.64$ ) was between the central and lateral incisor. In the mandible the highest BL, MD-B/L and BC, LC were found between the lateral incisor and canine ( $7.83 \pm 1.36$ ,  $3.89 \pm 1.33$ ,  $3.12 \pm 1.51$ ,  $1.24 \pm 0.19$  and  $2.36 \pm 0.53$  mm, respectively. Males and subjects in the older age group had significantly higher BL, BC, and PC in the maxilla. No significant differences were found in the mandible.

**CONCLUSION:** For mini-implant placement anteriorly, the highest bone dimensions and cortical thickness exists between the central and lateral incisor and lateral incisor and canine in the maxilla, and between the lateral incisor and canine in the mandible. Males and older subjects had significantly higher buccolingual bone depth and cortical bone thickness only in the maxilla.

## SP 186

Wednesday, June 16 – at 13:00

### WHITE SPOT LESIONS AFTER ORTHODONTIC TREATMENT – THREE MONTHS USE OF CASEIN-PHOSPHOPEPTIDE AMORPHOUS-CALCIUM-PHOSPHATE FLUORIDE PASTE

M W Beerens, M Stevens, A H Toxopeus, M H van der Veen, Academic Centre for Dentistry Amsterdam (ACTA), Netherlands

**AIM:** Casein-phosphopeptide amorphous-calcium-phosphate fluoride paste (CPP-ACP-F) may promote remineralization of white-spot lesions (WSL) by supplying bioavailable calcium and promoting a healthier plaque. The aim of this study was to evaluate the effect of CPP-ACP-F paste versus a placebo paste, when WSL were present after fixed appliance treatment. Changes in the size and depth were monitored using quantitative light fluorescence during a three month period.

**SUBJECTS AND METHOD:** Initial buccal carious lesions were determined in 51 adolescent orthodontic patients at debond. Subjects with WSL were given CPP-ACP-F paste ( $n = 26$ ) or a placebo paste ( $n = 25$ ) to use once a day at bedtime. At debond and after three months, carious lesions were examined by QLF. Integrated fluorescence loss (IFL) was determined for every lesion.

**RESULTS:** A total of 30 female and 21 male adolescent orthodontic patients, with an average age at debond of 15 years 3 months, participated. A total of 945 elements were monitored, of which 59.8 per cent were affected. Two hundred lesions in the CPP-ACP-F-group and 365 in the placebo-group were found. Reduction of total IFL was seen in 65.4 per cent of subjects in the CPP-ACP-F group and 56 per cent in the placebo group. Overall there was no significant difference in improvement between the groups ( $P = 0.391$ ).

**CONCLUSION:** It is assumed that lesions developed during treatment tend to improve after appliance removal. The use of CPP-ACP-F paste; however, may improve the lesions by decreasing the lesion size and depth after appliance removal and this is supported by the data, however lesion reduction in size and depth were not statistically significantly different between the groups after three months. A randomized clinical trial with a longer follow-up should be performed.

## SP 187

Wednesday, June 16 – at 13:00

### EFFECT OF CASEIN-PHOSPHOPEPTIDE AMORPHOUS-CALCIUM-PHOSPHATE FLUORIDE PASTE ON PLAQUE COMPOSITION AFTER ORTHODONTIC TREATMENT

N C W van der Kaaij, L S Bonhof, M W Beerens, M H van der Veen, Academic Centre for Dentistry Amsterdam (ACTA), Netherlands

**AIM:** Casein-phosphopeptide amorphous-calcium-phosphate fluoride paste (CPP-ACP-F) may promote remineralization of white spot lesions (WSL) by supplying bioavailable calcium and promoting a healthier plaque. The aim of this pilot study was to test the effect of CPP-ACP-F versus a placebo paste on plaque composition after fixed-appliance therapy in patients with multiple WSL after orthodontic treatment.

**SUBJECTS AND METHOD:** Plaque composition was determined at debonding in 34 adolescent orthodontic patients. Patients with multiple WSL were given CPP-ACP-F ( $n = 10$ ) or a placebo ( $n = 10$ )

to use once-a-day at bedtime. WSL were determined by quantitative light-induced fluorescence. Bacterial counts in plaque composition were determined (anaerobic, 37°C, 72 hours) on blood agar (total counts), BHI (pH 5, aciduric bacteria), Rogossa (lactobacilli) and TYCSB (*S. mutans*).

**RESULTS:** The number of WSL at debonding was comparable: CPP-ACP-F 86; placebo 97. Total counts for all groups and times were 5.107 CFU. Aciduric bacteria and *S. mutans* comprised 46 and 11.5 per cent of plaque in the caries group compared with 26 and 3.2 per cent in the caries-free group ( $P < 0.05$ ). After 3 months, aciduric bacteria and *S. mutans* in the CPP-ACP-F group reduced significantly to 25.6 and 3.8 per cent ( $P < 0.05$ ) but were stable for the placebo group. Small colonies of lactobacilli were found that comprised 0.1 per cent of the total flora.

**CONCLUSION:** Plaque flora in orthodontic patients contained a large proportion of acidic bacteria. The level of cariogenic bacteria had not reduced in the placebo group 12 weeks after debond. CPP-ACP-F contributes to a healthier plaque composition. A longer follow-up to investigate the effect of CPP-ACP-F on lesion reduction is indicated.

## SP 188

Wednesday, June 16 – at 13:00

### RE-BOND STRENGTH OF THE BONDED LINGUAL WIRE RETAINER

K van Westing<sup>1</sup>, T Algera<sup>1</sup>, C J Kleverlaan<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Dental Materials Science and Orthodontics (ACTA), Amsterdam, Netherlands

**AIM:** Although orthodontic retainers fail frequently, there is no optimal procedure for re-bonding. The aim of this *in vitro* study was to evaluate the initial bond strength and re-bond strength of retainers bonded to enamel surfaces with and without composite remnants.

**MATERIALS AND METHOD:** The initial bond strength and re-bond strength of Pentaflex lingual wire retainers bonded with Excite and Tetric Flow were determined. The retainers were bonded on three different surfaces; (i) clean enamel, (ii) enamel on which the composite had been removed by a tungsten carbide bur, and (iii) enamel with cured composite remnants. Bond strength was determined with a cantilever/tensile bond strength test (in N) with a repeated crossover design where each tooth was re-bonded twice and tested three times ( $n = 6 \times 19$ ). Fracture was evaluated using the adhesive remnant index (ARI).

**RESULTS:** Comparison of mean bond strengths did not show significant differences for the three different enamel surfaces, but the specimens with the cured composite remnants showed a higher standard deviation. ARI scores showed that 96.5 per cent of bond fractures occurred at the retainer/resin interface. The standard deviations for the re-bonded on composite remnants indicated that leaving composite remains on the surface will lead to a clinically less reliable and less effective bond. In contrast to the *in vitro* ARI score, the clinical ARI scores also showed failure at the resin/enamel interface. This type of failure is most likely caused by wet bonding conditions.

**CONCLUSION:** To achieve optimal bonding properties for re-bonding, the enamel surface should be free of old composite remnants bonded in a moisture-free environment.

## SP 189

Wednesday, June 16 – at 13:00

### INVESTIGATION OF MANDIBULAR ROTATION WITH VARIOUS CEPHALOMETRIC PARAMETERS

O Uslu Akçam<sup>1</sup>, Z Novruzov<sup>2</sup>, <sup>1</sup>Department of Orthodontics, School of Dentistry, Ankara University, Turkey and <sup>2</sup>Clinic of Stomatology, Department of Pediatric Stomatology, Azerbaijan Medical University, Baku, Azerbaijan

**AIM:** The most commonly used measurement to determine the vertical position of the mandible in cephalometric analysis is GoGn/SN angle. Gonial angle is an alternative to this measure-

ment. The aim of this study was to evaluate the indicative factors of mandibular rotation (GoGn/SN, GoMe/FH, gonial angles) and the relationship between them.

**SUBJECTS AND METHOD:** Sixty-nine patients who had completed growth and who did not have craniofacial deformities or congenitally missing teeth. Twenty-four were skeletal Class I (ANB between 0-4°), 20 skeletal Class II (ANB >4°) and 25 skeletal Class III (ANB <0°). Measurements were considered on the pre-treatment lateral cephalometric radiographs.

**RESULTS:** For GoGn/SN angle, the smallest values were observed in the Class I group, while the largest values were in Class II group. GoMe/FH angle was increased in Class I, Class II and Class III cases, whilst gonial angle was increased as Class I, Class III and Class II respectively. As GoGn/SN angle was associated with the location of Nasion, it may not reflect the actual position of the mandibular plane.

**CONCLUSION:** GoMe/FH angle is more reliable in determining the vertical position of the mandible.

## SP 190

Wednesday, June 16 – at 13:45

### POSTERIOR CROSSBITE AND SUCKING HABITS IN CHILDREN FOLLOWED BETWEEN 3 AND 7 YEARS OF AGE

L Dimberg<sup>1</sup>, L Bondemark<sup>2</sup>, B Söderfeldt<sup>3</sup>, B Lennartsson<sup>1</sup>,  
<sup>1</sup>Department of Orthodontics, Postgraduate Dental Education Center, Örebro and Departments of <sup>2</sup>Orthodontics and <sup>3</sup>Oral Public Health, Faculty of Odontology, Malmö University, Sweden

**AIMS:** Longitudinal studies following the development of occlusion are rare. Therefore a longitudinal study was started in 2003 to observe the development of the occlusion and sucking habits between 3 and 7 years of age. The aims were to analyze the prevalence of a posterior crossbite in a group of children followed longitudinally, and to evaluate if the posterior crossbite is influenced by sucking habits.

**SUBJECTS AND METHOD:** Four hundred and fifty seven, 3-year-old children (234 girls, 223 boys) examined by one experienced orthodontist. In the follow-up, 386 children at 7 years of age were re-examined by the same examiner. Data from the clinical examination and a questionnaire were used to determine malocclusion traits and sucking habits.

**RESULTS:** The prevalence of a posterior crossbite was 19 per cent (89/457) at 3 years of age and 18 per cent (68/386) at 7 years of age. If the child had a posterior crossbite at 3 years of age it had a 28 times increased risk of having a posterior crossbite also at 7 years of age. In 6 per cent of the children with a normal transverse relationship at baseline, a new posterior crossbite was developed between 3 and 7 years of age. Spontaneous correction occurred in 27 of the 76 children with a posterior crossbite. The prevalence of sucking habits decreased from 66 to 3 per cent between 3 and 7 years of age.

**CONCLUSION:** When a posterior crossbite is present at 3 years of age, the possibility was high that it remained at 7 years of age. The influence of sucking habits on the prevalence of posterior crossbite at 3 years of age was high, but not confirmed at 7 years of age.

## SP 191

Wednesday, June 16 – at 13:45

### CLINICAL EFFICACY OF POLYVINYLPIRROLIDONE-SODIUM HYALURONATE FOR THE TREATMENT OF ULCERS CAUSED BY ORTHODONTIC APPLIANCES

S-M Lee, B-H Yang, M-S Yun, J-H Lee, Y-H Jung, Department of Clinical Orthodontics, Graduate School of Clinical Dentistry, Hallym University, Korea

**AIM:** To evaluate the efficacy of polyvinylpyrrolidone-sodium hyaluronate (Alocclair™) in reducing pain caused by orthodontic appliances.

**SUBJECTS AND METHOD:** Sixty consecutive patients who were prescribed Alocclair™. The patients were divided into three groups and their pain and discomfort levels were investigated using

three different questionnaires. The three questionnaires included the following: 1. An additional appliance was used; 2. Patients with ulcers caused by orthodontic appliance; 3. Initial placement of brackets. Before and after pain were compared. Visual analogue scales were used to assess the extent of pain.

**RESULTS:** The mean time for reducing pain and the lasting effect of Alocclair™ was 7 minutes and 3.2 hours, respectively. Most patients reported rapid pain relief and found Alocclair™ easy and more pleasant to use than wax. Alocclair™ provided rapid relief of resting and provoked lower pain.

## SP 192

Wednesday, June 16 – at 13:45

### EFFECTS OF SALVIA MILTIORRHIZA ON OSTEOBLASTIC CELL LINE OF MC3T3-E1

A Chin, Y Q Yang, L Chai, R W K Wong, A B M Rabie, Department of Orthodontics, University of Hong Kong, SAR China

**AIM:** Orthodontic tooth movement is mediated by bone remodelling. Sufficient bone is essential for the success and effectiveness of orthodontic treatment. It has previously been found that *salvia miltiorrhiza* extract (SM), a commonly used Chinese herb in cardiac disease to enhance blood perfusion, could increase bone formation *in vivo*. The aim of this study was to investigate the effects of SM on bone cells *in vitro*, to obtain a better understanding on how SM can promote bone remodelling.

**MATERIALS AND METHOD:** MC3T3-E1, an osteoblastic cell line, was cultured with SM for different time intervals (24, 48 and 72 hours), whereas the control group consisted of cells cultured without any intervention. The mRNA expressions of alkaline phosphatase (ALP), osteocalcin (OCN), osteoprotegerin (OPG) and the receptor activator of nuclear factor kappa β ligand (RANKL) were examined by real-time polymerase chain reaction.

**RESULTS:** The expression of ALP showed an early increase at 24 and 48 hours by 50 and 13 per cent, respectively ( $P < 0.001$ ). OCN was decreased by 22 per cent at 24 hours ( $P < 0.001$ ) but increased by 50 and 88 per cent at 48 and 72 hours, respectively ( $P < 0.001$ ). OPG was up-regulated by 10 per cent at 48 hours ( $P < 0.01$ ) and 68 per cent at 72 hours ( $P < 0.001$ ), while RANKL showed an early increase at 24 and 48 hours by 45 ( $P < 0.001$ ) and 36 ( $P < 0.01$ ) per cent, respectively. OPG/RANKL was first down-regulated and then up-regulated.

**CONCLUSION:** SM enhances bone remodelling by regulating the gene expressions of ALP, OCN, OPG and RANKL. It is a potential medicinal herb that could be utilized in the application that requires stimulation in bone cell activities.

## SP 193

Wednesday, June 16 – at 13:45

### CEPHALOMETRIC ASSESSMENT OF RAPID MAXILLARY EXPANSION IN CLASS II DIVISION 1 MALOCCLUSION SUBJECTS\*\*\*

M Pacurar<sup>1</sup>, I N Zetu<sup>2</sup>, A M Jurca<sup>1</sup>, S Kecsed<sup>1</sup>, Universities of Medicine and Pharmacy, <sup>1</sup>Tg. Mures and <sup>2</sup>Iasi, Romania,

**AIM:** Rapid maxillary expansion (RME) is routinely used for correction of transverse maxillary deficiency, a posterior crossbite, crowding and snoring with nasal obstruction. A common secondary effect of RME is anterior mandibular sliding also necessary in Class II malocclusion correction. The aim of this study was to assess the amount of mandibular anterior sliding as a result of the use of RME, by analysis of facial profile parameters.

**SUBJECTS AND METHOD:** Twenty-three patients (12 females, 11 males) with a Class II division 1 malocclusion, who underwent RME. All subjects had a maxillary width deficiency, at least half a cusp distalization associated with a posterior crossbite. Their average age was 12-15 years. The subjects and/or their parents activated the screw once daily. The widening procedure continued until 3-5 mm over-expansion was obtained (in maxillary intermolar distance). To evaluate Class II malocclusion correction, the facial and occlusal improvements and the lateral cephalograms

obtained before and after treatment (McNamara, Burstone and Ricketts method) were assessed. Horizontal and vertical mandibular length (Go-Pog), Ar-Go, convexity of the skeletal profile A-Npog and soft tissue parameters chin thickness (Pog-Pog) upper and lower lip thickness (UL-E line, LL-E line) were determined.

**RESULTS:** RME resulted in spontaneous forward movement of the mandible and Class II correction in 14 patients (the skeletal parameters were increased). The soft tissue modifications were less in comparison with dentoalveolar and bone. The remaining nine subjects required associated Class II interarch mechanics. No significant differences were found in the soft tissue parameters following use of the Class II mechanics.

**CONCLUSION:** There are many favourable effects of RME in Class II division 1 malocclusion patients, one being anterior mandibular growth with less facial profile convexity.

## SP 194

Wednesday, June 16 – at 13:45

### POSTURAL INVESTIGATIONS IN CHILDREN AND JUVENILES WITH AND WITHOUT A CROSSBITE

**S Kopp**, B Büntemeyer, D Ohlendorf, Department of Orthodontics, Goethe-University, Frankfurt am Main, Germany

**AIM:** The success of crossbite therapy is not always guaranteed. The reasons are varied. Therefore, it is important to determine if the sensorimotor system may have an influence on therapy. The aim of this research was to determine the interdependency between a lateral crossbite and the sensorimotor system in children and juveniles.

**SUBJECTS AND METHOD:** Sixty-five children and juveniles (28 males, 37 females). Thirty-two had a crossbite. All subjects were posturographic examined and compared with each other. The Interactive Balance System (Germany) was used. It differentiated on four force plates vertical forces in the forefoot and backfoot areas. For eight standardized test items analytical frequency treatment of the force-time-signal was possible. The parameters were evaluated with the Statistical Package for Social Sciences, version 16.00 (SPSS Inc., Chicago, Illinois, USA) and BIAS, version 9. After Kolmogorov-Smirnov testing, as the results showed that the samples followed a normal distribution, the t-test was used.

**RESULTS:** No significant differences were found between the two groups in the parameters of the sensorimotor system. No difference in the function of sensorimotor subsystems was found between patients with or without a crossbite by posturometric measurements.

**CONCLUSION:** A crossbite does not seem to influence either postural stability or the fall index in children and juveniles. It is feasible that a crossbite can cause body asymmetry after many years of manifestation and thus early preventive treatment should be carried out.

## SP 195

Wednesday, June 16 – at 13:45

### ORAL HYGIENE IN ORTHODONTIC PATIENTS – A RANDOMIZED CLINICAL TRIAL

**L Golland**<sup>1</sup>, T Imfeld<sup>2</sup>, T Attin<sup>2</sup>, M Schätzle<sup>1</sup>, P R Schmidlin<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and Pediatric Dentistry and <sup>2</sup>Preventive Dentistry, Periodontology and Cariology, University of Zurich, Switzerland

**AIM:** To verify, in a single-blind, randomized clinical trial, the oral hygiene efficacy of different tooth brushes in orthodontic patients. The brushes tested had been chosen after prior *in vitro* tests.

**SUBJECTS AND METHOD:** Thirty-five regular orthodontic patients with a minimum of six brackets bonded to the upper anterior teeth were randomly assigned to three experimental groups of different manual toothbrushes. The best performing *in vitro* tooth brushes with a staged (Parodin plus former Candia Parodin, Migros, 12 patients) or a V-shaped (Oral-B Ortho, Procter & Gamble, 12 patients) brush head design were compared with a planar control brush for cleaning efficacy (Paro M 43, Esro, 11 patients).

The participants were advised to brush their teeth twice daily, for 2 minutes, irrespective of the toothbrush used, for 3 weeks. A standard toothpaste (elmex Junior, Gaba) was provided. Prior to the start and at the end of the study the Gingival Index (GI) was undertaken at the mesial, buccal and distal tooth sites (18 sites per patient) to evaluate the level of oral hygiene.

**RESULTS:** Of the 35 subjects who started the clinical investigation, two (one from the control group and one from the Oral-B cohort) did not attend after 3 weeks and were excluded. At baseline, the control group had a significantly better oral hygiene level (39 sites with GI  $\geq 2$ ) than the two test groups (Parodin 76 sites, Oral-B 70 sites). After 3 weeks of manual tooth brushing with the respective devices, the control group showed no improvement in oral hygiene level (40 sites with GI  $\geq 2$ ), whereas the Parodin-cohort (55 sites with GI  $\geq 2$ ) and the Oral-B group (42 sites with GI  $\geq 2$ ) showed an improvement of 27 to 40 per cent, respectively.

**CONCLUSION:** Participants using staged and V-shaped brush head designs showed a significant improvement in their level of oral hygiene and gingival health.

## SP 196

Wednesday, June 16 – at 13:45

### DOES ORTHODONTIC TREATMENT HAVE ANY EFFECT ON MORPHOPSYCHOLOGY?

**A Karamifar**, M Oshagh, P Salehi, S Momeni Danaei, M M Naseri, Department of Orthodontics, Shiraz Dental School, Iran

**AIM:** The relationship between physical and morphological appearance and psychological characteristics is discussed in morphopsychology. In dentistry, the size and width of the mouth is considered in morphopsychology. A protrusive mouth indicates strong ambition, while a retracted mouth is usually suggestive of being affected by rules. The aim of this study was to evaluate the morphologic aspects of orthodontic treatment.

**MATERIALS AND METHOD:** Pre- and post-treatment frontal photograph of 30 patients treated with standard edgewise and 30 with straight edgewise (MBT) appliances. All had a Class I malocclusion (ANB = 2-4, Wits appraisal = -1-1 and moderate crowding). The proposed treatment plan was extraction of four first premolars. On each photograph the distance between the most superior point of the upper lip and the most inferior point of the lower lip and the inter commissural width were measured using Photoshop software. The ratio between lip thickness and intercommissural width was calculated in order to avoid magnification errors. This ratio was calculated between the pre- and post-treatment photographs for both methods, and the data were statistically analyzed using SPSS software.

**RESULTS:** The mean pre- and post-treatment lip thickness/intercommissural width ratio in patients with standard edgewise appliance was  $0.3830 \pm 0.1191$  and  $0.3125 \pm 0.06668$  respectively, which was statistically significant ( $P \leq 0.001$ ), and the same pre- and post-ratio for the second group was  $0.3313 \pm 0.08753$  and  $0.3009 \pm 0.0559$  which was statistically significant ( $P = 0.019 < 0.05$ ). The mean reduction in the ratio was not statistically significant between either method ( $P = 0.127$ ).

**CONCLUSION:** Extraction of teeth can cause retrusion of the lips. It is suggested that in borderline cases non-extraction treatment should be considered.

## SP 197

Wednesday, June 16 – at 13:45

### BUCCAL CORRIDOR RATIO – DOES IT FOLLOW A RULE?

**A Karamifar**, P Salehi, M Oshagh, M M Naseri, **S Saadat**, Orthodontic Research Centre, Shiraz Dental College, Iran

**AIM:** One of the most important aspects of a 'beautiful' smile is amount of tooth shown during smiling and the negative space created between the teeth and cheeks. The aim of this study was to determine the relationship between the amount of tooth shown during smiling and intercommissural width (buccal corridor ratio) in Class 1 malocclusion patients.

**SUBJECTS AND METHOD:** One hundred and thirteen patients with a Class I malocclusion with crowding (ANB 2-4, Wits = -1 to 1, Class I canine relationship). For each patient three photographs were taken during posed smiling and the photograph that was considered as being more close to a posed smile was selected. The ratio between the amount of tooth displayed and intercomisural width was determined and measured as the buccal corridor ratio using the Photoshop cs2 software. The data were analysed using the Statistical Package for Social Sciences.

**RESULTS:** It seems that in Class I malocclusion subjects the ratio is approximately 1.3.

**CONCLUSION:** Since the amount of tooth shown during smiling is an important compartment of smile aesthetics it should be considered during reconstructive treatment. In Class I malocclusion patients it follows a rule. Buccal corridor ratio can be used as the transverse limitation of tooth positions.

## SP 198

Wednesday, June 16 – at 13:45

### CRANIOFACIAL MORPHOLOGY IN SLEEP APNOEA PATIENTS WITH DIFFERENT CERVICAL COLUMN FUSIONS

**P Svanholt**<sup>1</sup>, L Sonnesen<sup>1</sup>, N Petri<sup>2</sup>, I Kjær<sup>1</sup>, <sup>1</sup>Department of Orthodontics, University of Copenhagen and <sup>2</sup>ENT Department, Nykøbing County Hospital, Denmark

**AIMS:** Scientific investigations have not revealed a definitive aetiological and morphological background for patients with sleep apnoea (OSA), and it has not been possible to subclassify craniofacial phenotypes and relate these to specific aetiological factors. The aims of this study were to subgroup patients with OSA according to cervical column morphology and to analyze craniofacial profiles and head posture in the different groups.

**SUBJECTS AND METHOD:** The OSA group consisted of 16 females and 75 males. Craniofacial profiles and head posture were examined for 74 of the males OSA was diagnosed by overnight polysomnography. The control group comprised 15 females and six males. Lateral profile radiographs in standardized a head posture were taken. Visual assessment of the cervical column and standardized cephalometric analyses was performed. The male OSA group was divided into four groups according to fusions in the cervical vertebrae: group I, no fusions; group II, fusion C2 and C3; group III, occipitalization; group IV, block fusion.

**RESULTS:** In the OSA group 46.2 per cent had fusion anomalies. As 14.3 per cent in the control group had fusion anomalies, significantly more fusion anomalies were seen in the OSA group. In the male OSA group no significant differences in craniofacial morphology and head posture were seen between groups I and III. Between groups I and II a significant difference was seen in jaw relationship and between groups I and IV anterior face height and head posture deviated significantly.

**CONCLUSION:** Morphological deviations of the upper cervical vertebrae are associated with deviations in the craniofacial profile. OSA individuals with block fusions in the cervical vertebrae differed significantly in craniofacial profile and head posture compared with other OSA patients.

## SP 199

Wednesday, June 16 – at 13:45

### IMPACTED MAXILLARY CANINES – AN EPIDEMIOLOGICAL STUDY OF TREATED PATIENTS

**J Cotič**<sup>1</sup>, S Zupančič<sup>2</sup>, M Ovsenik<sup>1</sup>, <sup>1</sup>Department of Orthodontics, Faculty of Medicine, University of Ljubljana and <sup>2</sup>Zobozdravstvo Zupančič d.o.o., Ljubljana, Slovenia

**AIM:** Impacted maxillary canines (IMC) occur in approximately 2 per cent of the population. The condition is thought to be genetic and related to an Angle Class II division 2 malocclusion. Extraction of the primary canine is a possible preventive measure and is most successful in patients under 13 years of age. The aim of this study was to collect epidemiological data regarding IMC in the Slovenian population. Emphasis was placed on the age at the beginning

of orthodontic treatment. A further interest was whether the primary canines had been extracted by a general practitioner and if a correlation exists between age and severity of displacement.

**MATERIALS AND METHOD:** Data from the dental casts and radiographic images of 35 patients (21 girls, 14 boys) with IMC. Two age groups were formed: up to 13 years and older than 13 years. The position of the IMC and the presence of the primary canine were recorded. Dental casts were assessed for crowding with a calliper and classified according to Angle's classification. From panoramic radiographs, among other data, canine angulation to the midline was measured using a digital screen protractor.

**RESULTS:** The mean age at the start of treatment was 15.4 years. Twenty patients (57%) were older than 13 years. Both canines were impacted in five patients (14%). Palatal impaction was present in 28 patients (80%) and buccal impaction in seven patients (20%). Twenty-two patients (63%) still had the primary canines. The occurrence of RII was 46 per cent. Crowding of 4 mm or more was correlated with buccal impaction. Angulation to the midline exceeded 15 degrees significantly more frequently in patients over 13 years of age.

**CONCLUSION:** The majority of patients start treatment for IMC too late; the primary canines are still present. Early intervention is vital because IMC tend to shift mesially in time, complicating treatment and inducing root resorption of adjacent teeth.

## SP 200

Wednesday, June 16 – at 13:45

### FORCE DEGRADATION OF ELASTOMERIC LIGATURES AND ELASTIC SEPARATORS USED AS AN ACTIVE TIEBACK

**F Mahmoudi**<sup>1</sup>, A Mohammadi<sup>2</sup>, Departments of Orthodontics, <sup>1</sup>Tehran Azad University of Dental Science and <sup>2</sup>Tabriz University of Medical Science, Iran

**AIM:** To evaluate the initial force and force decay patterns of two commercially available elastomeric products including elastomeric ligatures and separator elastic used as an active tieback.

**MATERIALS AND METHOD:** Twelve elastomeric ligatures and also elastic separators from each of three manufacturers (Dentaurum, Rocky Mountain, 3M Unitek) were stretched 100 per cent and 12 samples of each brand were stretched 150 per cent of their inner diameter. Force levels were measured initially and at 3 minutes, 24 hours, and 1, 2, 3 and 4 weeks. All samples were placed in a synthetic saliva bath at 37°C. Data was analyzed statistically by univariate analysis of variance and Tukey *post hoc* test.

**RESULTS:** The mean initial forces of the elastomeric ligatures and separators stretched 100 per cent of their inner diameters were 199 to 305 g and 330 to 416 g; respectively, whereas stretched to 150 per cent the loads were 286 to 422 g and 433 to 540 g, respectively. Ten to eighteen per cent of the initial force was lost within the first 3 minutes, with 29-63 per cent of force decay occurring in the first 24 hours. The force decay rate then decreased; however 60-81 per cent of initial force was lost in 4 weeks. Although the force decay pattern was identical for all products, the initial force and force decay of Dentaurum elastomeric ligature and separators was less than similar products from RMO and 3M Unitek ( $P < 0.05$ ). The force decay of separator elastic at different time points was almost similar to the elastomeric ligatures of the same company.

**CONCLUSION:** Regarding the initial force and force decay of elastomeric ligature and separator elastics, they can be used to apply orthodontic force in the form of active tieback.

## SP 201

Wednesday, June 16 – at 13:45

### APPLICATION OF A ER:YAG LASER FOR ORTHODONTIC BONDING. A PRELIMINARY IN VITRO STUDY

**G Di Carlo**<sup>1</sup>, G Ierardo<sup>1</sup>, R Kornblit<sup>1</sup>, P Rocca<sup>2</sup>, A Polimeni<sup>1</sup>, <sup>1</sup>Department of Oral Science, University of Rome 'Sapienza', Italy and <sup>2</sup>University of Nice 'Sophia-Antipolis', France

**AIM:** The use of lasers in orthodontics was, for many years, limited to surgery of the pathological fraenum and the uncovering of

enclosed teeth. Recently, with the introduction of the Er:YAG laser in dentistry for ablation of hard tissue such as enamel and dentine, this wavelength was studied for enamel treatment before orthodontic bracket bonding.

**MATERIALS AND METHOD:** Sixty teeth, extracted for orthodontic reasons, with no decay or previous treatment. The samples were stored in a saline solution at 4°C to avoid dehydration or alteration of the enamel tissue, for a period not exceeding 21 days. The roots were then included in a resin block to hold the specimen and to ease bonding procedure. The teeth were divided into five groups treated with different bonding procedure. Group 1 was prepared with orthophosphoric acid, group 2 were treated with a Er:YAG laser 80 mJ and group 3 with a laser at 80 mJ and orthophosphoric acid. Groups 4 and 5 were treated using same procedures as for groups 2 and 3 but at a lower energy, i.e. 40 mJ.

**RESULTS:** After debonding significant differences were found difference between the five adhesion groups.

**CONCLUSION:** The application of a Er:YAG laser (2940 nm) for bonding should be combined with orthophosphoric acid.

## SP 202

Wednesday, June 16 – at 14:30

### BONDING AND REBONDING OF DIFFERENT CERAMIC BRACKETS

**F Falkensammer**<sup>1</sup>, I Mischak<sup>2</sup>, H-P Bantleon<sup>1</sup>, <sup>1</sup>Medical University, Bernhard Gottlieb University Dental Clinic, Vienna and <sup>2</sup>Medical University, Dental Clinic, Graz, Austria

**AIM:** Ceramic brackets have become an established aesthetic option in orthodontic dentistry. In contrast to mesh designed metal bracket bases, ceramic brackets show different base designs. Little evidence exists about the bond strengths and rebonding capacity of these brackets using diverse reconditioning procedures. The aim of this study was to investigate the bond strengths of three different ceramic brackets as well as the rebond strengths using two different conditioning procedures.

**MATERIALS AND METHOD:** Three different ceramic brackets [Clarity SL (3M), Innovation-C (GAC) and Fascination 2 (Dentaurum)] were bonded on feldspathic ceramic specimens using a standard bonding procedure. Sandblasting or sandblasting in combination with a recently developed ceramic conditioner were included as rebonding procedures. Shear testing was performed with a universal testing machine, followed by stereomicroscope investigation of the debonded brackets.

**RESULTS:** All ceramic brackets (bonded/rebonded) exceeded the minimum bracket bond strength of 6-8 MPa. A significant difference ( $P < 0.001$ ) was found between the three standard bonded brackets. Sandblasting showed varying, but still high bond strengths compared with standard bonding. Adding the ceramic conditioner to the sandblasted brackets increased and significantly ( $P < 0.001$ ) stabilized the bond strengths.

**CONCLUSION:** Bonding ceramic brackets with different designed bracket bases leads to different but high bond strengths. Reconditioning ceramic brackets using the above procedures seems to be appropriate for rebonding.

## SP 203

Wednesday, June 16 – at 14:30

### PROFILE CHANGES AFTER MAXILLARY POSTERIOR IMPACTION

F Sarkarat<sup>1</sup>, A Jamilian<sup>2</sup>, A Karamifar<sup>2</sup>, A Masoudi<sup>2</sup>, T Taban<sup>2</sup>, Department of Orthodontics, <sup>1</sup>Shiraz Dental School and <sup>2</sup>Islamic Azad University, Tehran, Iran

**AIM:** The treatment of long face patients is considered difficult and a large number require orthognathic surgery: The aim of this study was to evaluate and predict profile changes after Le Fort 1 osteotomy, including maxillary posterior impaction.

**SUBJECTS AND METHOD:** Fifteen patients (12 females, 3 males, mean age 24.2 years) with a Class II division 1 malocclusion and an open bite (more than 4 mm). In three patients the second premolars had been extracted while in 12 patients no extractions had

been performed. After bonding with a 0.022 standard edgewise system, levelling, aligning and arch decompensation, all patients underwent a Le Fort 1 osteotomy for maxillary posterior impaction. Pre- and post-operative lateral cephalograms were evaluated, and analyzed by Wilcoxon test.

**RESULTS:** Go-GN/SN and nasolabial angle were decreased  $3 \pm 0.5$  ( $P < 0.01$ ) and  $1.4 \pm 1.6$  ( $P < 0.8$ ) degrees, respectively. The lower lip was shortened  $3.2 \pm 0.1$  mm ( $P < 0.003$ ) and its protrusion was reduced  $1.6 \pm 0.5$  mm ( $P < 0.02$ ). Upper lip length decreased  $0.4 \pm 2$  mm ( $P < 0.3$ ) and its protrusion was decreased  $1.2 \pm 0.2$  mm ( $P < 0.2$ ). Anterior face height decreased,  $5.9 \pm 1.4$  mm ( $P < 0.001$ ).

**CONCLUSION:** After maxillary posterior impaction, anterior face height, GO-GN/SN, lower lip length and lower lip protrusion, were significantly decreased. Upper lip length, nasolabial angle and upper lip protrusion did not change significantly.

## SP 204

Wednesday, June 16 – at 14:30

### ASSESSMENT OF THE PERMANENT DENTITION IN ADULTS WITH AN IDEAL OCCLUSION

**M Ribackova**, L S Persin, T Repina, A Zinchenko, I Popova, Moscow State University of Medicine and Dentistry, Russia

**AIM:** It has been suggested that the permanent dentition should be evaluated using angular parameters Po-N-I and Po-N-M (Persin, 2009). Angle Po-N-I characterizes the position of interincisor point I with regard to the PoN reference line and angle Po-N-M the position of the occlusal contact point (M) between the upper and lower first molars with regard to the PoN reference line. The aim of this study was to evaluate the permanent dentition in adults with an ideal occlusion, using these parameter angles.

**SUBJECTS AND METHOD:** Forty-five adults, aged 19 to 25 years, with an ideal occlusion. Plaster casts were obtained and measured. Angular measurements were obtained from lateral cephalograms. Correlation and statistical analyses were undertaken.

**RESULTS:** The average means of the parameters were  $52.3 \pm 2.6$  degrees (PoNI) and  $76.3 \pm 1.0$  degrees (PoNM). Correlations between parameters, characterized by the sum of the mesiodistal widths of the teeth and angles, PoNI, PoNM were high.

**CONCLUSION:** The use of the parameters Po-N-I and Po-N-M is recommended for assessment of the permanent dentition in adults with an ideal occlusion.

## SP 205

Wednesday, June 16 – at 14:30

### PATHWAYS BETWEEN SUBJECTIVE ORAL HEALTH, FACIAL PAIN, TEMPOROMANDIBULAR DISORDERS AND OCCLUSAL FEATURES

**J Rusanen**<sup>1</sup>, K Sipilä<sup>2,4</sup>, A-S Silvola<sup>1</sup>, P Pirttiniemi<sup>1,4</sup>, S Lahti<sup>3,4</sup>, Departments of <sup>1</sup>Oral Development and Orthodontics, <sup>2</sup>Prosthetic Dentistry and Stomatognathic Physiology and <sup>3</sup>Community Dentistry, Institute of Dentistry, University of Oulu and <sup>4</sup>Oral and Maxillofacial Department, Oulu University Hospital, Finland

**AIM:** To examine the pathways between subjective oral health, facial pain, temporomandibular joint disorders (TMD) and occlusal features in patients with severe malocclusion.

**SUBJECTS AND METHOD:** Ninety-four adults (mean age 38 years, 64% female, 36% male) referred for orthodontic or surgical-orthodontic treatment. All had severe, diagnosed malocclusions with considerable functional disorders. Subjective oral health was measured with the Oral Health Impact Profile-14, facial pain using a visual analogue scale, TMD with Helkimo's clinical dysfunction index and occlusal features with the Peer Assessment Rating Index. A hypothetical model of the interrelationships between these factors was constructed based on the conceptual model of biological, behavioural and psychosocial consequences of oral diseases. The associations were studied with path analysis.

**RESULTS:** In contrast to the hypothetical model, among females the occlusal features were not directly associated with subjective

tive oral health but the association was mediated via facial pain. Among males the occlusal features were directly associated with subjective oral health. Females reported poorer subjective oral health, higher pain levels and had more severe TMD than males, but the gender difference was statistically significant only for pain and TMD.

**CONCLUSION:** TMD, occlusal features, and facial pain are associated with subjective oral health among patients with severe malocclusions, but the association of the occlusal features to oral health differ between genders. Therefore, these associations should be studied separately among genders.

## SP 206

Wednesday, June 16 – at 14:30

### ARRESTED ERUPTION OF THE PERMANENT LOWER SECOND MOLAR

**N Toseska-Spasova<sup>1</sup>**, J Gjorgova<sup>1</sup>, C Misevska<sup>1</sup>, Z Spasov<sup>2</sup>,  
<sup>1</sup>Department of Orthodontics, Faculty of Dentistry, University of Ss. 'Cyril and Methodius' and <sup>2</sup>Private practice 'Prosthetics-S&S', Skopje, Former Yugoslav Republic of Macedonia

**AIM:** To investigate craniofacial morphology, the frequency of dental anomalies, and the inclination of the lower second molar (LM2) and the adjacent first molar in patients with arrested eruption of LM2. The overall goal was to elucidate the aetiology of arrested tooth eruption and to present the characteristics of these patients in order to improve diagnosis and treatment planning.

**MATERIALS AND METHOD:** Radiographs of 20 patients with arrested eruption of LM2, aged 13 to 19 years. Seventeen cephalometric variables were evaluated on lateral cephalograms. The frequencies of dental anomalies of the retained LM2 (crown and root morphology) and of the inclination of LM2 were evaluated on dental pantomograms.

**RESULTS:** Patients with arrested eruption of LM2 had an increased sagittal jaw relationship. The mandibular prognathism was less, the mandibular gonial angle smaller, the mandibular alveolar prognathism enlarged and the maxillary incisor inclination less than in the reference group. This group of patients had a more frequent occurrence of morphological tooth anomalies, such as root deflections, invaginations and taurodontism. There was no agenesis of the lower third molar. There was no association between the degree of inclination of the LM2 and that of the first molar in the same region.

**CONCLUSION:** Craniofacial morphology and deviations in the dentition are associated with arrested eruption of LM2. It is important to evaluate these conditions in future diagnosis and treatment planning of patients with arrested eruption of LM2.

## SP 207

Wednesday, June 16 – at 14:30

### INCIDENCE OF OROFACIAL CLEFTS IN THE CZECH REPUBLIC FROM 2000-2006

**W Urbanova<sup>1</sup>**, M Kotova<sup>1</sup>, Z Vankova<sup>2</sup>, <sup>1</sup>Department of Orthodontics and Cleft Anomalies, Stomatological Clinic 3rd Medical Faculty and <sup>2</sup>2nd Faculty of Medicine Charles University, Prague, Czech Republic

**AIM:** Orofacial clefts are the most common craniofacial congenital birth defect. There is a wide variation in incidence of various cleft anomalies between different populations and genders. The aim of this study was to determine an incidence of orofacial cleft defects in the Czech Republic among infants born during 2000-2006.

**MATERIALS AND METHOD:** Data were collected from the Institute of Health Information and Statistics, Czech Statistical Office and Czech Health Statistics Yearbooks. The incidence of orofacial cleft anomalies and gender differences in live born infants were calculated.

**RESULTS:** There were a total of 1086 infants with an orofacial cleft among 673,802 of newborns between 2000-2006. The incidence of orofacial cleft defects was on average 1.61 per 1000 live births.

The incidence of a cleft lip was 0.38, of a cleft palate (CP) 0.67 and of a cleft lip and palate (CLP) 0.55 per 1000 live born infants. There were almost twice as many boys with a cleft lip (169 boys, 90 girls, incidence 0.49 and 0.28, respectively per 1000 live births) and CLP in girls (248 boys, 126 girls, incidence 0.71 and 0.39, respectively per 1000 live births). The overall number of boys and girls with a CP was 231 and 222, respectively (incidence 0.67 in boys and 0.68 in girls per 1000 live births).

**CONCLUSION:** The incidence of orofacial clefts in the Czech Republic among infants born during 2000-2006 was lower than that reported in previous studies (1.74 per 1000 newborns) from 1962-1996, oscillating with small deviations around 1.61 per 1000 live born infants. Boys are affected with orofacial clefts more frequently than girls, with the exception of CP patients, where both genders are equally affected. Despite the rising birth rate in the Czech Republic during 2000-2006, the incidence of orofacial clefts in the time period remained stable.

## SP 208

Wednesday, June 16 – at 14:30

### EFFECTS OF FLUORIDE-CONTAINING CASEIN PHOSPHOPEPTIDE-AMORPHOUS CALCIUM PHOSPHATE ON SHEAR BOND STRENGTH OF ORTHODONTIC BRACKETS

**S B Çehreli<sup>1</sup>**, Ç Sar<sup>2</sup>, Ö Polat-Özsoy<sup>2</sup>, B Ünver<sup>3</sup>, S Özsoy<sup>4</sup>,  
Departments of <sup>1</sup>Pediatric Dentistry and <sup>2</sup>Orthodontics, Başkent University, <sup>3</sup>Department of Mining and Engineering, Hacettepe University, and <sup>4</sup>Ankara Topraklık Ministry of Health Dental Hospital, Turkey

**AIM:** To investigate the effect of enamel pre-treatment with a new 900 ppm fluoride-containing casein phosphopeptide-amorphous calcium phosphate (CPP-ACP) complex on the shear bond strength (SBS) of brackets bonded with etch-and-rinse or self-etching adhesive systems.

**MATERIALS AND METHOD:** Extracted human premolars were randomly divided into six groups (n = 11/group) with respect to the enamel pre-treatments and adhesive systems employed. Group I, No pre-treatment was performed on the enamel and the brackets were bonded with an etch-and-rinse adhesive system (Transbond XT). Group II, the enamel was pre-treated with a fluoride-containing CPP-ACP paste (MI Paste-Plus) and the brackets bonded with Transbond XT. Group III, the enamel was pre-treated with a non-fluoride CPP-ACP paste (MI Paste) and the brackets bonded with Transbond XT. Group IV, no pre-treatment was performed and the brackets were bonded with a self-etching adhesive system (Transbond-Plus). Groups V and VI, the enamel was pre-treated as for groups II and III, respectively, and the brackets bonded with Transbond Plus. Bonded specimens were subjected to thermocycling (X1000) and 6 weeks water degradation before SBS testing. The residual adhesive on the enamel surface was evaluated after debonding with the adhesive remnant index (ARI).

**RESULTS:** Enamel pre-treatment with either the fluoride or non-fluoride CPP-ACP paste had no significant effect on the SBS of the self-etching adhesive system ( $P > 0.05$ ). Enamel pre-treatment with non-fluoride CPP-ACP significantly reduced the SBS of the etch-and-rinse adhesive ( $P < 0.001$ ), while pre-treatment with the fluoride-containing CPP-ACP paste did not affect debonding values ( $P > 0.05$ ).

**CONCLUSION:** The fluoride-containing CPP-ACP did not compromise the bond strength of brackets bonded with the tested etch-and-rinse and self-etching systems, but its non-fluoride version significantly decreased the SBS of the etch-and-rinse adhesive system.

### QUALITY AND VARIATION IN ORTHODONTIC POSTGRADUATE EDUCATION IN EUROPE

**E Santigli**<sup>1</sup>, L Bolamperti<sup>2</sup>, O Annousaki-Thormann<sup>2</sup>, M G Laursen<sup>4</sup>, <sup>1</sup>Department of Orthodontics and Maxillofacial Orthopedics, Medical University Graz, Austria, <sup>2</sup>Department of Orthodontics, University of Insubria, Italy, <sup>3</sup>Switzerland and <sup>4</sup>Department of Orthodontics, University of Aarhus, Denmark

**AIM:** Clinical excellence in orthodontics begins with quality in orthodontic education. At present, postgraduate education in orthodontics varies throughout Europe both in duration and content, hence presumably in quality. The purpose of the study was to identify existing disparities in European orthodontic postgraduate education.

**MATERIALS AND METHOD:** An inquiry was conducted during two European Postgraduate Student Orthodontic Society meetings in 2008 and 2009. One hundred and fifteen postgraduate students volunteered to answer a comprehensive questionnaire concerning their current education including admission criteria, clinical training, theoretical education and terms of completion. Descriptive statistics were applied to the data.

**RESULTS:** Thirty-eight institutions in 23 countries were represented. The number of residents in a department ranged from 3 to 100. The education was full-time for 93 per cent of the students. The terms of completion included a thesis for 83.3 per cent, and a final examination for 81.6 per cent. The duration of the programmes ranged between 2 and 5 years. The number of started patients ranged from 4 to 200 and the number of finished cases from 0 to 198. Chair time as the first operator was reported by 94.8 per cent. Of these, 1.7 per cent had less than 5 hours/week and 67.7 per cent more than 20 hours/week as first operator. Mandatory theoretical lesions were reported by 90.3 per cent of respondents. Of these 8.1 per cent had less than 2 hours/week and 20 per cent more than 20 hours/week.

**CONCLUSION:** The data revealed inconsistencies with respect to hours of presence, chair time, number of patients, theoretical lessons and terms of completion, indicating substantial disparities in orthodontic training for European postgraduate students. The study serves as a starting point for further investigations, with the ultimate aim to promote standardization and overall quality improvement of postgraduate orthodontic education in Europe.

### ORTHODONTIC FEATURES IN CEREBRAL PALSY

**E Pozzani**, S Di Raffaele, F Tezza, Special Care Dentistry Department, Verona, Italy

**AIM:** Cerebral palsy (CP) refers to anyone of a number of neurological disorders that appear in infancy or early childhood and permanently affect body movement and muscle coordination. The aim of this study was to evaluate the relationship between cerebral palsy and orthodontic defects and the presence of atypical swallowing.

**MATERIALS AND METHOD:** The clinical records of 36 patients with CP aged from 2 to 24 years. The data were compared with those of a randomly selected control group (CG). The clinical parameters investigated were dental Class, crossbite, overbite, dental crowding and atypical swallowing.

**RESULTS:** 1) Class I: CP 27.8 per cent, CG 55.6 per cent; Class II: CP 72.2 per cent, CG 30.6 per cent; Class III: CP 0 per cent, CG 13.9 per cent. 2) Crossbite: CP 8.3 per cent, CG 30.6 per cent; 3) Normal overbite: CP 55.6 per cent, CG 77.8 per cent; 4) Open bite: CP 38.9 per cent, CG 11.1 per cent; 5) Deep bite: CP 5.6 per cent, CG 11.1 per cent; 6) Dental crowding: CP 13.9 per cent, CG 30.6 per cent; 7) Atypical swallowing: CP 58.2 per cent, CG 27.8 per cent.

**CONCLUSION:** CP patients must have a dental check-up starting from an early age in order to control the risk of developing a malocclusion; most of all in relation to the high incidence of Class II malocclusions and skeletal open bites.

### INFLUENCE OF ORTHODONTIC TREATMENT ON CANDIDA COLONISATION IN THE ORAL CAVITY

**A Wojtaszek-Słomińska**<sup>1</sup>, A Kędzia<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Oral Microbiology, Medical University of Gdansk, Poland

**AIM:** Fixed appliances make oral hygiene care difficult by creating new retention sites where dental plaque and tartar accumulate. As microbiologic studies have shown, the sites especially permitting adhesion of bacteria and yeast are orthodontic brackets, bands and adhesive. The aim of this research was to examine how orthodontic treatment with fixed appliances affects the frequency of candida occurrence in the oral cavity of adults.

**SUBJECTS AND METHOD:** One hundred and fourteen adult patients (85 females, 29 males) divided into two groups. In group I (62 subjects) the material for microbiological examination was collected three times: before treatment and one and six months after the start of treatment and in group II (52 patients) after 18 months of treatment. The material was collected from four sites in the oral cavity: the buccal surface of tooth 16, the labial surface of tooth 21, the buccal surface of tooth 36 and the dorsal surface of the tongue. The material was obtained with the use of a disposable sterile matrix. A sterile swab moistened in 0.9 per cent salt solution was used and delivered to a microbiology laboratory within 1 hour.

**RESULTS:** In group I, 55 candida strains were cultured in 24 (38.7%) patients before treatment, 57 strains in 23 subjects (37.1%) at one month, while 87 candida strains were cultured in 32 patients (51.6%) at six months. In group II, 120 candida strains were found in 39 (75%) of patients. The most frequently observed species were *Candida albicans* (52.97%).

**CONCLUSION:** The number of candida species in the oral cavity of patients treated with fixed appliances increases over treatment time. The most frequently found candida species was *C. albicans*.

### ASSESSMENT OF STAINING OF ORTHODONTIC AESTHETIC BRACKETS DUE TO FOOD COLOURANTS

**K Malkiewicz**<sup>1</sup>, E Malkiewicz<sup>2</sup>, <sup>1</sup>Department of Conservative Dentistry, Medical University of Warsaw, and <sup>2</sup>Department of Public Health, College of Finance and Management, Siedlce, Poland

**AIM:** To assess if food dyes significantly affect the colour of different brands of orthodontic brackets, both composite and ceramic.

**MATERIALS AND METHOD:** Eight brands of orthodontic brackets, four composite: Rave, Crystal Clear Plastic Bracket, Silkon Plus, Orthoflex, and four ceramic: Reflections, Pure, Contour and Miso were investigated. Ten samples of each bracket were assessed. Five brackets of each series were stored in coffee and five in red wine for 24 hours. The colour change after storage was analyzed for each sample with the use of a dental spectrophotometer according to the CIE L\*a\*b\* colour scale. The lightness, redness, yellowness and complete colour change ( $\Delta E$ ) of the brackets were statistically analyzed at the level of  $P = 0.05$ .

**RESULTS:** The greatest colour change ( $\Delta E$ ) was observed in composite brackets (mean level 13.84 points) compared with porcelain brackets (mean level 3.02 points). Various colour changes were observed both among composite brands (mean  $\Delta E$  ranged from 6.94 to 18.19) and among the porcelain group (mean  $\Delta E$  ranged from 2.11 to 3.78).

**CONCLUSION:** Food colourants cause significant colour changes of both composite and ceramic orthodontic brackets. Composite orthodontic brackets are significantly more susceptible to food discolouration than ceramic brackets.

**THREE-DIMENSIONAL STUDY OF THE DUMBO RAT**

**M Charezinski**<sup>1</sup>, N Vanmuylder<sup>2</sup>, S Van Sint Jan<sup>2</sup>, R Glineur<sup>1</sup>, S Louryan<sup>2</sup>, <sup>1</sup>Clinique D'Orthodontie - Hopital Erasme and <sup>2</sup>Laboratoire d'Anatomie et d'Embryologie, Faculte de Medecine, Universite Libre de Bruxelles, Belgium

**AIM:** A variation of the domestic rat, the dumbo rat, has similar traits to human craniofacial disorders such as Treacher Collin syndrome or other pharyngeal arch disorders. The dumbo rat's malpositioned ears and retrognathic aspect strongly suggest a defect in the development of the first and second pharyngeal arch. Possible causes of such abnormal development may be faulty migration of neural crest cells, a genetic mutation (e.g. TCOF1) or external factors such as teratogens (e.g. retinoids). Beside an embryologic and genetic approach focused on possible causes, the anatomical traits of the dumbo rat need to be precisely defined. The aim of this three-dimensional (3D) cephalometric analysis was to validate a new 3D rat analysis and to define more precisely the anatomical malformations of the dumbo rat in comparison with the Wistar rat.

**MATERIALS AND METHOD:** Amira 4.0 software was first used to analyze the computed tomographic scan data (dicom files) and to create a 3D model of the head of the rat. The second phase, with LHPBuilder software, consisted of anatomical landmarks on the 3D model. Numerous linear and angular measurements were inferred from these landmarks. 3D cephalometric results obtained from a sample of dumbo rats were compared with a control group of Wistar rats.

**CONCLUSION:** The 3D study of dumbo rat could be a first step in using this rodent as an animal model for better understanding human pharyngeal arch disorders.

**AN IN VITRO STUDY OF FRICTION OF CONVENTIONAL LIGATURES WITH SELF-LIGATING BRACKETS**

**M R Ricchiuti**, F Ballanti, C Pavoni, Department of Orthodontics University of Rome, Italy

**AIM:** The overall resistance to sliding in orthodontic appliances is a combination of classic friction, archwire-bracket binding, and archwire notching. The aim of this study was to compare the frictional force generated by self-ligating (SL) and standard (S) brackets coupled with stainless steel (SS) wires when conventional elastomeric (E) or SS ligatures (L) were used.

**MATERIALS AND METHOD:** Four types of brackets were selected, one standard (Standard Boston, Leone), one passive SL (Damon 3MX, Ormco) and two active SL (InOvation R, GAC; Time 3, American Orthodontics). For each type of bracket one molar tube and two upper premolars were used in combination with three different sizes of wire (0.016 X 0.022, 0.017 X 0.025 and 0.019 X 0.025 inch SS). Testing was performed with an Instron 3344 machine on brackets bonded with an epoxy adhesive to a Perspex block. Each bracket/wire combination was tested with EL and SSL. Tests performed with SL brackets were also carried out without conventional ligatures. ANOVA and Tukey tests were used to analyze the results for the different archwire-bracket-ligature combinations.

**RESULTS:** The lowest friction values were registered for the SL method. No significant differences were found in active SL bracket/0.016 X 0.022/EL or SSL assembly when compared with S bracket/0.016 X 0.022/EL combination. Active SL bracket/0.017 X 0.025 or 0.019 X 0.025/SSL showed significantly higher frictional force values than the S bracket for the same combinations. The passive SL bracket showed significantly lower friction values than the S bracket for each archwire/ligature assembly.

**CONCLUSION:** The use of SSL on active SL brackets produced a significantly higher level of frictional force than in combination with the S bracket for the 0.017 X 0.025 and 0.019 X 0.025 archwire. Standard ligatures in combination with SL brackets could be used to obtain anchorage in the dental arch.

**SELF-LIGATING VERSUS CLEAR ALIGNERS ANALYSIS OF DENTOALVEOLAR EFFECTS**

**G Laganà**, M Mucedero, M R Ricchiuti, Department of Orthodontics University of Rome Tor Vergata, Italy

**AIM:** Changes in the transverse measurements of the maxillary arch have an impact on arch perimeter with the consequent space created to accommodate teeth in non-extraction treatment protocols. The aim of this study was to evaluate the changes in transverse dimensions and maxillary arch perimeter produced by low friction self-ligating brackets (Time 3) compared with the Invisalign technique.

**SUBJECTS AND METHOD:** Both the self-ligating and Invisalign groups comprised 20 subjects, evaluated at the beginning (T0) and completion (T1) of therapy. All had a Class I malocclusion with mild crowding in the permanent dentition, without craniofacial anomalies, missing teeth or a history of orthodontic treatment. The following measurements were made on the maxillary dental casts at T0 and T1: intercanine width between the tips of the cusp (CWC) and the most lingual point on the lingual surface (CWL), first interpremolar width between the central fossae (FPWF) and the most lingual point on the lingual surface (FPWL), second interpremolar width between the central fossae (SPWF) and the most lingual point on the lingual surface (SPWL), intermolar width between the mesial fossae on the occlusal surface of the maxillary first molars (MWF) and the lingual fissure location on the lingual surface (MWL), arch perimeter (AP) and arch depth (AD). Significant differences between the treated groups were assessed with an independent samples t-test ( $P < 0.05$ ).

**RESULTS:** Statistically significant differences between the self-ligating and Invisalign group were recorded for CWC, FPWF, FPWL, SPWF, SPWL, and AP measurements. No significant changes were found for CWL, MWF, MWL, and AD values.

**CONCLUSION:** The low friction self-ligating system produced statistically significant different outcomes in transverse dentoalveolar width and maxillary arch perimeter when compared with the Invisalign technique.

**THE SUSCEPTIBILITY TO STAINING OF ORTHODONTIC ELASTICS LIGATURES**

**K Malkiewicz**<sup>1</sup>, B Pietrzak-Bilinska<sup>2</sup>, Departments of <sup>1</sup>Conservative Dentistry and <sup>2</sup>Orthodontics, Medical University of Warsaw, Poland

**AIM:** To assess if food dyes significantly affect the colour of four different brands of orthodontic elastic ligatures.

**MATERIALS AND METHOD:** Eight brands of elastic ligatures: Ortho Organizers, Dentaurum, Ortho Technology and Dentech Corp., divided according to four colours: transparent, white, yellow and silver, were investigated. Twenty samples from each manufacturer were stored in coffee and 20 in red wine for 24 hours. The colour change after storage was analyzed for each sample with the use of a dental spectrophotometer according to the CIE L\*a\*b\* colour scale. The complete colour change ( $\Delta E$ ) of examined ligatures was statistically analyzed at the level of  $P = 0.05$ .

**RESULTS:** The greatest colour change ( $\Delta E$ ) was observed for Ortho Organizers elastics (mean level of 12.81 points) and the lowest for Ortho Technology elastics (mean level 10.93 points). No significant differences among the manufacturers were observed. Significant differences in colour change caused by food dyes were observed in relation to the original colour of the elastics. The greatest colour change was observed for white ligatures (mean level 20.70 points) and the lowest for silver products (mean level 4.91 points).

**CONCLUSION:** Susceptibility to discolouration of elastic ligatures did not depend on the brand under the conditions of this study. The level of discolouration of orthodontic elastic by food dyes is strongly related to the original colour of the material.

**SP 217**

Thursday, June 17 – at 12:15

**PHOTOGRAMMETRIC PROFILE TREATMENT CHANGES OF CLASS II DIVISION 1 MALOCCLUSIONS**

**A Moullas**, A Athanasiou, G Kolokithas, S Sidiropoulou-Chatziagianni, Department of Orthodontics, School of Dentistry, Aristotle University of Thessaloniki, Greece

**AIM:** To assess soft tissue profile changes associated with various orthodontic treatment protocols in growing individuals presenting a Class II division 1 malocclusion.

**MATERIALS AND METHOD:** Initial and final, coloured, profile, standardized photographs taken in the natural head position of 110 Class II division 1 growing patients were analyzed by means of 15 angular and one ratio measurement. Different orthodontic treatment protocols were used: one phase/two extractions ( $n = 21$ ), one phase/four extractions ( $n = 14$ ), one phase/no extractions ( $n = 50$ ) and two phase/no-extractions ( $n = 25$ ). Statistical analysis included paired samples t-tests for each pair of observations in each group as well as assessment of the magnitude of the method error using Dahlberg's formula.

**RESULTS:** Statistically significant ( $P < 0.05$ ) findings included: reduction of facial convexity as well as increase of nasolabial angle, angle of the lips and mentolabial angle, in all four groups. No significant change was observed in the ratio of upper to lower face height in any group. Nasofrontal and cervicomental measurements showed that they constitute parts of the dentofacial system, which were not significantly influenced by the treatment protocols performed.

**CONCLUSION:** Soft tissue profile was improved by the orthodontic treatment in all groups regardless of the specific therapeutic approach used. Angular photogrammetric profile analysis can be a useful mean for assessing the various soft tissue profile regions of the face.

**SP 218**

Thursday, June 17 – at 12:15

**FACIAL PROPORTIONS IN SERBIAN FEMALE ADULTS**

**N Nedeljković**, J Milutinović, Z Stamenković, Department of Orthodontics, University of Belgrade, Serbia

**AIM:** Facial attractiveness is important in interhuman communication because beauty means social power and success and has a positive influence in all areas of civilized society. The aim of this study was to compare vertical and horizontal proportions of the face – frontal view, with golden proportions in adult Serbian females.

**MATERIALS AND METHOD:** Facial photographs in the frontal view of 120 female dental students. The *en face* photograph of each subject was assessed using horizontal thirds and vertical fifths. Four transverse and six vertical facial distances, were measured and compared with the corresponding calculated divine distances. The relationship between the measurements on the photographs and golden facial proportions was evaluated.

**RESULTS:** A significant difference between the measured photographs and golden proportions was found for both lower facial third distances and the lateral fifths of the face.

**CONCLUSION:** The main inequality from the golden proportions was in the lower facial third, as the most impressive attribute of facial attractiveness.

**SP 219**

Thursday, June 17 – at 12:15

**IS A SECULAR TREND IN CRANIAL FORMATION AN EXPLANATION OF THE INCREASE IN MALOCCLUSIONS?**

**M Svabova**<sup>1</sup>, J Racek<sup>1</sup>, I Chourova<sup>2</sup>, <sup>1</sup>Charles University in Prague, First Faculty of Medicine and <sup>2</sup>Private practice, Strazne, Czech Republic

**AIM:** The increasing prevalence of malocclusions is characteristic for recent populations. The aim of this study was to compare craniofacial and dental characteristics of contemporary and his-

torical populations and elucidate some aetiological aspects of malocclusion.

**SUBJECTS AND METHOD:** A contemporary cohort of 703 university students and three historical samples (73 skulls from the 9th, 344 skulls from 10th-14th and 210 skulls from 14th-18th centuries) were examined. Measurements of craniometric and anthropometric points were undertaken. The width of the jaws was examined in Pont's points. Björk's method for epidemiological registration of malocclusion was used and teleroentgenograms were examined.

**RESULTS:** Broader dental arches, regardless of the type of skull, and a significantly lower frequency of serious malocclusions were found in the historical population.

**CONCLUSION:** The increase in severe malocclusions in the contemporary population is most probably caused by civilising factors than by secular trends in the formation of skull.

**SP 220**

Thursday, June 17 – at 12:15

**PERCEPTION OF PAIN DURING ORTHODONTIC TREATMENT**

**J Petrovska**, D Petrovski, Department of Fixed Prosthodontics, Dental clinic PHO 'Sv. Pantelejmon', University 'Sv.Kiril i Metodij', Skopje, Former Yugoslav Republic of Macedonia

**AIM:** To investigate the intensity, location and duration of patient discomfort, following insertion of orthodontic appliances.

**SUBJECTS AND METHOD:** Thirty-two females and 25 males (mean age: 15.5 years) treated with fixed appliances. They were given oral and written instructions on how to complete a questionnaire, which consisted of eight questions. The patient was requested to respond to each question by placing a mark along a 10 cm long visual analogue scale. The questionnaire was to be filled out 4 hours after the appointment and then at 24 hourly intervals for the next 7 days.

**RESULTS:** The majority of patients experienced pain 4 hours after archwire placement, which peaked at 24 hours and then declined. The pain usually lasted 2-3 days and gradually decreased in its intensity by day 5 or 6. Patients reported more pain experience in the anterior than in the posterior teeth, because of the differences in root surface area. The overall findings indicate that analgesics are still the main treatment modality to reduce orthodontic pain. There were no differences between females and males in the evaluation of their facial and dental appearance. The influence of pain on diet was significant.

**SP 221**

Thursday, June 17 – at 12:15

**FORCE SYSTEM OF REVERSE CURVE NICKEL TITANIUM WIRES IN CONVENTIONAL AND SELF-LIGATING BRACKETS**

**I Sifakakis**<sup>1</sup>, N Pandis<sup>2</sup>, M Makou<sup>1</sup>, T Eliades<sup>3</sup>, C Bouraue<sup>4</sup>, <sup>1</sup>University of Athens, <sup>2</sup>Private practice, Corfu, <sup>3</sup>University of Thessaloniki, Greece and <sup>4</sup>University of Bonn, Germany

**AIM:** To comparatively evaluate the intrusive forces and buccolingual moments generated at the maxillary incisors by a 0.017 X 0.025 inch reverse curve nickel titanium (NiTi) wire using self-ligating and conventional appliances.

**MATERIALS AND METHOD:** Ten 0.017 X 0.025 inch reverse curve NiTi wire specimens were used with each of the following 0.022 inch bracket systems: Titanium Orthos™, In-Ovation R™ and Damon System 3MX™. The wires were inserted on bracketed maxillary Frasco models, segmented mesially to the canines. Simulated intrusion from 0.0-1.0 mm was performed on the Orthodontic Measurement and Simulation System, which recorded the intrusive forces and the buccolingual moments at 0.05 mm increments. The data were analyzed by means of ANOVA and Scheffe test.

**RESULTS:** The intrusive forces were significantly different between all bracket types. The highest force was recorded with the conventional brackets (8.2 N), followed by the Damon 3MX™ (6.3 N) and the In-Ovation R™ (5.5 N) system. The moments were found to be significantly different between the conventional and

self-ligating brackets but not between the two types of self-ligating brackets. The highest moments were recorded with the self-ligating brackets (16.6-16.9 Nmm), followed by the conventional brackets (10.8 Nmm).

**CONCLUSION:** The intrusive forces exerted on the incisors by the 0.017 X 0.025 inch reverse curve NiTi wire specimens in unlevelled dental arches were beyond biologically safe limits. Lower intrusive forces but higher buccolingual moments were recorded with the self-ligating brackets in comparison with the conventional brackets. These findings could be attributed to the different ligation methods as well as to the width differences between the various bracket types.

## SP 222

Thursday, June 17 – at 12:15

### DEVELOPMENTAL ANALYSIS OF MANDIBULAR FOSSA CIRCUMFERENCE EMPLOYING SAGITTAL COMPUTED TOMOGRAPHIC IMAGES\*\*\*

**R Oyake**, Y Shinohara, N Koizumi, School of Dental Radiology, Meikai University, Chiba, Japan

**AIM:** To assess the relationship between mandibular fossa circumference developmental stage and occlusal development.

**MATERIALS AND METHOD:** Dried human skulls served as the experimental material; five samples were selected from each stage and a total of 50 samples were examined. Computed tomographic (CT) imaging of the mandibular fossa circumference was performed with the Somatom Emotion 6 (Siemens). Imaging conditions were as follows: X-ray tube voltage 130 kV, tube current 63 mA in bone mode, scan duration 2.7 seconds and slicing interval 2 mm. The images of the sagittal view at the midline and the axial view at the FH plane were acquired at WW3000;WL480. Analyzed images were transferred to the high-speed three-dimensional analysis device, Virtual Place Advance Plus (AZE Ltd), in order to construct images for measurement. The area and growth rate of the sagittal image and the area of the mandibular fossa circumference were then measured on the images and evaluated.

**RESULTS:** 1. The area of the right tuberculum articulare sagittal image at stage IA, which was approximately 0.09 cm<sup>2</sup>, increased to approximately 0.5 cm<sup>2</sup> by stage VA. The growth rate at stage VA was approximately 5.6-fold based on stage IA. The area of the left tuberculum articulare sagittal image at stage IA, which was about 0.068 cm<sup>2</sup>, increased to approximately 0.46 cm<sup>2</sup> by stage VA. The growth rate at stage VA was approximately 6.8-fold based on stage IA. 2. The area of the right aspect of the posterior articulare sagittal image at stage IA, which was about 0.054 cm<sup>2</sup>, increased to approximately 0.258 cm<sup>2</sup> by stage VA. The growth rate at stage VA was approximately 5.5-fold based on stage IA. The area of the left tuberculum articulare sagittal image at stage IA, which was about 0.054 cm<sup>2</sup>, increased to approximately 0.258 cm<sup>2</sup> by stage VA. The growth rate at stage VA was approximately 4.8-fold based on stage IA.

## SP 223

Thursday, June 17 – at 12:15

### HUMAN PRIMARY TEETH REVEAL THE INFLUENCE OF EPITHELIAL CELLS AND INNERVATION IN RESORPTION

**M-L B Bille**, I Kjær, Department of Orthodontics, School of Dentistry, University of Copenhagen, Denmark

**AIM:** Dentitions with ectodermal characteristics are predisposed to root resorption (Kjær, 1995). Studies indicate that the ectodermal derived epithelial cells of Malassez protect against root resorption (Brice *et al.*, 1991; Lambrechts *et al.*, 1993; Kat *et al.*, 2003; Kjær and Nolting, 2009). A connection between resorption in primary teeth and permanent teeth indicates an inborn aetiology (Bille *et al.*, 2008). The aim of this study was to describe the epithelial cells of Malassez and innervation in three different root surface areas without resorption, areas with resorption, and areas with repair after resorption.

**MATERIALS AND METHOD:** Paraffin sections of 13 primary molars and six primary canines from four boys and six girls aged 8-15 years. Sections were stained using REAL TMEnvision™ Detection System (K5007, Dako). Wide spectrum screening (Z0622, Dako) diluted 1:5000 in antibody diluent (S2022, Dako) was used for immunohistochemical detection of epithelial cells. Neuronal nuclei (MAB377, Chemicon/Millipore) diluted 1:50 (S2022, Dako) was used for detection of nerves.

**RESULTS:** The epithelial cells of Malassez were scattered in small islands along root surfaces without resorption and along root surfaces with repair after resorption, but epithelial cells were not present in areas with resorption. Innervation was seen along all three root surface areas and also within the epithelial cells of Malassez.

**CONCLUSION:** Human primary teeth reveal that the epithelial cells of Malassez and innervation may play a role in root resorption.

## SP 224

Thursday, June 17 – at 12:15

### DENTOSKELETAL CHARACTERISTICS IN SUBJECTS WITH ORTHODONTIC TREATMENT NEED IN TOOTH AGENESIS CASES

**G Laganà**<sup>1</sup>, C C Lombardi<sup>1</sup>, A Marino<sup>2</sup>, Department of Orthodontics, University of Rome Tor Vergata, and <sup>2</sup>University of Sapienza Rome, Italy

**AIM:** Tooth agenesis, the congenital absence of at least one tooth, is the most frequent dental anomaly with a prevalence in the permanent dentition of 1.0-9.6 per cent in Caucasians, 6.6 per cent in Japanese, and 7.7 per cent in the Black population. The female-to-male ratio is 3:2 for Caucasians and 2:1 for blacks. In the primary dentition, hypodontia occurs less often (under 1%) and has no significant gender distribution difference. The aim of this study was to analyse skeletal and dental and arch morphology in subjects with dental agenesis compared with a control group.

**MATERIALS AND METHOD:** The records of 84 patients with agenesis and an orthodontic treatment need and a control group of 84 subjects with a complete dentition. Skeletal, dental, morphological and aesthetic parameters were studied on latero-lateral and postero-anterior telerradiographs and dental casts.

**RESULTS:** The teeth most often missing were the maxillary lateral incisors (72.61%) and lower second premolars. The female-to-male ratio was 5:3. Transverse and sagittal measurements showed a significant decrease in the size of the maxilla in the agenesis group. No changes were observed in the mandible.

**CONCLUSION:** Consideration should be given to the altered craniofacial dimensions when planning treatment for subjects with agenesis.

## SP 225

Thursday, June 17 – at 12:15

### IN VITRO INVESTIGATION OF THE TENSILE BOND STRENGTH OF THREE DIFFERENT BRACKET TYPES

**R J Speel**, C J Kleverlaan, T J Algera, Academic Centre for Dentistry Amsterdam (ACTA), Netherlands

**AIM:** To compare, in vitro, the tensile bond strength (TBS) and debonding characteristics [Adhesive Remnant Index (ARI) scores] of metal, ceramic, and polycarbonate brackets.

**MATERIALS AND METHOD:** Forty-five freshly extracted bovine teeth, divided into three equal groups, were ground flat (grit 600) and used as substrate for bracket bonding. Mini Twin metal brackets (Ormco Corporation), Allure ceramic brackets (Dentsply GAC International), and Elegance polycarbonate brackets (Dentaurum) were bonded according to the manufacturers' instructions. After 30 second etching with 37 per cent phosphoric acid, the enamel was washed for 20 seconds and dried for 10 seconds. Brackets were bonded with the Padlock bonding system (Reliance) and light cured for 10 seconds with a visible curing unit (Satelec). After 1 hour the specimens were debonded in a tensile mode using a universal testing machine (Instron) with a crosshead speed of

1 mm/minute. ARI scores were assessed after debonding with a X25 magnifying microscope.

**RESULTS:** Significant differences ( $P < 0.01$ ) in TSB were found between metal ( $4.8 \pm 0.6$  MPa), ceramic ( $9.2 \pm 1.9$  MPa) and polycarbonate ( $3.6 \pm 1.3$  MPa) brackets. Lower ARI scores were found for polycarbonate brackets than for the other groups ( $P < 0.01$ ).

**CONCLUSION:** Higher forces are needed for debonding ceramic brackets than for metal or polycarbonate brackets. Polycarbonate brackets showed significantly less residual resin after debonding than metal or ceramic brackets. Bracket material and geometry appear to play a significant role in debonding characteristics. Changing these properties may improve bonding properties.

## SP 226

Thursday, June 17 - at 13:00

### CHANGES IN ARCH FORM AND DENTITION WITH DIFFERENT RETAINERS AFTER ORTHODONTIC TREATMENT

**S-S An, Y J Choi, C Chung, K-H Kim,** Department of Orthodontics, College of Dentistry, Gangnam Severance Dental Hospital, Seoul, Korea

**AIM:** To evaluate the long-term change in the dentition and dental arch form 5 years after fixed orthodontic treatment using different retention devices.

**MATERIALS AND METHOD:** In group 1 (fixed retainer group) a multistranded wire retainer was bonded to the lingual surface of the anterior segment with removable type retainers fitted after debonding ( $n = 54$ ). In group 2 (removable retainer group) a removable type retainer (circumferential type) was fitted after debonding ( $n = 32$ ). Study models were taken before treatment, after debonding, and 5 years after treatment. Changes in intercanine width, intermolar width, arch length and irregularity index were measured.

**RESULTS:** The irregularity index of the upper and lower incisors significantly increased in both groups, but the amount of increase was significantly greater in the removable retainer group ( $P < 0.05$ ). Intercanine width and arch length significantly decreased 5 years after orthodontic treatment in the removable retainer group ( $P < 0.05$ ). There was no significant change in the fixed retainer group for the same variables ( $P > 0.05$ ). Intercanine width and anterior irregularity index showed a significant correlation during the retention period (upper  $r = 0.27$ ,  $P < 0.05$ , lower  $r = -0.30$ ,  $P < 0.05$ ). There was no significant correlation between arch length change and anterior irregularity index during the retention period ( $P > 0.05$ ). There was no significant difference in intercanine width, intermolar width and arch length between the extraction and non-extraction cases.

**CONCLUSION:** Post-treatment changes in arch form and dentition are significantly less with fixed than removable retainers.

## SP 227

Thursday, June 17 - at 13:00

### ELECTRON BEAM IRRADIATION OF POLYMER BRACKETS

**A Faltermeier, M Behr, C Reicheneder, P Proff, P Römer,** University of Regensburg, Germany

**AIM:** Polymer brackets have some disadvantages because of their reduced colour stability and insufficient load resistance. The aim of this study was to improve the mechanical stability of polymer brackets via polymer cross-linking using electron beam irradiation.

**MATERIALS AND METHOD:** The electron beam irradiation processor was manufactured in Bruchsaal (Germany, Beta Gamma Service). An acceleration voltage 10 MeV was chosen. The energy dose was 100 kGy. Irradiated samples provided from commercially available polymer brackets and polymer bracket raw materials (PC, POM, UDMA, HDPE) were identified. Fracture toughness, Vickers hardness and wear resistance were investigated. Statistical analysis was carried out using the Mann-Whitney  $U$  test. The level of significance was set at  $\alpha = 0.05$ .

**RESULTS:** Electron beam irradiation caused an improvement of mechanical properties of some samples. HDPE revealed a significant increase of wear resistance and Vickers hardness. However, polyoxymethylene brackets showed almost no change in their mechanical properties compared with the untreated control.

**CONCLUSION:** Electron beam irradiation seems to be able to change the mechanical properties of polymer brackets. Polymer brackets with lower mechanical properties appear to benefit more from electron beam post-curing.

## SP 228

Thursday, June 17 - at 13:00

### THE INFLUENCE OF ORTHODONTIC ADHESIVE FILLER LEVEL ON BACTERIAL ADHESION AND BOND STRENGTH

**A Faltermeier, C Reicheneder, P Proff, M Behr, P Römer,** University of Regensburg, Germany

**AIM:** To evaluate the effect of different filler contents of orthodontic adhesives on colour stability, shear bond strength (SBS), Adhesive Remnant Index (ARI), and bacterial adhesion.

**MATERIALS AND METHOD:** Four experimental adhesive groups were created: UDMA containing different volumes of SiO<sub>2</sub> filler: 0, 30, 50 and 70 per cent. Cylindrical specimens were incubated with *streptococcus mutans* and colonies were quantified using a luminescence kit and a plate reader. Other samples were exposed to artificial ultraviolet (UV) light for 72 hours. Tomato ketchup, coke and tea were chosen as food colourants, and colour measurements were performed according to the CIE L\*a\*b\* system. SBS and ARI were examined after bonding stainless steel brackets to extracted human third molars using the various adhesives. The means and standard deviations were calculated. Statistical analysis was performed using the Mann-Whitney  $U$  test. The level of significance was set at  $\alpha = 0.05$ .

**RESULTS:** After exposure to food dyes or UV light, higher filled adhesives presented the least and unfilled adhesive the greatest colour changes. Higher adhesive filler levels showed a greater bond strength between the enamel and brackets. Bacterial adhesion was also influenced by varying the filler level of the adhesives.

**CONCLUSION:** The properties of orthodontic adhesives can be strongly influenced by varying filler level. Bacterial adhesion, SBS and discolouration were altered using different amounts of SiO<sub>2</sub>.

## SP 229

Thursday, June 17 - at 13:00

### ROOT RESORPTION IN ORTHODONTICS: COMPARISON OF TWO TREATMENT TECHNIQUES

**E Pontarolo, G Masotto, F A Miotti,** University of Padova, Italy

**AIM:** Root resorption is a iatrogenic problem associated with orthodontic treatment. The low-friction Synergy (SY) system allows the use of light forces compared with the conventional straight-wire (SW) technique that requires heavier forces. The aim of this study was to compare root resorption after treatment with the conventional SW technique and the SY system.

**MATERIALS AND METHOD:** Pre- and post-treatment radiographs of 32 patients (16 SW, 16 SY) were analyzed. The radiographs were standardized and image distortion was compensated for. Measurements were undertaken by the same operator of root length (apex-cemento-enamel junction) and tooth length (apex-crown point). ANOVA tests were used to study the influence of gender and possible individual predisposition.

**RESULTS:** The SY system appeared to cause more frequent (SY 46.4%, SW 35.7%), though apparently less, severe resorption. Considering tooth length before and after treatment, canines and premolars appeared to show some increase, indicating a possibly reduced risk of damage in immature teeth. The anterior segments were more affected, with the upper lateral incisors showing the greatest root loss (4.3%). No correlation between gender and root damage was observed. The different distribution of resorption within individuals in the two groups appeared to indicate some predisposition to root resorption, more frequently observed in

the SW patients. Considering the suggested threshold of 2.5 mm as a clinically significant indication of resorption, 13 per cent of the teeth in the SY group were shortened above this value, 18 per cent in the SW group. In the SW group the roots in the posterior segments presented more significant resorption. In the SY group tooth damage was greater in males.

**CONCLUSION:** Less severe root damage could be expected with the use of the light-force low-friction SY system. As undetectable individual predisposition could be present in some cases, clinicians should always apply light forces independent of the type of mechanics used.

## SP 230

Thursday, June 17 - at 13:00

### EFFECT OF ORAL APPLIANCES ON INFLAMMATORY MARKERS IN OBSTRUCTIVE SLEEP APNOEA

**A Niżankowska-Jędrzejczyk<sup>1</sup>**, A Kania<sup>2</sup>, P Nastalek<sup>2</sup>, A Undas<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Medicine, Jagiellonian University, Cracow, Poland

**AIM:** Growing evidence demonstrates enhanced systemic inflammation in obstructive sleep apnoea (OSA) that might contribute to atherosclerotic vascular disease. Inflammatory markers including proinflammatory cytokines could be involved in this process. While continuous positive airway pressure treatment is known to decrease systemic inflammation in OSA, data on the effect of oral appliance (OA) treatment on cytokines are scarce. The aim of this study was to evaluate selected interleukins and hs-CRP in OSA patients prior to and following OA treatment.

**SUBJECTS AND METHOD:** Mild and moderate OSA patients [apnoea-hypopnoea index (AHI) 5-30/hours] and apparently healthy controls (HC) underwent polysomnography at baseline and then after 3 and 6 months. OSA patients were treated with a custom-made appliance, Klearway™, over 6 months. Interleukin-1 beta (IL-1β), IL-6, IL-10 and hs-CRP were measured in all subjects at all three time points. Interleukins were determined in plasma using ELISA (RAD systems) and serum hs-CRP using nephelometry (Dade Behring).

**RESULTS:** A total of 15 male OSA patients (mean age 54 years, AHI = 22/hours) and 6 HC (mean AHI 2/h) were analyzed. Demographic variables did not differ between the groups. OA treatment reduced symptoms of OSA and mean AHI from 22 to 9 hours after 6 months ( $P < 0.01$ ). No serious side-effects of treatment were observed. The baseline levels of IL-1β and IL-10 were significantly higher in OSA patients and decreased by 29 and 57 per cent, respectively (both  $P < 0.05$ ) after 6 months of OA. There were no changes in IL-6 and hs-CRP levels between OSA patients and HC at baseline and these variables remained unaltered during OA treatment.

**CONCLUSION:** OA therapy not only diminishes OSA symptoms and AHI but also modulates the inflammatory state via mechanisms independent from IL-6.

## SP 231

Thursday, June 17 - at 13:00

### STABILITY OF OPEN BITE CORRECTION IN CLASS II DIVISION 1 ADULTS WITH AND WITHOUT OSTEOTOMY

**R B Kuitert**, A Zentner, Department of Orthodontics, ACTA, Universiteit van Amsterdam and VU University, Amsterdam, Netherlands

**AIM:** To compare the stability of overbite correction after surgical and non-surgical correction of Class II division 1 malocclusion subjects with an open bite, hypothesizing more stability after surgical correction.

**SUBJECTS AND METHOD:** Thirty-six adults with a Class II division 1 malocclusion with open bites ranging 3.5 to 11.5 mm. Nineteen underwent orthodontic treatment only, and 17 had surgical-orthodontic treatment (a Le Fort I osteotomy in 10 cases, or Le Fort I and bilateral sagittal split osteotomy in 7 cases). Pre- and post-treatment and long-term cephalograms eight years after treatment were analyzed by measuring dental and skeletal variables.

**RESULTS:** Significant differences were found only for the increase during treatment in lower face height and the palatoman-dibular angle in the surgery group. Post-treatment and long-term no significant differences in overbite or in other variables were found. Post-treatment overbite changes in both groups varied from -2.0 to +1.0 mm. The non-surgery group showed stability in 37 per cent, relapse in 55 per cent and overbite increase in 8 per cent, while in the surgery group these were 21, 64 and 15 per cent, respectively. The non-surgery group showed significantly more extrusion of the upper incisors during treatment. Extrusion of the upper and lower incisors during treatment was largely maintained after treatment. The non-surgery group showed significantly more retroclination of the upper and lower incisors, and these changes remained relatively stable. Correlation analyses in the non-surgery group did not show any relationship between post-treatment overbite changes and changes in other variables during or after treatment. Post-treatment overbite changes in the surgery group were correlated with pre- and post-treatment vertical maxillary anterior dimensions and post-treatment lower face height.

**CONCLUSION:** Both orthodontic treatment only and combined surgical-orthodontic treatment seem to provide equal overbite stability after correction of Class II division 1 open bite malocclusions.

## SP 232

Thursday, June 17 - at 13:00

### DIMENSIONAL CHANGES OF THE MAXILLARY APICAL BASE IN UNTREATED CLASS I SUBJECTS

**J van der Vlag**, R B Kuitert, A Zentner, Department of Orthodontics, ACTA, Universiteit van Amsterdam and VU University, Netherlands

**AIM:** To assess maxillary apical base dimensional changes from the primary to the permanent dentition in untreated Class I subjects.

**MATERIALS AND METHOD:** Maxillary plaster casts of 38 patients were obtained from the material of a growth study carried out in the 1960s and assessed using Digimodel®. Inclusion criteria were high quality study casts with a full buccal sulcus corresponding to the following stages of dental development full primary dentition (T1; erupted first permanent molars allowed), early mixed dentition (T2) and permanent dentition (TR3; eruption of all permanent teeth mesial of the first permanent molar). The apical base was defined as a line running through the most concave point of the buccal sulcus in the region of the tooth apices. The apical base measurements were: 1) molar width (MW), the point just mesial of the first permanent molar (or distal of the second premolar), 2) canine width (CW), measured at the buccal cusp tip, 3) the circumference of the outline running from the first molar to the contralateral molar and, 4) the surface of this area enclosed by the molar width line.

**RESULTS:** T1-T2, MW and CW increased by 5 and 11 per cent, respectively. No decrease in any measurement was found. Circumference measurement remained constant, the surface measurement decreased by 10 per cent. T2-T3, MW did not change, circumference and CW decreased by 8 and 10 per cent, respectively, while CW returned to its original value. The surface decreased by 42 per cent.

**CONCLUSION:** Over time, MW increased first and then decreased, to end below the T1 value. For CW, a small increase was seen, ending below the T1 value. Circumference remained constant from T1 to T2 and ended below its T1 value. The surface measurement showed a decrease over time, to end around 50 per cent of its T1 value.

### FACIAL TRANSVERSE AND VERTICAL DEVELOPMENT DURING PRE-PUBERTAL GROWTH IN YOUNG BRAZILIANS

P S G Quaglia, A M Quaggio, C F A Salvador, R Bigliuzzi, **K Faltin Jr.**, Department of Orthodontics, University Paulista, São Paulo, Brazil

**AIM:** To compare transverse growth at different radiographic times and establish average reference values and sexual dimorphism for angular cephalometric measurements at the end of the observation period.

**MATERIALS AND METHOD:** Cephalometric measurements obtained from frontal telerradiographs of 32 subjects (13 males, 19 females) with an initial mean age of 7.7 years. All were in the mixed dentition phase with satisfactory dental occlusion and without a history of previous orthodontic or orthopaedic treatment. Each child was followed for an average period of 46 months. Three radiographs were obtained at three different time points – a total of 96 radiographs. The average interval between T0 and T1 was 566 days and between T1 and T2, 779 days.

**RESULTS:** There was an increase of the average values for the following cephalometric measurements: facial transverse, maxillary transverse, mandibular transverse, occlusal vertical and total vertical dimensions. The average values for angular cephalometric measurements at T2 were: maxillary posture  $40.70 \pm 4.09$ ; mandibular posture  $73.20 \pm 2.69$ ; and craniofacial posture  $15.13 \pm 2.49$ .

**CONCLUSION:** There is sexual dimorphism for maxillary posture cephalometric measurement but no statistically significant difference between genders for the mandibular posture and craniofacial posture.

### IN VITRO STUDY LABORATORY TEST OF THE SHEAR BOND STRENGTH OF METAL BRACKETS BONDED TO ENAMEL

M Di Lorenzo, **S I Penco**, A Nakhleh, Italy

**AIM:** To compare, *in vitro*, the shear bond strength (SBS) of five different types of metal orthodontic brackets with diverse mechanical retention on the enamel surface of bovine teeth.

**MATERIALS AND METHOD:** The lateral incisor teeth of 50 young cattle with undamaged buccal enamel, no cracks caused by the extraction forceps, and no defects observed by two investigators with a magnifier. The teeth were polished with pumice and water containing no fluoride, and embedded in a self-curing acrylic resin. Five types of orthodontic metal brackets were selected, 10 of each type (G1: miniequilibrium, Dentaureum, Pforzheim, Germany; G2 elite opti-MIM mini-twin regular profile, Ortho Organizers, San Marcos, California, USA.; G3: mini-dynalock, 3M Unitek, Monrovia, California, USA.; G4: minisprint, Forestadent, Pforzheim, Germany; G5 mini-diagonal, Leone, Italy) and bonded to the enamel surfaces of the teeth with Pad Lock (Reliance Orthodontic Products, Inc., Illinois, USA). The specimens were tested for SBS with an Instron universal testing machine. The base surface of the brackets after debonding was observed using a microscope at X20. Data obtained in Newton and Megapascal were analyzed with descriptive statistics. The adhesive fracture site was classified with the adhesive remnant index (ARI).

**RESULTS:** All groups showed statistically differences from each other, but had sufficient SBS to resist orthodontic forces. G1 showed the greatest bond strength, both in Newton and Megapascal. The ARI confirmed the greater retention capacity of the surface and the lower percentage of residual composite on the buccal enamel after debonding.

**CONCLUSION:** The different mechanical retention of the bracket bases influences bond strength. Although the dimensions of the brackets were small, all groups showed adequate bond strength.

### LONG-TERM OUTCOME AND STABILITY IN TREATED ANGLE CLASS II DIVISION 1 PATIENTS

**K Nordås**, G Slette, M Midtbø, O E Bøe, Department of Clinical Dentistry, Orthodontic Division, University of Bergen, Norway

**AIM:** To evaluate long-term hard and soft tissue changes in Class II division 1 patients treated with fixed orthodontic appliances.

**MATERIALS AND METHOD:** Lateral cephalograms of 68 patients, 40 extraction and 28 non-extraction cases. Lateral cephalograms taken pre-treatment (T1), post-treatment (T2), post-retention (T3), and 10 years post-retention (T4) were digitized and traced.

**RESULTS:** No statistically significant changes were found between the extraction and non-extraction group for soft and hard tissue variables, except for gonial angle. Significant longitudinal changes occurred from T1 to T2. ANB and SNA showed a marked reduction during T1-T2 and remained stable thereafter. SNB and SNPog increased during the whole observation period (T1-T4). Interbasal angle (ML/NL) decreased continuously from T1-T4. Both upper and lower incisors showed significant uprighting during treatment. Post-treatment there was a tendency towards relapse. Upper and lower lip distance to the E-line increased progressively from T1-T4, most marked from T1-T2 and the Holdaway angle decreased.

**CONCLUSION:** Prior to treatment, during treatment and post-treatment, the extraction and non-extraction Class II division 1 subjects presented identical hard and soft tissue profile characteristics. There was a tendency towards dental relapse 10 years post-retention, a marked lip retrusion relative to the E line, and a decrease in the Holdaway angle.

### IMPACTED MAXILLARY CANINES: IMPLICATIONS FOR ORTHODONTIC TREATMENT ON ROOT LENGTH

**E M G Brusveen**, M Mavragani, P Brudvik, O E Bøe, Department of Clinical Dentistry, Division of Orthodontics, University of Bergen, Norway

**AIM:** The incidence of impacted maxillary canines varies from 1 to 3 per cent. Root resorption of maxillary incisors is a common sequela of maxillary canine impaction. Treatment of impacted canines is often challenging and causes an additional burden on neighbouring teeth. The aim of this study was to evaluate the effect of maxillary canine impaction on root resorption caused by orthodontic treatment. The effect of several characteristics of the impacted canine on root shortening of the adjacent incisors was also studied.

**MATERIALS AND METHOD:** Orthodontic records of 64 treated patient. Thirty-two patients had a unilateral impacted maxillary canine, while the rest did not have any impaction. All were treated with fixed appliances. Crown and root lengths of the maxillary incisors were measured on pre- and post-treatment periapical radiographs corrected for image distortion. The percentage of root shortening and root length loss in millimetres was then calculated. The inclination of the eruption path of the impacted canine, relative to the midline, the long axis of the lateral incisor, the occlusal plane and the nasal line, was recorded on dental pantomograms (DPTs) and lateral cephalometric films taken before any treatment. Horizontal and vertical impacted canine position was also evaluated on the DPTs. Impacted canine root development and width of the dental follicle were also recorded.

**RESULTS:** No significant difference was found in root resorption of the maxillary incisors during orthodontic treatment between patients with impacted canines and those without impaction.

**CONCLUSION:** It seems that the treatment of impacted canines does not cause any additional burden to the roots of the neighbouring incisors, than normal orthodontic treatment of dentitions without impacted canines.

### AUTOMATED TRACKING OF FACIAL LANDMARKS FOR THREE-DIMENSIONAL ANIMATED MODELS, USING STEREOPHOTOGRAMMETRY

**E Di Gioia**<sup>1</sup>, R Deli<sup>1</sup>, L M Galantucci<sup>2</sup>, G Percoco<sup>2</sup>, <sup>1</sup>Scuola di Specializzazione in Ortognatodonzia, Università Cattolica del Sacro Cuore di Roma and <sup>2</sup>DIMEG Lab. Prototipazione Rapida e Reverse Engineering, Politecnico di Bari, Italy

**AIM:** Correct three-dimensional (3D) diagnostic classification of developmental disharmonies is essential for assessing rehabilitation of facial soft tissue and skull-facial structures, and for monitoring the changes induced by surgical, aesthetic and functional treatment. The aim of this study was to evaluate a simple and economic digital close range photogrammetric system developed for measurement and tracking of facial movements. The system produces a 3D animated model (wire frame, tasselled and textured surfaces), which accurately reproduces the subject's facial movements during opening and closing of the mouth.

**MATERIALS AND METHOD:** Three Canon EOS 40D, calibrated cameras were used. A 12 bit coded target L frame provided the scale factor and also the fixed reference points during data acquisition. The main software used was PhotoModeler Video. A laser scanner Konica Minolta Vivid was positioned between the left and middle camera; in order to compare only the photogrammetrically acquired static data to the laser scanned acquired static data. Biometric measurements were first determined on a dummy face, and then on a subject's face to study the transformations of distances, angles and relationships between landmarks during motion.

**RESULT:** Recognition and tracking of facial landmarks was achieved by automated procedures that required minor human intervention. An animated 3D triangulated surface was produced, which also accurately reproduced the subject's movements. Accuracy values in the order of fractions of a millimetre can be achieved.

**CONCLUSION:** The use of a laser scanner allows 3D biometric point positions to be determined only statically and a set of values to be calculated, but its accuracy is much lower when compared with the photogrammetric technique, and motion capture is not possible.

### PULPAL REPAIR AFTER SUBCUTANEOUS TRANSPLANTATION OF CRYOPRESERVED IMMATURE TEETH IN RATS

**L Temmerman**<sup>1</sup>, A Vral<sup>2</sup>, E Deschepper<sup>3</sup>, G De Pauw<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Anatomy, Embryology, Histology and Medical Physics and <sup>3</sup>Biostatistics Unit, Ghent University, Belgium

**AIM:** Pulpal tissue of an immature tooth with an open apex can maintain its viability during cryopreservation. However, pulpal tissue reactions after autotransplantation of cryopreserved teeth are not yet well understood. The purpose of this study was to evaluate in vivo whether revascularisation after transplantation of cryopreserved immature teeth occurs, comparable with non-cryopreserved transplanted teeth.

**MATERIALS AND METHOD:** In 24, four-week-old Wistar rats the first and second maxillary molars were extracted bilaterally. In 12 rats, the molars were cryopreserved before transplantation (test group). In the remaining 12 rats, the molars were transplanted subcutaneously in the abdomen immediately after extraction (control group). Material was collected for the test and control animals at intervals of 1, 2, 4 and 10 weeks after transplantation, and histological examination was performed.

**RESULTS:** In the first week after transplantation, degeneration of the original pulpal tissue was observed in the test and control teeth. At the root apex, repair started by ingrowth of highly vascular granulation tissue. Two weeks after transplantation, ingrowth of

new connective tissue was observed at the apex originating from the adjacent periodontal tissues. In the cryopreserved teeth degenerated pulpal tissue was still present coronally, while in most of the control teeth the newly formed connective tissue had replaced the original degenerated tissue. Cellular mineralized tissue at the apex and dentine-like tissue in the root canal were present in most of the controls, while these tissues were rarely seen in the test teeth. At week 4, these differences were still recognizable, but in general bone-like and dentine-like hard tissues increased in the root canal and coronal pulp of all the teeth. At week 10, no differences between the test and control teeth were observed.

**CONCLUSION:** Regeneration of pulpal tissue after autotransplantation of cryopreserved immature teeth was similar to immediately transplanted teeth, although the healing process was slower.

### THIN-PLATE SPLINE ANALYSIS OF THE DIFFERENT FACIAL TYPES IN FRONTAL NORM TELERADIOGRAPHS

E A Walverde<sup>1</sup>, C F A Salvador<sup>2</sup>, A Quaggio<sup>2</sup>, R Bigliuzzi<sup>2</sup>, **K Faltin Jr.**<sup>2</sup>, <sup>1</sup>Private Practice and <sup>2</sup>Department of Orthodontics, University Paulista, São Paulo, Brazil

**AIM:** To evaluate facial morphology in teleradiographs with respect to normal values derived from frontal radiographs using thin plate spline (TPS) analysis.

**SUBJECTS AND METHOD:** Thirty subjects (15 females, 15 males), all in the mixed dentition phase with ages ranging from 8.4 to 10 years with satisfactory occlusion and harmonic profile with no history of orthodontic or orthopaedic facial treatment. The sample was divided into three groups according to the facial types proposed by Ricketts (1982). The coordinates of the anatomical points of the traces scanned from the teleradiographs were obtained using appropriate software (tpsDig2). The average craniofacial configurations of each group were obtained by superimposing orthogonal Procrustes, eliminating the differences in size, and analysing only the shape differences between groups by the visualization of the grid of deformations of the TPS method.

**RESULTS:** Statistically significant differences were observed between the three facial types but they were more marked between mesofacial types than between the mesofacial and brachyfacial subjects.

**CONCLUSION:** TPS morphometric analysis is efficient for accurate visualization of transverse and vertical differences among facial types even before the peak of pubertal growth.

### A TWO-JAW ORTHODONTIC APPLIANCE FOR A CLASS III MALOCCLUSION PATIENT TREATMENT

**M Postnikov**, G Stepanov, N Pankratova, L S Persin, T Odnokova, Samara State Medical University and Moscow State University of Medicine and Dentistry, Russia

**AIM:** To improve methods of orthodontic treatment in children with a Class III malocclusion combined with early extraction of molars.

**SUBJECTS AND METHOD:** Forty patients aged 6-9 years with a Class III malocclusion combined with early molar extraction. Clinical, anthropometric, morphometric and roentgenologic methods were used. Space deficiency for premolars was estimated and space in the extraction region was measured using the Michigan University scheme. A multifunctional two-jaw orthodontic prosthetic device for Class III and deep-bite treatment (patent 12307620 from 10 October 2007) was used. The appliance consisted of two basic plastic plates (upper and lower), functional, fixed and strength elements and contained artificial teeth to replace the missing molars. The upper and lower basic plates were connected by Class III elastics.

**RESULTS:** The occlusal anomaly was corrected. Orthodontic treatment improved interpersonal relationship and occlusal function.

**SP 241**

Thursday, June 17 - at 13:45

**A SYSTEMATIC REVIEW OF THE CORRELATION BETWEEN THE STOMATOGNATHIC SYSTEM AND BODY POSTURE****G Perinetti**, P Gabrieli, L Contardo, R Di Lenarda, Department of Biomedicine, Unit of Orthodontics, University of Trieste, Italy

**AIM:** Correlations between the stomatognathic system and whole-body posture has recently been reported. Among the different recording methods used is surface electromyography. This method is able to monitor the muscle activity of both the stomatognathic system and other areas of the body. The aim of this systematic literature review was to assess the scientific evidence for detectable correlations between the stomatognathic system and whole-body posture as investigated through surface electromyography.

**MATERIALS AND METHOD:** A literature survey was performed using the Medline database, covering the period from January 1966 to September 2009, and using the medical subject headings. After selection, seven articles qualified for final analysis.

**RESULTS:** Only one study was judged to be of medium quality, with all the rest classified as of low quality design. None of the studies included had a follow-up. Overall, all of the studies reported detectable correlations between the stomatognathic system and body posture for at least one of the parameters used. However, after reappraisal of most of the data provided in these studies, a weak correlation that reached biological relevance is apparent. Therefore, with limitations due to the poor methodological quality of the published studies, a correlation between the stomatognathic system and whole-body posture can be detected, at least under experimental conditions.

**CONCLUSION:** While more investigations with improved levels of scientific evidence are needed, the current evidence does not support either the existence of clinically significant correlations between the stomatognathic system and whole-body posture.

**SP 242**

Thursday, June 17 - at 13:45

**THE POSITION OF THE HYOID BONE INVESTIGATED BY DIFFERENT METHODS OF ANALYSIS OF A RADIOGRAPH****I Kirovski**<sup>1</sup>, G Curcievea-Cuckova<sup>2</sup>, S Popovski<sup>2</sup>, R Dimovska<sup>3</sup>, T Kirovska<sup>4</sup>, <sup>1</sup>Stomatological Department, <sup>2</sup>Stomatological Faculty, University 'St. Cyril and Methodius', <sup>3</sup>'Kruna' Private practice, <sup>4</sup>'Dr. Tatjana Kirovska' Private practice, Skopje, Former Yugoslav Republic of Macedonia

**AIM:** To investigate the position of the hyoid bone.

**SUBJECTS AND METHOD:** Sixty subjects, evenly divided into two groups: those with a normal posterior position of the mandible and a control group with a normal mandibular position. The position of the hyoid bone was determined using different methods suggested in the literature and compared with the spine, mandible and cranium.

**RESULTS:** In general, the hyoid bone followed the posterior position of the mandible, though less in proportion. Comparison of the results with published studies was not possible due to the different methodologies used.

**SP 243**

Thursday, June 17 - at 13:45

**APPRAISAL OF THE GEOMETRIC EFFECTS ON TWO CEPHALOMETRIC INDICES OF SKELETAL CLASS****E Di Maria**, M Ceschi, G Perinetti, A Castaldo, L Contardo, Department of Biomedicine, Unit of Orthodontics, University of Trieste, Italy

**AIM:** The importance of accurate anteroposterior measurement of the maxillo-mandibular relationships, i.e. skeletal Class, is critical in orthodontic diagnosis. Among the most popular cephalometric indices are ANB angle and Wits appraisal on the functional occlusal plane. However, these indices are geometrically sensitive and can give false results. The present investigation was thus

aimed at providing quantitative knowledge for the most relevant geometric effects on these indices, which can be corrected on the individual skeletal morphological features.

**MATERIALS AND METHOD:** One-hundred-forty-four lateral cephalograms of 8-to-14-year-old subjects seeking orthodontic treatment were traced. Seventeen different cephalometric variables were recorded and used as independent variables in backward multiple regressions; the outcomes were either ANB angle or Wits appraisal. The most relevant associations were thus retrieved for both these two indices of skeletal Class.

**RESULTS:** ANB was mostly correlated with the mandibular plane angle to S-N plane and SNA angle. Wits appraisal was mostly correlated with the mandibular plane and bisected occlusal plane angle to the S-N plane. Through the multiple regression models, correction coefficients were retrieved for each of the indices. Subsequently, corrected indices were calculated on an individual basis. In the whole sample, the correlation coefficients for the uncorrected ANB and Wits appraisal was 0.72, and reached 0.96 between the corresponding corrected indices. Through elimination of geometric distorting effects, the increase in agreement between the indices of skeletal Class is expected to be consistent with a more accurate diagnosis that would be based on individual corrections.

**CONCLUSION:** Cephalometric indices of skeletal Class are subject to geometric effects making individual corrections useful to avoid bias in diagnosis.

**SP 244**

Thursday, June 17 - at 13:45

**IS CASEIN PHOSHOPEPTIDE-AMORPHOUS CALCIUM PHOSPHATE BENEFICIAL TO ORTHODONTIC PATIENTS?****S Momeni Danaei**, A Karamifar, H Pakshir, **S Saadat**, F Khaje, Orthodontic Research Centre, Shiraz Dental College, Iran

**AIM:** Demineralization of enamel adjacent to orthodontic brackets can be an aesthetic and functional problem. A number of strategies are available to treat these lesions such as casein phosphopeptide-amorphous calcium phosphate (CPP-ACP). The aim of this study was to evaluate the effect of Tooth Mousse™ topical cream in orthodontic patients.

**SUBJECTS AND METHOD:** Thirty patients divided into three equal groups. No patient had any white spot lesions (WSL) and all underwent oral hygiene instruction. Group 1 used a toothbrush and superflossed twice daily, group 2 used a toothbrush, superflossed twice daily, and used a fluoride mouth rinse before sleep, group 3 used a toothbrush, superflossed twice daily and used a fluoride mouthrinse and tooth mousse before sleep. The patients were examined at 1, 3 and 6 monthly intervals to evaluate the incidence of WSL on the buccal surfaces of 20 teeth from the central incisor to the second premolar in each quadrant by a specialist, and the number of WSL were recorded.

**RESULTS:** After statistical analysis, the mean prevalence of WSL after 6 months was group 1, 7.3, group 2, 4.7 and group 3, 1.6. There was a statistically significant difference between groups 1 and 2, groups 1 and 3, and also between groups 2 and 3.

**CONCLUSION:** CPP-ACP agents such as Tooth Mousse™ can help in protecting teeth from WSL. Tooth Mousse™ topical cream has additive effects to fluoride mouth rinses and helps remineralize teeth by delivering calcium substrate to tooth structure.

**SP 245**

Thursday, June 17 - at 13:45

**ORTHODONTIC STATUS AND ORTHODONTIC TREATMENT NEED AMONG SEVENTH GRADE STUDENTS****E Deljo**, Healthcare 'Dr Isak Samokovlija', Goražde, Bosnia and Herzegovina

**AIMS:** To determine the prevalence of malocclusion and orthodontic treatment need among students of seventh grade classes in Gorazde (Bosnia and Herzegovina) and to compare the results with other populations.

**SUBJECTS AND METHOD:** Three hundred and fifty eight students from the seventh grade classes in Bosnian-Podrinje Canton (Bosnia and Herzegovina) aged from 12-13 years. The measurement instrument was the Dental Aesthetic Index (DAI).

**RESULTS:** The incidence of orthodontic anomalies was 60.44 per cent and the average value of the DAI 25.25 (SD 8.771). The percentage of students requiring orthodontic therapy (DAI  $\geq$  31) was 29.61 per cent. There were no statistically significant differences in DAI score between genders [girls 24.70 (SD: 8.784), boys 25.71 (SD: 8.755)].

**CONCLUSION:** The distribution of DAI scores in this sample of students is not very different to that reported for other European populations.

## SP 246

Thursday, June 17 - at 13:45

### BOTULINUM TOXIN INFLUENCES THE HISTOLOGY AND GENE EXPRESSION OF DYSTROPHIC MASSETER MUSCLE

**T Gedrange**<sup>1</sup>, A Spassov<sup>1</sup>, T Gredes<sup>1</sup>, T Klinke<sup>2</sup>, C Kunert-Keil<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Prosthodontics, EMAU Greifswald, Germany

**AIM:** Duchenne Muscular Dystrophy (DMD) and its murine model, mdx, are characterized by Ca<sup>2+</sup> induced muscle damage and muscle weakness. The disease is caused by mutations in the gene encoding dystrophin. Masticatory muscles of DMD patients show hypertrophic effects followed by distorted dentofacial morphology. Masseteric hypertrophy can be treated with a botulinum toxin A (BTX-A) injection into the masseter muscle. The aim of this study was to evaluate the changes in muscle histology and in the MyHC mRNA expression in muscle tissue specimens from masseter, temporal and tongue muscles of control and mdx mice after injection of BTX-A.

**MATERIALS AND METHOD:** One hundred day-old mice (12 unilateral, 12 bilateral treated animals, 12 untreated animals of both mice strains, mdx and controls). BTX-A was injected into the right or both masseter(s) to initialise total immobilisation. After 6 weeks, muscle tissue was taken from the left and right side of the masseter, temporal, and tongue muscles using a standardized method. The muscle cross-sections were stained with haemalaun/eosin or with Sirius Red and morphometrically analyzed. The MyHC mRNA levels were studied using quantitative RT-polymerase chain reaction.

**RESULTS:** The muscles adapted to such stress by changing fibre types and MyHC mRNA content. Paralyzed masseters displayed atrophic changes, while other masticatory muscles showed hypertrophic changes as demonstrated by changes in the mean fibre diameter. The diameters of the muscle fibres of the BTX-A groups were more irregular and an increase in the amount of muscle fibres with central nuclei was observed in both mice strains after BTX-A treatment.

**CONCLUSION:** Treatment with BTX-A may lead to structural changes in affected and unaffected muscles. These changes may cause changes in muscle balance and could compensate the dystrophic effects.

## SP 247

Thursday, June 17 - at 13:45

### EXPRESSION OF GENES INVOLVED IN CALCIUM HOMEOSTASIS IN DYSTROPHIC MASTICATORY MUSCLES

**C Kunert-Keil**, T Gredes, S Lucke, T Gedrange, A Spassov, Department of Orthodontics, EMAU Greifswald, Germany

**AIM:** Duchenne muscular dystrophy (DMD) and its murine model, mdx, are characterized by calcium ions (Ca<sup>2+</sup>) induced muscle damage and muscle weakness. The disease is caused by mutations in the gene encoding dystrophin. DMD patients have distorted dentofacial morphology that could be a result of changed masticatory muscles force balance due to unequal dystrophic changes in various masticatory muscles. Recently it was shown, that the masticatory muscles of mdx mice are differentially affected by the disease. Since an altered Ca<sup>2+</sup> homeostasis seems to

underlie mdx muscle pathology, the expression of genes involved in the Ca<sup>2+</sup> homeostasis, in particular, ion channels, ion exchangers and Ca<sup>2+</sup> handling proteins were studied.

**MATERIALS AND METHOD:** The mRNA levels of the sarcoplasmic reticulum Ca<sup>2+</sup> ATPases SERCA1 and SERCA2, the plasma membrane Ca<sup>2+</sup> ATPases Atp2b1 and Atp2b4, the sodium/calcium exchanger NCX1, the ryanodine receptor 1, parvalbumin, sarcolipin, phospholamban and the L-type Ca<sup>2+</sup> channel Cav1.1 were studied using quantitative RT polymerase chain reaction in the masseter and temporalis muscles and the tongue of control and mdx mice.

**RESULTS:** In mdx masseter muscle no differences were found in the expression of any tested genes compared with the control mice. In contrast, the mRNA amounts of SERCA2 and RYR1 were significantly reduced in mdx temporalis muscle. In mdx, the tongue a down-regulation of the sarcolipin and parvalbumin mRNA expression was found, whereas the phospholamban mRNA level was significantly increased compared with the controls.

**CONCLUSION:** mdx masticatory muscles showed an unequal modified expression of genes involved in Ca<sup>2+</sup> homeostasis. This could cause the functional imbalance of masticatory muscles in DMD patients followed by morphological changes, which are observed in this disorder.

## SP 248

Thursday, June 17 - at 13:45

### INFLUENCE OF THE MAGNITUDE OF INTRUSIVE FORCE ON EXTERNAL APICAL ROOT RESORPTION AND CHANGES IN ALVEOLAR CREST HEIGHT

**I Marek**, M Kamínek, M Spidlen, K Langova, Orthodontic Department, Dental Faculty, University of Palacky, Olomouc, Czech Republic

**AIM:** Determination of the interdependence of the extent of external apical root resorption (EARR) and bone remodelling of the alveolar ridge on the amount of applied orthodontic intrusive force and the correlation between resorption and the extent of intrusion.

**SUBJECTS AND METHOD:** Seventeen patients who required fixed appliance treatment together with the extraction of the maxillary first premolars. A unilateral force of 50 g was applied for six months to the premolars and a force of 150 g to the contralateral teeth. The teeth were then extracted and measured. Computed tomographic (CT) scans were obtained before and after treatment. On all CT scans the length of the tooth, root, and crown were measured pre- and post-treatment.

**RESULTS:** The mean intrusion was 1.69 mm with an intrusive force of 50 g and 4.86 mm with an intrusive force of 150 g. The extent of tooth intrusion depended on the extent of the force. The dependence was statistically significant. With an intrusive force of 50 g, the mean change in vestibular root length was 0.44 mm and with a force of 150 g, 0.73 mm. The values were not statistically significant. The mean value for the sulcus was 0.63 mm (50 g force), and 2.17 mm (150 g force). With a force of 50 g, bone sulcus represented 41 per cent of intrusion and with a force of 150 g, 44 per cent of the intrusion.

**CONCLUSION:** EARR does not depend on the extent of the intrusive force. However, force is a statistically significant factor in the amount of tooth intrusion. During intrusion, the alveolar ridge dislocated in the direction of the tooth movement, and the tooth was partially shifted into the bone.

## SP 249

Thursday, June 17 - at 13:45

### OCCUSAL RELATIONSHIPS IN THE EARLY BRONZE AGE IN SOUTH EASTERN EUROPE

**T Pajevic**, J Juloski, D Cubrilo, I Scepan, B Glisic, Department of Orthodontics, University of Belgrade, Serbia

**AIM:** Inheritance is a significant causal factor in the aetiology of malocclusions. However, environmental factors, in particular the type of diet, also have an impact on teeth, bones, neuromuscular

tissues and soft tissues. The territory of today's Serbia, in the early Bronze Age was the settlement of farmers, who domesticated plants and animals. Their diet consisted of softer food, similar to contemporary human diets. The aim of this study was to examine the occlusal relationships and presence of orthodontic anomalies in skull remains of an early Bronze Age society.

**MATERIALS AND METHOD:** Three hundred and twelve skulls of individuals from the necropolis in Mokrin (Serbia) dating between 2100 and 1800 BC. Well-preserved skulls, where occlusal relationships could be reconstructed were used. Analysis of sagittal and transverse relationships was undertaken. Sagittal relationships were analyzed based on Angle's classification – the relationship of the mesiobuccal cusp of the upper first molar and buccal groove of the lower molar was observed. In the transverse plane, the bucco-oral relationship of the upper and lower molars was analyzed. Malocclusions in the vertical plane were not investigated because of the *post-mortem* loss of the anterior teeth in most individuals. The Statistical Package for Social Sciences, version 12.0 (SPSS Inc., Chicago, Illinois, USA was used for statistical analysis.

**RESULTS:** A Class I occlusal relationship was observed in 54.5 per cent of the skulls, while 45.5 per cent had a Class II relationship. Normal transverse relationships were found in 75 per cent, while 25 per cent exhibited a unilateral crossbite.

**CONCLUSION:** Malocclusions were present 4000 years ago. These findings indicate that life style and dietary habits, including the type of food and its consistency, could be important factors in the aetiology of malocclusions.

## SP 250

Thursday, June 17 - at 14:30

### PREVALENCE OF MALOCCLUSION IN PATIENTS WITH TURNER SYNDROME

**C Misevska<sup>1</sup>**, M Kocova<sup>2</sup>, N Toseska<sup>1</sup>, J Gjorgova<sup>1</sup>, E Bajraktarova Valjakova<sup>3</sup>, <sup>1</sup>Department of Orthodontics, <sup>2</sup>Pediatric Clinic and <sup>3</sup>Department of Prosthetics, Ss. Cyril and Methodius University, Former Yugoslav Republic of Macedonia

**AIM:** To assess the frequency of orthodontic malocclusions in patients with Turner syndrome.

**SUBJECTS AND METHOD:** Thirty Turner syndrome patients aged 8-18 years (mean age 14.7). The sample was subdivided according to karyotype.

**RESULTS:** A malocclusion was found in 70 per cent of patients with Turner syndrome. A Class II division 1 malocclusion was most frequently observed (30%). Crowding and a bilateral crossbite were found in 6.7 per cent of the subjects, respectively. A unilateral crossbite was present in 13.3 per cent whereas an open bite was observed in only 3.3 per cent. The most significant differences were found between 45X patients and mosaic and isochromosome for the long arm of X karyotypes. The prevalence of a Class II division 1 malocclusion was the most frequently anomaly in 45X patients, whereas Class II division 2 malocclusion was most often found in patients with mosaic karyotypes.

**CONCLUSION:** Patients with structural and/or numerical aberrations of the X chromosome develop a specific pattern of malocclusion with deviations in the sagittal, vertical and transverse directions. The findings show the importance of early diagnosis of oral anomalies and timely orthodontic treatment of patients with Turner syndrome.

## SP 251

Thursday, June 17 - at 14:30

### SOFT TISSUE PROFILE CHANGES IN THE EARLY TREATMENT OF A DEEP BITE WITH MYOFUNCTIONAL APPLIANCES

**M Jämsä<sup>1</sup>**, A Poikela<sup>1</sup>, S Kärkkäinen<sup>1</sup>, P Pirttiniemi<sup>2</sup>, <sup>1</sup>Oral Health Services, Oulu and <sup>2</sup>Department of Oral Development and Orthodontics Institute of Dentistry University of Oulu, Finland

**AIM:** To investigate, in a longitudinal randomized study, the soft tissue profile changes in deep bite malocclusion subjects when treatment is started during the early mixed dentition with a myofunctional appliance.

**SUBJECTS AND METHOD:** Twenty-eight children (16 females, 12 males, mean age 6.3 years) randomly divided into two groups. In the myofunctional appliance group (MA) treatment was initiated immediately while the control group received no orthodontic treatment. Digital profile photographs were taken of all subjects at the beginning of the study and after one year. The photographs were traced and the landmarks identified manually.

**RESULTS:** The mean change in soft tissue ANB angle in the MA group was -1.7 degrees (SD 0.97) and in the controls 0.1 (SD 1.44). The difference between the groups was significant ( $P < 0.001$ ). The mean change in the labiomental angle in the MA group was 5.6 degrees (SD 7.14) and in the controls 0.1 degrees (SD 5.41;  $P < 0.003$ ).

**CONCLUSION:** Treatment with myofunctional appliances changed the soft tissue profile by straightening the profile in the MA group when compared with the controls.

## SP 252

Thursday, June 17 - at 14:30

### ECTOPIC ERUPTION OF THE MAXILLARY FIRST PERMANENT MOLAR

**M Zuzelova<sup>1</sup>**, M Smileva-Nacevska<sup>2</sup>, G Kovacevska<sup>3</sup>, M Maneva<sup>1</sup>, G Curcieva-Cuckova<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>3</sup>Prosthodontics, PHO University Dental Clinical Center, <sup>2</sup>Private Dental Practice, Skopje, Former Yugoslav Republic of Macedonia

**AIM:** Ectopic eruption reflects the eruption of a tooth in an abnormal position. The most frequently found ectopic teeth are the maxillary first permanent molars and canines followed by the mandibular canine, mandibular second premolar, and maxillary lateral incisors. Ectopic eruption of the maxillary first molars shows a variable frequency between 2 and 6 per cent. The aim of this retrospective study was to describe the characteristics and occurrence of ectopic eruption of the maxillary first permanent molar and its correlation with pathological resorption of the maxillary second primary molar in a group of untreated normal growing Macedonian children.

**SUBJECTS AND METHOD:** Five hundred and twenty one consecutive patients in the first phase of the mixed dentition. They were reviewed from January 2005 to May 2009 as a routine procedure. All were evaluated with a dental pantomograms and intraoral radiographs, taken every six months. The position of the maxillary first permanent molar and the amount of pathological resorption of the second maxillary primary molar was evaluated and the radiographs of each patient were compared.

**RESULTS:** The frequency of occurrence was 5.3 per cent with no differences in gender. Of these, 46.7 per cent were unilateral and 53.6 per cent bilateral with a right-left relationship of 2:1. Almost half (52%) of ectopic molars spontaneously self-corrected.

**CONCLUSION:** Early diagnosis and treatment of the maxillary first permanent molar results in the prevention of premature loss of the second maxillary primary molar and resulting malocclusion.

## SP 253

Thursday, June 17 - at 14:30

### MAXILLARY SECOND AND THIRD MOLAR INCLINATION AFTER EXTRACTION OF THE FIRST MOLARS IN CLASS II DIVISION 1 SUBJECTS

**C Livas<sup>1,2</sup>**, D Halazonetis<sup>3</sup>, J W Boonij<sup>4</sup>, C Katsaros<sup>1</sup>, <sup>1</sup>Department of Orthodontics and Dentofacial Orthopaedics, University of Bern, Switzerland, Departments of Orthodontics, <sup>2</sup>251 Hellenic Air Force and Veterans General Hospital and <sup>3</sup>Dental School, University of Athens, Greece, <sup>4</sup>Department of Orthodontics and Oral Biology, Radboud University Nijmegen Medical Centre, Netherlands

**AIM:** To assess the change in the inclination of the maxillary second (M2) and third (M3) molars after orthodontic treatment of a Class II division 1 malocclusion with extraction of the maxillary first molars.

**SUBJECTS AND METHOD:** Two groups of individuals were studied. The experimental group consisted of 37 subjects (18 boys, 19 girls, mean age  $13.2 \pm 1.62$ ). The inclusion criteria were Caucasian origin, Class II division 1 malocclusion, sagittal overjet of  $\geq 4$  mm, no missing teeth or agenesis, and maxillary third molars present. All patients were treated with extraction of the maxillary first molars and treated using the Begg technique. Standardized lateral cephalometric radiographs were taken at the start of active treatment (T1) and at least 3.7 years post-treatment (T2). The control group was drawn from the archives of the Nittedal Growth Material (Oslo University) and included 54 untreated Class I and Class II individuals (18 boys, 36 girls, mean age  $13.4 \pm 1.99$ ) followed up for a minimum period of 3.6 years. Second and third maxillary molar inclination relative to the palatal (PP) and functional occlusal plane (FOP) was measured and compared between groups and time periods.

**RESULTS:** In the control group there was no significant change in the initial distal inclination of M2, whereas in the extraction group, although more distally inclined at T1, it was mesially inclined at T2 (M2-FOP T1  $14.2 \pm 4.62^\circ$ , T2  $-6.2 \pm 6.10^\circ$ ,  $P < 0.0001$ ). M3 inclination was similar between groups at T1 (M3-FOP control  $17.3 \pm 9.35^\circ$ , M3-FOP experimental  $19.6 \pm 7.37^\circ$ ), and improved significantly in both groups however M3 uprighting was almost 4 times greater in the extraction group (M3-FOP T2-T1:  $5.6$  versus  $19.9^\circ$ ). The largest distal inclination of M3 at T2 in the extraction group was  $9.4$  degrees, a value attained by only 43 per cent of the control group. **CONCLUSION:** Extraction of the maxillary first molars in Class II division 1 subjects results in significant uprighting of second and third molars and ensures the normal eruption of M3.

## SP 254

Thursday, June 17 - at 14:30

### KNOWLEDGE ON BIOSTATISTICS OF ORTHODONTIC POSTGRADUATE STUDENTS IN EUROPE

A Polychronopoulou<sup>1</sup>, T Eliades<sup>2</sup>, K Taoufik<sup>1</sup>, M Papadopoulos<sup>2</sup>, A Athanasiou<sup>2</sup>, <sup>1</sup>Department of Preventive and Community Dentistry, University of Athens and <sup>2</sup>Department of Orthodontics, Aristotle University of Thessaloniki, Greece

**AIM:** To explore the level of knowledge in biostatistics of orthodontic postgraduate students.

**MATERIALS AND METHOD:** A four part questionnaire, which included a knowledge test on biostatistics and epidemiology, was developed and distributed to postgraduate programme directors of 21 European universities in 10 countries who delivered it to students; the questionnaire was completed during an in-class examination. The frequency distributions of demographic characteristics were examined, and the percentages of participants who agreed or strongly agreed with each attitudinal statement were calculated; percentages of participants who felt fairly to highly confident for each confidence statement were also determined. Knowledge scores were calculated as the percentage of correct answers; missing values were counted as wrong answers. Data were analyzed with the Student's *t*-test or one-way analysis of variance, and multiple linear regression modelling was employed to determine the adjusted/unconfounded effect of possible knowledge score predictors.

**RESULTS:** One hundred and twenty seven orthodontic students from a total of 129, responded to the questionnaire. The mean correct answers of the participants were 43.8 per cent with a 95 per cent confidence interval (95% CI) of 40.2-47.3 per cent. This score was not influenced by gender, years elapsed from graduation, other advanced degree or year of study; the sole parameter, which seemed to influence this score was attendance at a biostatistics/epidemiology course (51.9 versus 39.5% score of participants who had previously taken a course versus those who had not,  $P < 0.001$ ). A surprising finding was the inability of the responders to identify the appropriate use of a chi-square test (11.8%, 95% CI 6.1-17.5%).

**CONCLUSION:** Knowledge on biostatistics of orthodontic postgraduate students in Europe is only influenced by previous relevant education and should be considered inadequate.

## SP 255

Thursday, June 17 - at 14:30

### NASOLABIAL ANGLE VARIATIONS DUE TO UPPER INCISOR TORQUE AND POSITIONAL CHANGES

J Seoane, University of Oviedo, Spain

**AIMS:** To find a direct relationship between pre-and post-treatment changes on upper incisor position and angle, and the modifications on the nasolabial angle in non-extraction orthodontic cases.

**SUBJECTS AND METHOD:** One hundred patients aged from 17 to 30 years, all orthodontically treated without extractions. Lateral telerradiographs from before and after treatment were cephalometrically measured. The points and planes for the following measurements were drawn: upper incisor angle based on the maxillary plane (ANS-PNS/SI), upper incisor anteroposterior position based on a plane perpendicular to the Frankfort horizontal plane that crosses point N (N<sup>^</sup>Po-Or) and nasolabial angle (me-sn-ls).

**RESULTS:** Based upon the differences pre- and post-treatment (final minus the beginning telerradiograph measurements), the nasolabial angle changed when there were modifications of the upper incisors.

**CONCLUSION:** There is an inverse relationship between nasolabial angle and upper incisor position (the farther distance between the upper incisor and the N<sup>^</sup>Po-Or, the smaller the nasolabial angle) as well as between nasolabial angle and upper incisor angle (more incisor torque results in a smaller nasolabial angle).

## SP 256

Thursday, June 17 - at 14:30

### INFLUENCE OF SELF-ESTEEM ON QUALITY OF LIFE IN CHILDREN SEEKING ORTHODONTIC TREATMENT

E De Baets<sup>1</sup>, H Lambrechts<sup>1</sup>, J Lemiere<sup>2</sup>, L Diya<sup>3</sup>, G Willems<sup>1</sup>, Department of <sup>1</sup>Orthodontics, <sup>2</sup>Child and Adolescent Psychiatry and <sup>3</sup>Interuniversity Institute for Biostatistics and Statistical Bioinformatics, Katholieke Universiteit Leuven, Belgium

**AIM:** There are children who complain about minor aesthetic orthodontic problems, while others with a severe malocclusion are not even aware of it. Does self-esteem (SE) contribute to differences in perception of orthodontic problems? The aim of this study was to investigate the possible role of SE as an influencing factor on oral-health-related quality of life (OHRQoL) in children seeking orthodontic treatment.

**SUBJECTS AND METHOD:** A cross-sectional study of 78 children (43 boys, 35 girls) aged 11 to 16 years. SE was assessed by the domain 'sense of dignity' of the Competence Perception Scale for Adolescents (Dutch translation of Harte's Self-Perception Profile). OHRQoL was scored with the use of the Child Perception Questionnaire [CPQ 11-14 4 domains, oral symptoms (OS), functional limitations (FL), emotional well-being (EW) and social well-being (SW) and total score]. The Index of Orthodontic Treatment Need [ION, Dental Health Component (DHC) and Aesthetic Component (AC)] was used to quantify treatment need. Multiple regression analysis was undertaken to evaluate the impact of SE on the relationship between orthodontic treatment need and OHRQoL. The results were obtained after correction for age and gender.

**RESULTS:** SE had a significant impact on the relationship between FL and the DHC of ION in the group of children with low SE values; OHRQoL was lower for those who needed treatment (DHC 4-5) compared with those who did not (DHC 1-3). However, for high SE values the reverse occurred in that children with a high treatment need experienced a better OHRQoL. For all the other analyzed relationships, except that between SW and the DHC of the ION, a similar pattern was found, although these interactions were not significant.

**CONCLUSION:** There appears to be no significant impact of SE on the relationship between OHRQoL and orthodontic treatment need, with the exception of the relationship between FL and the DHC of the ION. Longitudinal data in a larger sample will be of value to further elucidate the role of SE as a moderator.

### A SURVEY OF LEVELS OF DENTAL ANXIETY AMONGST INDIVIDUALS WITH CLEFT LIP AND PALATE

**K Al-Moammar**<sup>1</sup>, F McDonald<sup>1</sup>, P Haers<sup>2</sup>, T Newton<sup>1</sup>, 1King's College London Dental Institute and <sup>2</sup>South Thames Cleft Service, London, England

**AIM:** Dental fear is common in the general population and related to early experience of traumatic dental treatment. The prevalence of dental fear in cleft lip and palate (CLP) subjects is unknown but might be hypothesized to be elevated given the increased likelihood of dental or surgical treatment. The aim of this study was to assess the level of dental and medical setting anxiety among children with CLP.

**SUBJECTS AND METHOD:** One hundred and eighty two individuals with a CLP drawn from the South Thames Cleft Service Database were compared with a control group of 150 healthy individuals attending the Orthodontic Department, Guy's Hospital. The participants were identified at three ages: 5-10, 11-15 and 16-20 years. A postal questionnaire survey was undertaken for the individuals with CLP, while the questionnaires were completed by the healthy individuals at the orthodontic department. The questionnaires used were the Facial Image Scale (FIS) (Buchanan and Niven, 2002) and the Modified Dental Anxiety Scale (MDAS) (Wong *et al.*, 1998).

**RESULTS:** For all age groups, dental fear scores were higher for individuals with a CLP than for the control group: 5-10 year-old, mean FIS CLP = 2.17 (SD 1.04), controls = 1.39 (SD=0.53),  $P < 0.001$ ; 11-15 year-old, mean MDAS CLP = 21.56 (SD 6.80), controls = 16.40 (SD 4.35)  $P < 0.001$ ; and 16-20 year-old, mean MDAS CLP = 22.08 (SD 8.68), controls = 13.44 (SD 2.89)  $P < 0.001$ .

**CONCLUSION:** Individuals with a CLP have high levels of dental fear in comparison with their age-matched peers. Interventions should focus on prevention and management of fear in this group of patients.

### VERIFYING THE TRACING OF TRUE HORIZONTAL AND TRUE VERTICAL USING THE INTRACRANIAL REFERENCE PLANE

L Polma, **V Lomakina**, V Cheremisova, Department of Orthodontics, Moscow State University of Medicine and Dentistry, Russia

**AIM:** Any morphometric analysis is based on the construction of reference lines. The problem is which of them should be used for cephalometric measurements. The true horizontal (TH) and true vertical (TVL) lines presuppose accuracy in taking lateral cephalograms in the natural head position (NHP). The aim of this research was to determine which intracranial plane shows the best correlation with TH.

**MATERIALS AND METHOD:** Profile photographs and lateral cephalograms of 71 patients with malocclusions were taken in the NHP (25 patients had a Class I, 25 a Class II and 21 a Class III malocclusion). The following lines were constructed: Frankfort horizontal traced through the soft (FH1) and hard tissue (FH2) landmarks, TH, the TVL and the optical plane (Opt). The angles measured were: FH1/FH2; FH2/TH; FH2/Opt; Opt/TH; Opt/TVL and N-S/TH.

**RESULTS:** There was no statistically significant difference between the groups. Frankfort horizontal (FH1 and FH2) were not identical; the mean value of FH1/FH2 angle was  $5.87 \pm 1.02$  degrees. FH2 had a different position according to TH (FH2/TH =  $5.03 \pm 3.79^\circ$ ). Opt plane was almost parallel to TH (Opt/TH =  $0.14 \pm 0.35^\circ$ ). The average mean value of the OptL/TVL angle was  $89.86 \pm 0.35$  degrees and of N-S/TH angle  $-9.1 \pm 4.07$  degrees.

**CONCLUSION:** Application of Opt as the intracranial reference plane helps prevent errors in tracing TH and TVL.

### TREATMENT EFFECTS ON THE SOFT TISSUE PROFILE IN PATIENTS TREATED WITH THE HERBST APPLIANCE

L Polma, **V Lomakina**, V Cheremisova, Department of Orthodontics, Moscow State University of Medicine and Dentistry, Russia

**AIM:** To study soft tissue profile changes during Herbst appliance treatment of patients with distal occlusion.

**MATERIALS AND METHOD:** Profile photographs and lateral cephalograms of 36 patients aged 12-24 years with distal occlusion were taken in the natural head position; mean treatment time was 11 months. To prevent maxillary molar distalization and consolidation of the upper dental archwire, brackets were used. Standard cephalometric landmarks were used to construct the reference lines and measurements. The patients were divided into three groups according to the position of the maxillary incisors: group I, patients with incisor retrusion ( $U1/NL > 75^\circ$ ); group II, with a normal position of the incisors ( $75^\circ \geq U1/NL \leq 65^\circ$ ) and group III, with incisor protrusion ( $U1/NL < 65^\circ$ ).

**RESULTS:** There was a significant decrease of ANB angle ( $1.68 \pm 0.83^\circ$ ,  $P < 0.05$ ) while  $U1/NL$  angle increased ( $11.22 \pm 4.46^\circ$ ,  $P < 0.01$ ) in group I. Changes of C/sn/UL angle were not statistically significant ( $P > 0.05$ ). Furthermore that group of patients had a gl/sn/pg angle of  $-3.90 \pm 1.81$  degrees ( $P < 0.05$ ), while UL/sn/pg angle and UL-B increased ( $-32.03 \pm 13.33^\circ$ ,  $P < 0.01$ ;  $-1.29 \pm 0.65$  mm,  $P < 0.05$  correspondingly). In group II the decrease of ANB angle ( $5.11 \pm 0.70^\circ$  to  $3.50 \pm 0.58^\circ$ ,  $P < 0.05$ ) was typical, gl/sn/pg angle did not change, the increase of UL/sn/pg angle was not statistically significant ( $P > 0.05$ ).  $U1/NL$  angle decreased in group III ( $7.88 \pm 3.95^\circ$ ,  $P < 0.05$ ) but gl/sn/pg angle did not change ( $P < 0.05$ ) and the increase of UL/sn/pg angle was not statistically significant ( $P > 0.05$ ).

**CONCLUSION:** Treatment planning for patients with distal occlusion should take into consideration maxillary incisor position. Patients with maxillary incisor retrusion demonstrate a statistically significant improvement of aesthetic parameters during orthodontic treatment.

### QUANTITATIVE ASSESSMENT OF RELAPSE AFTER RAPID MAXILLARY EXPANSION

**L Vaida**<sup>1</sup>, M Pacurar<sup>2</sup>, A Porumb<sup>1</sup>, C Corega<sup>1</sup>, A Pirte<sup>1</sup>, <sup>1</sup>University of Oradea and <sup>2</sup>University of Targu Mures, Romania

**AIM:** Rapid maxillary expansion (RME) is a method of enlarging the upper dentoalveolar arch, primarily involving the jawbone mass, with possibilities of application in all subjects with a narrow arch associated with a lack of space, regardless of the existing dento-maxillary anomalies (Angle Class I, II or III malocclusion). The purpose of this research was to assess the degree of relapse present 6 and 12 months after the active phase of RME.

**SUBJECTS AND METHOD:** Fifty-two patients selected according to the following criteria: aged 10-18 years; use of the same disjunctor fixed on the first permanent molars and premolars or on the first primary molars; widening the arch by 0.9-1.2 mm using RME; activation of the screw once a day (for 4-5 weeks); a 3-month period with the disjunctor *in situ* and a 9-month retention period with a rigid transpalatal arch. To determine arch enlargement and the degree of relapse, intermolar distance was measured. The degree of relapse was assessed by comparing measurements taken 3 months after disjunction (when the transpalatal arch replaced the disjunctor) and at 6 and 12 months, after retention (transpalatal arch).

**RESULTS:** The average degree of relapse at 6 months was 2.8 mm and at 12 months 3.1 mm.

**CONCLUSION:** The degree of relapse after RME is significant, even with a retention period of 12 months (disjunctor + transpalatal arch), mainly due to the secondary effect of coronal vestibular expansion when applying RME.

**SP 261**

Thursday, June 17 - at 14:30

**EN MASSE RETRACTION OF ANTERIOR TEETH WITH THE BIDIMENSIONAL TECHNIQUE****A Bodini**, M Greco, A Giancotti, Roma-Isola Tiberina-Ospedale Fatebenefratelli-Università Tor Vergata, Italy

**AIM:** The bidimensional technique emphasizes the three-dimensional (3D) control of maxillary incisors and space closure in extraction treatment by sliding mechanics, but needs double steps of the retraction phases (first the canines then the incisors). This presentation demonstrates the 3D control during en masse retraction (one single step) with sliding mechanics using the bidimensional technique in extraction treatment.

**SUBJECTS AND METHOD:** Sixteen adolescent patients (mean age  $13.9 \pm 1.42$  years) with a Class II division 1 malocclusion, normal skeletal pattern, increased overjet (7 mm) and no crowding in the lower arch treated with two upper first premolar extraction. Orthodontic treatment was carried out using traditional fixed orthodontic appliance with bidimensional prescription. The brackets consisted of two different slot sizes (0.018 X 0.025 inch on the anterior teeth and 0.022 X 0.028 inch on the lateral-posterior teeth). During *en masse* retraction a 0.018 X 0.022 inch stainless steel wire was used, with crimpable hooks distal to the canines, molar toe-in and accentuated curve of Spee, in order to manage torque control of the incisors. In the lateral segments the same archwire was undersized allowing easy sliding mechanics during space closure with a NiTi coil spring (300 g) extended from the first molar hooks to the archwire hooks. Posterior anchorage was controlled by means of a transpalatal bar. Panoramic and lateral radiographs were obtained for all patients at the beginning and end of treatment.

**RESULTS:** A Class I canine and Class II molar relationship was achieved in all patients. All patients showed bodily movement of the upper incisor with a good torque control evidenced on the cephalometric superimposition following Björk's method, and a mean variation of 1-ANS/PNS of 12 degrees.

**CONCLUSION:** The modified two-dimensional technique with *en masse* retraction on sliding mechanics using a 0.018 X 0.022 inch stainless steel wire and traditional brackets could be considered an effective and less time-consuming option in extraction treatment.

**SP 262**

Friday, June 18 - at 10:45

**IS THERE A RELATIONSHIP BETWEEN ORTHODONTIC AND PERIODONTAL TREATMENT NEED?****R Nalcaci**<sup>1</sup>, S Demirel<sup>2</sup>, O Sökücü<sup>3</sup>, F Öztürk<sup>4</sup>, A B Kocoğlu Altan<sup>4</sup>, Departments of Orthodontics, <sup>1</sup>Cumhuriyet University, Sivas, <sup>2</sup>Gaziantep University and <sup>4</sup>Inönü University, Malatya and <sup>3</sup>Department of Periodontology Kırıkkale University, Turkey

**AIM:** It is believed that an orthodontic malocclusion is one factor in the aetiology of periodontal disease. With orthodontic treatment the periodontal health may improve because of the ideal occlusion. The aim of this study was to determine the relationship between orthodontic and periodontal treatment need.

**SUBJECTS AND METHOD:** Eighty hundred and sixty children (384 males, 452 females) aged between 11-14 years. To ensure data from different socio-economic standards, the schools in Sivas divided into three demographic categories as: high, middle and low socio-economic standards. Using random sampling technique, seven of the 67 schools were selected representing a cross-section of socio-economic status. Eight hundred and sixty of 20781 students were also selected with a random sampling technique. Subjects who had previous orthodontic treatment were excluded from the examination. Four experienced orthodontists and two periodontist performed the clinical examinations. Treatment Priority Index (TPI) scores were used for determining the severity of malocclusion. For determining periodontal status, the Community Periodontal Index of Treatment Needs (CPITN) scores

were used. Associations between variables were analyzed by chi-square test.

**RESULTS AND CONCLUSION:** Thirty-six per cent of the students had normal occlusion, 41.3 per cent had a slight, 15.9 per cent a definite, 4.1 per cent a severe, and 2.7 per cent a very severe malocclusion. The outcome of this study revealed that there was no relationship between orthodontic treatment need and CPITN scores ( $\chi = 19.22$ ,  $P = 0.257$ ,  $P > 0,05$ ).

**SP 263**

Friday, June 18 - at 11:30

**ALLERGIES IN ORTHODONTICS – FACTS AND MYTHS****G Stefanović**<sup>1</sup>, L Vukosavljević<sup>2</sup>, N Stefanović<sup>1</sup>, <sup>1</sup>University of Belgrade, Serbia and <sup>2</sup>Private Dental Office, The Hague, Netherlands

**AIM:** Allergies are a rare occurrence in the orthodontic office. However, widespread use of nickel in cosmetic jewellery and body piercing, as well as the use of new materials, such as latex, has resulted in an increase in allergic occurrences among the general population and subsequently in orthodontic patients. The aim of this study was to review the major sources of allergens in the orthodontic practice and propose alternatives for a safer treatment and working environment.

**MATERIALS AND METHOD:** Literature review was performed using the PubMed and KoBSON websites. Ten articles and several book chapters were analyzed.

**RESULTS:** According to the literature review, allergic reactions to orthodontic materials are classified as Type I and Type IV (hypersensitivity). Both types of allergic reactions require several prolonged contacts with allergens. Dental materials that could act as allergens are metals, polymers and latex. Materials causing Type IV allergic reactions contain polymerized materials and/or catalysts, and are most commonly used in prosthodontics, orthodontics and restorative dentistry. Materials containing latex, which are used in orthodontics, were shown to be strong inductors of Type I allergic reactions.

**CONCLUSION:** Toxicity, allergenic potential and biocompatibility vary in different materials. Metals that are most commonly suspected to cause allergic reactions during orthodontic treatment are nickel and chrome. In patients allergic to those metals, arches containing nickel and/or chrome should be avoided or covered with plastic protectors. Increased allergenic potential in contemporary orthodontic practice could also be attributed to latex. Intraoral elastics and elastic ligatures that contain latex have relatively less potential to cause allergies than extraoral elastics. In such cases non-latex elastics should be used. Another problem is sensitisation of orthodontists and their assistants to latex. Special attention should be paid to such cases, since latex is the only material used in orthodontic practice that could cause anaphylactic shock.

**SP 264**

Friday, June 18 - at 11:30

**COMPARISON OF PLAQUE REMOVAL OF TWO POWER BRUSH HEADS VERSUS A MANUAL TOOTHBRUSH****C Erbe**<sup>1</sup>, M Klukowska<sup>2</sup>, I Tsaknaki<sup>1</sup>, H Timm<sup>3</sup>, H Wehrbein<sup>1</sup>, <sup>1</sup>Department of Orthodontics, University Medical Centre of the Johannes Gutenberg-University, Mainz, Germany and Procter & Gamble Company, <sup>2</sup>Mason, Ohio, USA and <sup>3</sup>Kronberg, Germany

**AIM:** To evaluate the plaque removal efficacy of a power toothbrush with an orthodontic brush head compared with a regular brush head and a manual toothbrush in orthodontic patients.

**SUBJECTS AND METHOD:** All 45 patients who completed the study had multibracket-appliances in the upper and lower arch. This study was a randomized, operator-blind, six-period crossover design, which examined plaque removal of three toothbrushes: Oral-B® Triumph™ 5000 with an Ortho brush head or a FlossAction™ brush head and an ADA manual toothbrush. Digital plaque imaging analysis (DPIA) was used to measure the pre- and post-

brushing plaque level. The scores were analyzed for group differences using a mixed model ANCOVA.

**RESULTS:** The mean age of the subjects was  $14.6 \pm 2.4$  (minimum 12 years, maximum 25 years). The average baseline DPIA scores were 46.5 per cent for the Triumph™ with Ortho head, 45.5 per cent for Triumph™ with the FlossAction™ head, and 46.1 per cent for the ADA manual brush, which were not statistically different from each other ( $P = 0.856$ ). The plaque reduction for the power brush with orthodontic brush head was 59.5 per cent and with the regular brush head 55.9 per cent. For the manual toothbrush plaque reduction was 47.0 per cent ( $P < 0.001$ ). The Triumph™ brush with the Ortho or FlossAction™ brush head was statistically superior to the ADA brush ( $P < 0.001$ ). The power brush with an orthodontic brush head showed statistically higher plaque reduction than the power brush with a regular brush head ( $P = 0.007$ ).

**CONCLUSION:** Both power brush/brush head combinations were found to remove statistically significantly more plaque compared with a manual toothbrush in patients with fixed appliances.

## SP 265

Friday, June 18 - at 11:30

### ROOT RESORPTION AFTER ORTHODONTIC TREATMENT

**E Kowalska**, P Pietrzak, Department of Orthodontics, University of Lodz, Poland

**AIM:** Resorption is defined as a loss of alveolar bone and hard tissues of the teeth, due to certain physiological or pathological factors. The most frequent and severe resorption is observed in the upper and lower anterior teeth. Roots which are prone to resorption are: dilacerated, bottle shaped and pointed. Treatment for a period of 3 years and a Class III malocclusion increase the risk of resorption. The aim of this study was to analyse the scale of resorption in patients, treated with fixed appliances. An attempt was also made to evaluate the causes of this phenomenon.

**MATERIALS AND METHOD:** Dental pantomograms (DPTs) of 60 patients aged 8 to 32 years, treated with fixed appliances. The extent of resorption was measured by comparing the root length on DPTs taken before and after treatment. Root length was measured from the radiological apex to the cemento-enamel junction parallel to the long axis of the tooth.

**RESULTS:** Resorption was observed on the roots of the upper and lower anterior teeth in 28 patients, most frequently on the roots of the upper lateral incisors. The greatest resorption occurred on the lower right medial incisor and the least on an upper right lateral incisor. The phenomenon of resorption was independent of gender. The risk of resorption increased with age. The average treatment time of patients in whom resorption occurred, was 2.9 years. Teeth with an atypical root shape (pointed, curved and narrow) predisposed to resorption. Extraction and Class II elastics increased the risk of resorption.

**CONCLUSION:** Root shape, type of malocclusion and patient's age should be taken into consideration when planning orthodontic treatment, using continuous forces.

## SP 266

Friday, June 18 - at 11:30

### PHOTOGRAPHIC PROFILE ANALYSIS OF PATIENTS WITH PIERRE-ROBIN SEQUENCE

**A Ifert**<sup>1</sup>, R Sader<sup>2</sup>, F Ifert<sup>1</sup>, S Kopp<sup>1</sup>, Department of <sup>1</sup>Orthodontics and <sup>2</sup>Oral, Maxillofacial and Plastic Surgery, Goethe University Frankfurt, Germany

**AIM:** Pierre-Robin sequence is characterized as a congenital combination of malformations of micrognathia, glossoptosis and a median cleft palate (60 to 80%). Every 8,000 to 20,000 births is a baby with Pierre-Robin sequence. The patients often suffer from obstructions in the respiratory tract. Newborns with Pierre-Robin sequence show a typical triad. The intention of this study was to compare growth of the mandible of patients with Pierre-Robin sequence during their first year of life.

**SUBJECTS AND METHOD:** Using the orthodontic planning software, Onyx Ceph3® for photographic profile analysis, six newborns with Pierre-Robin sequence were examined. All were treated with a modified upper plate with an individual pharyngeal spur. The babies were monitored to check their oxygen saturation during treatment. Photographs were taken before treatment with upper plate (T1), 4 months after insertion of the plate (T2) and after palatal surgical closure (T3). For the profile analysis, the angle between glabella (Gl), labiale superius (Ls) and soft pogonion (Pog) was measured.

**RESULTS:** The changes of the profile were statistically significant between the three examinations. Between the T1 and T2 the angle of Gl-Ls-Pog increased on average by 13.25 degrees. Between T1 and T3, Gl-Ls-Pog angle changed, on average, by 36.45 degrees.

**CONCLUSION:** Conservative orthodontic treatment of Pierre-Robin sequence patients with a modified upper plate resulted in a major change in the profile. There was a positional change of the retrognathic mandible. Facial growth of these patients, especially of the lower jaw, should be controlled routinely over a longer time period.

## SP 267

Friday, June 18 - at 11:30

### EVALUATION OF THE CRANIAL BASE ANGLE IN ACHONDROPLASTIC DWARFS – A CEPHALOMETRIC ANALYSIS

**M de Mezer**<sup>1</sup>, Z Krysiński<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Prosthodontics, University of Medical Sciences Poznań Poland

**AIM:** Achondroplasia is the most common cause of dwarfism. This condition is genetically based, autosomal, dominant with full penetration. It affects 1 in 15000 to 30000 live births. Almost every case of achondroplasia is associated with a mutation of FGFR-3. Short stature is one of the most characteristic features of achondroplasia. Skull changes include hypoplasia of the jaw bones, enlargement of the cerebral skull and closing of cranial base angle. Cranial base growth affects the relationships between the maxilla and mandible. The aim of this study was to determine measurements of the cranial base angle of achondroplastic dwarfs.

**MATERIALS AND METHOD:** Cephalograms of 18 achondroplastic dwarfs. Achondroplasia was genetically confirmed for every patient. Two cephalometric parameters were analyzed, N-S-Ba and N-S-Ar angle, which describe skull base flexure. The mean values and standard deviations for these angles were calculated, both for the whole group and separately for the genders.

**RESULTS:** The mean values for the angles were 128 degrees ( $\pm 9.1^\circ$ ) for S-N-Ba and 117.8 degrees ( $\pm 10.5^\circ$ ) for S-N-Ar. Differences in the mean angle values between genders were observed. For females these values were 126.7 degrees ( $\pm 9.5^\circ$ ) for N-S-Ba and 115.3 degrees ( $\pm 11.2^\circ$ ) for N-S-Ar and for males 129.6 degrees ( $\pm 8.8$ ) and 120.8 degrees ( $\pm 9.2^\circ$ ), respectively.

**CONCLUSION:** Cranial base angle in achondroplasts is within the normal range or slightly above. Further research is required to determine other factors that might be involved in the characteristic facial profile of achondroplastic dwarfs.

## SP 268

Friday, June 18 - at 11:30

### FAMOTIDINE AFFECTS HISTOMORPHOMETRIC AND BIOCHEMICAL PARAMETERS DURING TOOTH MOVEMENT

**A Meh**<sup>1</sup>, Š Sprogar<sup>2,3</sup>, J Marc<sup>4</sup>, G Drevenšek<sup>2</sup>, M Drevenšek<sup>1</sup>, <sup>1</sup>Department of Orthodontics, University Medical Centre <sup>2</sup>Institute of Pharmacology and Experimental Toxicology and <sup>4</sup>Department of Biochemistry, University of Ljubljana and <sup>3</sup>Ort-line, Institute for Orthodontics and General Dentistry, Ljubljana, Slovenia

**AIM:** Histamine is involved in bone modelling, which is the key process during orthodontic tooth movement. Osteoclasts, which resorb bone, and osteoblasts, the bone-forming cells, both express H2 receptors. H2 receptor antagonists are widely used in the treatment of gastrointestinal ulcers. The aim of the present

study was to determine the effects of famotidine, an H2 receptor antagonist, on bone modelling during orthodontic tooth movement in rats.

**MATERIALS AND METHOD:** Three groups of Wistar rats, a control group (n = 16), an appliance only group (n = 16) and a famotidine group (n = 16). Animals in the last two groups were fitted with a superelastic closed coil spring (force = 25 cN) between the upper left first molar and the upper incisors. Animals of the control group and the appliance only group were treated daily with saline, and those of the famotidine group with 10 mg kg<sup>-1</sup> of famotidine. Tooth movement was measured weekly from day 0 to day 42. The animals of all groups were sacrificed on day 42 and tissue samples were prepared for further analysis. Gene expression levels for bone turnover markers, cathepsin K and osteocalcin, were determined by means of RT-polymerase chain reaction. Alveolar bone, osteoblast and osteoclast volumes were determined histomorphometrically.

**RESULTS:** Famotidine decreased the amount of tooth movement from day 35 onwards ( $P < 0.01$ ) and also decreased osteoclast volume. An increase in osteoblast volume ( $P < 0.01$ ), osteoblast activity ( $P < 0.01$ ) and osteoclast activity ( $P < 0.01$ ) was observed in the famotidine group compared with the appliance only group. No statistically significant difference was observed in alveolar bone volume between the famotidine and the appliance only groups.

**CONCLUSION:** Famotidine affects bone resorption and bone formation processes during orthodontic tooth movement. The observed decrease in tooth movement in the late stage of orthodontic tooth movement is probably due to the reduction of osteoclasts.

## SP 269

Friday, June 18 - at 11:30

### RADIOGRAPHIC ASSESSMENT OF THE PRESENCE OF THIRD MOLAR GERMS IN MACEDONIAN CHILDREN

**G Curcievea-Cuckova**<sup>1</sup>, H Pendov<sup>2</sup>, M Maneva<sup>1</sup>, A Mojsoska<sup>2</sup>, L Jolevski<sup>2</sup>, <sup>1</sup>Department of Orthodontics, P.H.O. University Dental Clinical Center, Skopje and <sup>2</sup>Private Practice, Ohrid, Former Yugoslav Republic of Macedonia

**AIM:** To investigate the stages of calcification and eruption times of various teeth, especially third molars, in Macedonian children, correlating dental and chronological age.

**MATERIALS AND METHOD:** Dental panoramic radiographs of 1000 subjects (527 males, 473 females) born between 1985 and 2005 who were between 5 and 16 years of age at the time a dental pantomogram (DPT) were obtained. All children were placed in the age group closest to their chronological age. Calcification of the each tooth on the left side of the mandible was rated in eight stages of calcification, A to H, according to the method described by Demirijan.

**RESULTS:** Significant differences were found between chronological and dental age. The findings showed premature eruption times in girls, but delayed calcification in boys compared with Demirijan's stages of calcification. The earliest appearance of visible signs of lower third molars (start of follicle development) was found at 7.4 years for both genders. The latest appearance was found at 10.6 years in girls, and 13.2 years in boys.

**CONCLUSION:** Radiographic assessment of calcification times of the permanent dentition, including the third molars, is of importance.

## SP 270

Friday, June 18 - at 11:30

### DENTAL ANOMALIES IN CHILDREN REGISTERED WITHIN THE CLEFT PALATE CENTRE OF STRASBOURG

**O Matern**<sup>1</sup>, F Perrin-Schmitt<sup>2</sup>, E-A Sauleau<sup>3</sup>, W Bacon<sup>1</sup>, B Grollemund<sup>1</sup>, <sup>1</sup>Functional Unit of Dentofacial Orthopaedics and <sup>3</sup>Public Health Department, University Hospital, and <sup>2</sup>Inserm U977, Faculty of Medicine, Strasbourg, France

**AIM:** In the general population, the prevalence of hypodontia (excluding third molars) has been reported to range from 3 to 10

per cent. Dental anomalies have been found to be more prevalent in subjects with a cleft lip, with or without palate (CL/P) than in those without a CL/P. The aim of the study was to determine the frequency of these anomalies.

**SUBJECTS AND METHOD:** Alsatian children with CL/P registered within the Cleft Palate Centre of Strasbourg, France. Cleft types and dental anomalies (missing, supernumerary or peg-shaped teeth) were assessed clinically and radiographically. Exclusion criteria were: patients under 5 years of age and absence of or unusable panoramic radiograph. The investigation included records of 109 patients with CL/P, 79 boys and 30 girls (mean age: 12 years).

**RESULTS:** Fifty-nine per cent of the subjects had evidence of hypodontia. Cleft types were divided into four groups based on the extent of clefting: [cleft lip only (CLO) 13.8%, cleft lip and alveolus (CLA) 3.7%, CL/P (49.5%) and cleft palate only (CPO) 33%]. There were significantly more dental anomalies associated with CPO and CLP than with CLO. The maxillary lateral incisors and upper and lower premolars were the most frequently missing teeth. Eleven patients had supernumerary teeth in the region of the cleft, and 15 teeth were considered as being peg-shaped (maxillary lateral incisors were the most affected teeth). All these anomalies were found in proportionately higher frequencies as the severity of the cleft increased. A male dominant tendency was evident for all cleft types ( $P < 0.0001$ ) as well as a left side predominance for hypodontia ( $P = 0.0015$ ) whatever cleft sidedness.

**CONCLUSION:** A statistically significantly higher prevalence of hypodontia was found for this group of French children with CL/P than in the general population. A significant dominant tendency of hypodontia was observed on the left side irrespective of cleft side. It seems that the same environmental and genetic factors are related to tooth development and cleft genesis.

## SP 271

Friday, June 18 - at 11:30

### VITAMINS B6, B12 AND HOMOCYSTEINE IN CLEFT LIP AND PALATE: BIOCHEMICAL ANALYSIS

**A T Altug**<sup>1</sup>, E Ozdiler<sup>1</sup>, H Tastan<sup>1</sup>, Ö Sancak<sup>1</sup>, E Derelli<sup>2</sup>, <sup>1</sup>Department of Orthodontics, School of Dentistry and <sup>2</sup>Faculty of Biology, School of Science, Ankara University, Turkey

**AIM:** The vitamins B6 and B12 are essentially involved in the methylation cycle. Homocysteine is also an important metabolic parameter in the methylation cycle whose increase in tissue indicates a reduced methylation rate. A lack or deficiency of B6 and B12 and elevated homocysteine levels are thus considered a cause of malformations. However, evidence linking low levels of B group vitamins and oral clefting is presently equivocal. The aim of this pilot study was to investigate associations between non-syndromic oral clefts and biochemical measures of B6 (pyridoxine hydrochloride), B12 (cyanocobalamin) and homocysteine status.

**SUBJECTS AND METHOD:** A consecutive sample of 37 children (8 girls, 29 boys) born with non-syndromic cleft lip, with or without a cleft palate. Homocysteine, B6 and B12 concentrations in the blood were analyzed. They were studied with high-performance liquid chromatography and electro-chemiluminescence immunoassay.

**RESULTS:** According to age-adjusted ranges, 20 patients had increased, six decreased and 11 normal ranges of homocysteine. In one patient, B6 was decreased and in another B12 was decreased. Ten patients had B12 values close to the lower border of the normal range. Eighteen subjects had increased B6 values.

**CONCLUSION:** Due to the limited number of cleft patients included in the present study, no significant decrease in the levels of B6 and B12 was observed in blood. A slight lack of B12 in conjunction with an increase in homocysteine levels could be considered as a risk factor for non-syndromic orofacial clefting. As an increase in B group vitamins has been proven to cause a decrease in homocysteine levels, pregnant women should be informed about vitamin supplements and a well-balanced diet.

### THREE DIMENSIONS OF THE MAXILLA IN UNILATERAL VERSUS BILATERAL CANINE IMPACTION

Z Uiku<sup>1</sup>, D Luk<sup>2</sup>, <sup>1</sup>Division of Orthodontics, College of Dental Medicine, Columbia University and <sup>2</sup>Private practice, Brooklyn, New York, USA

**AIM:** The canines play an integral role in functional occlusion and an aesthetic smile. Both of these are compromised by impactions. Although maxillary width has been investigated in association with canine impaction, no comparison has been made between unilateral versus bilateral groups. The aim of this study was to compare positional variation, length and arch width of the maxilla in unilateral versus bilateral impaction groups.

**SUBJECTS AND METHOD:** A search of all orthodontic clinic records at Columbia University was performed. Subjects were selected with the criteria of a minimum age of 12.5 years, unerupted maxillary canines, no previous orthodontic treatment and non-syndromic patients or those with congenital defects. The age range for the unilateral (66 subjects) and bilateral (35 subjects) groups were from 12.5-28.5 years and 12.6-26.8 years, respectively. The comparison group with a mild Class I malocclusion (83 subjects) ranged from 12.5-21.2 years. Statistical analysis consisted of a two-sample t-test.

**RESULTS:** The unilateral versus bilateral group was not statistically different when assessed in the transverse dimensions. The difference was -1.2 mm in premolar width, and -0.6 mm in molar width. Maxillary position or length was not significantly different in either group and was similar to the non-impaction Class I group. Since the sagittal relationship of the maxilla in the canine impaction groups was very similar to the norms, the use of headgear in the treatment of canine impaction, as suggested in the literature, should be carefully considered in the absence of other indications.

**CONCLUSION:** Unilateral and bilateral impaction patients do not differ in anterior or posterior arch width. Absence of over-retained primary canines in unilateral impaction patients may cause a slight collapse of premolar width. Maxillary position and transverse dimension do not appear to influence the presence of either unilateral or bilateral canine impaction.

## SP 273

Friday, June 18 - at 11:30

### THE BENDING LOAD CAPACITIES OF RAPID MAXILLARY EXPANSION SCREWS – A PILOT STUDY

A P Muchitsch<sup>1</sup>, B Wendl<sup>1</sup>, H Winsauer<sup>2</sup>, M Pichelmayer<sup>1</sup>, M Payer<sup>3</sup>, <sup>1</sup>Department of Orthodontics, Medical University Clinic of Graz, <sup>2</sup>Private practice, Bregenz and <sup>3</sup>Technical University of Graz, Austria

**AIM:** In order to apply high, short-term, forces during rapid maxillary expansion (RME) to the medial palatal suture without causing unwanted side-effects (dentoalveolar tipping etc.), the appliances used should be as rigid as possible. Since the retention arms of the jack screws are vulnerable, stressed weak points, it was the aim of this study to investigate their bending strength.

**MATERIALS AND METHOD:** The retention arms of 16 RME screws comprising four arms and one with eight arms, were examined using a three-point bending test. The screws were classified according to their ability to absorb the applied bending loads, indicated by the parameters force (F), bending stress ( $\sigma$ ) and ductility (ffmax).

**RESULTS:** Fifteen retention arms (stainless steel), despite having the same diameter (1.48-1.49 mm), differed up to 69.81 per cent between the highest (288.0 N) and lowest (124.5 N) force parameters (F), and up to 66.40 per cent between the highest (3325.9 N/mm<sup>2</sup>) and lowest (1464.0 N/mm<sup>2</sup>) bending stress parameters ( $\sigma$ ). Due to optimum formability, though reduced rigidity, a titanium screw for nickel sensitive patients, displayed the lowest force and bending tension values. The stainless steel double arms of the 8-arm screw device displayed the highest force data. The mean

ductilities (ffmax) of the screws with the most and least rigid single steel arms differed by 22.77 per cent. Statistical analysis, using the Pearson correlation coefficient, revealed a significant indirect correlation between ductility and both maximum force ( $r = -0.780$ ,  $P < 0.001$ ) and maximum bending stress ( $r = -0.778$ ,  $P < 0.001$ ).

**CONCLUSION:** Despite having the same cross-sectional diameters, the single retention arms of all tested stainless steel expansion screws displayed variable loading capacities. Clinically screws with lower rigidity appear acceptable for RME during the pre-pubertal age, while in adolescence and early adulthood screws with higher loading capacities seem advisable.

## SP 274

Friday, June 18 - at 11:30

### RELATIONSHIP BETWEEN DELETERIOUS ORAL HABITS AND THE DEVELOPMENT OF MALOCCLUSION

A S Ladislau<sup>1</sup>, H A Barbosa<sup>1</sup>, S N David<sup>2</sup>, K Faltin Jr.<sup>1</sup>, C F Ortolani<sup>1</sup>, <sup>1</sup>Paulista University and <sup>2</sup>Cruzeiro do Sul University, São Paulo, Brazil

**AIM:** Several epidemiological studies have shown a high incidence of occlusal disharmonies in the Brazilian child population. Among the aetiological malocclusion factors, are deleterious oral habits such as mouth breathing, abnormal function of the tongue, and finger, pacifier and bottle sucking. Thus, the purpose of this cross-sectional study was to evaluate malocclusion development and its relationship with the presence of oral habits.

**SUBJECTS AND METHOD:** One thousand school children aged between 6 and 10 years. A questionnaire related to the child's health and the presence of habits was completed by the parents. On examination, data were obtained regarding occlusion and stomatognathic system functional alterations.

**RESULTS:** There was a significant statistical association between deleterious oral habits and malocclusion. It was found that habits can cause morphological changes resulting in the development of a malocclusion.

**CONCLUSION:** Oral habits are potential aetiological factors in the alteration of occlusion and the change in normal growth pattern.

## SP 275

Friday, June 18 - at 12:15

### INFLUENCE OF PACIFIER NECK DIMENSIONS ON FLEXIBILITY AND INTRAORAL FORCE DISTRIBUTION

M Bertl, H-P Bantleon, Department of Orthodontics, Medical University of Vienna, Austria

**AIM:** A pacifier may be seen as a child's first oral appliance, yet little research is available on the ideal properties that will facilitate natural dentoalveolar development. The aim of the study was to test the influence of pacifier neck dimensions on its flexibility and ability to allow for desirable intraoral force distribution.

**MATERIALS AND METHOD:** Thirteen different silicone pacifiers. The precise neck dimensions were obtained using an optical measuring method. Five samples of each pacifier were subsequently placed in a measuring machine which consisted of three sensors, that could be independently adjusted and measure force and distance in two dimensions. Two of the sensors represented the upper and lower dentition and enclosed the pacifier's neck with a defined force of 200 cN. The third sensor was used to apply force to and thereby elevate the head of the pacifier in place of the tongue.

**RESULTS:** Vertical neck dimensions ranged from 2.28 to 10.15 mm. A mean of  $46.55 \pm 17.08$  cN was required to lift the pacifier's head 2.6 mm. At this position, Pearson coefficient showed a strong and highly significant correlation ( $r = 0.882$ ;  $P < 0.01$ ) between the applied force and the vertical diameter of the pacifier neck. For linear regression, the coefficient of determination,  $r^2$ , was calculated at 0.778. Similar results were obtained after further elevation to 5 mm  $r = 0.845$  ( $P < 0.01$ );  $r^2 = 0.714$ .

**CONCLUSION:** There is a strong influence of the neck dimensions of a pacifier on overall flexibility. A more flexible pacifier should

provide less resistance to the natural position of the tongue and also less undesirable force on the alveolar processes and dentition during their development. While a thin neck should therefore be recommended, other parameters such as acceptance and durability also have to be considered in an ideal pacifier design.

### SP 276

Friday, June 18 - at 12:15

#### THREE-DIMENSIONAL MOVEMENT OF THE MAXILLARY FIRST MOLAR WITH THE QUADHELIX APPLIANCE\*\*\*

**K Terada**, N Sano, Y Hasegawa, Department of Orthodontics, The Nippon Dental University at Niigata, Japan

**AIM:** To investigate three-dimensional (3D) movement of the maxillary first molar with a quadhelix appliance.

**MATERIALS AND METHOD:** Fifty-two dental models obtained before and after treatment of 26 patients treated with the quadhelix appliance in two orthodontic offices. Surface images of the models were obtained using a laser scanner (Vivid 9i, Konica Minolta, Tokyo, Japan). The reference plane (X-Y plane) on each surface image was at the anterior point of the incisive papilla and the top of right and left mesio-lingual cusps of the maxillary first molar. The mesio-lingual cusp of the maxillary first molar was set as the origin. 3D changes of the other cusps were analyzed during treatment.

**RESULTS:** The mesio-buccal cusp of the maxillary first molar was moved 0.58 mm distally, 0.81 mm buccally, and extruded 0.20 mm. For the disto-buccal and disto-lingual cusps movement was 0.12, 0.80, 0.15 and 0.48, 0.10, 0.041 mm, respectively. While the same movement pattern was observed, the amount of tooth movement was different in the two orthodontic offices. This may have been due to the different treatment objectives

**CONCLUSION:** The maxillary first molars were inclined buccally and rotated distally. The centre of rotation was observed around the mesio-lingual cusp.

### SP 277

Friday, June 18 - at 12:15

#### ROOT RESORPTION AFTER 2.5 AND 15 DEGREE BUCCAL ROOT TORQUE FOR 4 WEEKS: A MICRO-COMPUTED TOMOGRAPHIC STUDY

N Bartley<sup>1</sup>, T Turk<sup>2</sup>, **C Colak<sup>2</sup>**, S Elekdag-Turk<sup>2</sup>, M A Darendeliler<sup>1</sup>, <sup>1</sup>Discipline of Orthodontics, Faculty of Dentistry, University of Sydney, Australia and <sup>2</sup>Department of Orthodontics, Faculty of Dentistry, University of Ondokuz Mayıs, Samsun, Turkey

**AIM:** Root resorption is an undesirable but inevitable consequence of orthodontic tooth movement. The severity is unpredictable and despite extensive research, the aetiology remains unknown. Torque has been acknowledged as a risk factor for root resorption. The aim of this study was to evaluate and quantify the extent of root resorption following the application of 2.5 and 15 degrees buccal root torque for 4 weeks.

**SUBJECTS AND METHOD:** Fifteen patients requiring extraction of bilateral upper first premolars for orthodontic treatment. Using a standardized experimental protocol, the right and left premolars were randomly subjected to either 2.5 or 15 degree buccal root torque. At the end of the four-week experimental period the premolars were extracted. A volumetric analysis of the extent of root resorption was performed using micro-computed tomography and measured using specially designed software.

**RESULTS:** Overall, the amount of root resorption was comparable after the application of 2.5 or 15 degree buccal root torque ( $P = 0.59$ ). A significant difference between the two force levels existed only at the apical region ( $P = 0.034$ ). More root resorption occurred in areas of compression than in areas of tension. The variables of age and gender were not statistically significant.

**CONCLUSION:** (1) Root resorption was evident after four weeks of buccal root torque application. (2) More root resorption was seen at the apical than at the middle and cervical regions. (3) Higher magnitudes of torque have the potential to cause more root resorption, particularly in the apical region.

### SP 278

Friday, June 18 - at 12:15

#### A CLINICAL STUDY OF THE DELAYED ERUPTION OF THE MAXILLARY FIRST MOLAR\*\*\*

**N Sano**, K Terada, Y Hasegawa, E Yajima, Y Terashima, Department of Orthodontics The Nippon Dental University School of Life Dentistry at Niigata, Japan

**AIM:** The first molar is the key of the occlusion. However few cases have been reported regarding congenitally missing and delayed eruption of the maxillary first molars. The aim of this study was to investigate the longitudinal formation of the maxillary first molars that were considered to be delayed in terms of eruption.

**MATERIALS AND METHOD:** Delayed eruption of the maxillary first molars was defined as a delay in formation 2 years later than average development. Longitudinal panoramic radiographs of 100 teeth from 53 subjects with delayed maxillary first molar eruption and without genealogical disease. The formation of the teeth were evaluated according to the method of Moorrees.

**RESULTS:** Forty-seven subjects (89%) had bilateral delayed maxillary first molar eruption; six (11%) were unilateral. Thirty-nine subjects (74%) had an Angle Class III, four (8%) an Angle Class I, and two (4%) an Angle Class II. Eighty-three teeth (83%) were erupted at R1/2 stage of Moorrees development, all teeth were erupted at R3/4.

**CONCLUSION:** The delayed maxillary first molars were erupted until R3/4 stage of Moorrees development. It is suggested that this investigation was useful for making the treatment plan in the mixed dentition.

### SP 279

Friday, June 18 - at 12:15

#### FINITE ELEMENT ANALYSIS OF MINI-IMPLANT AND PERIPHERAL BONE DURING UPPER CANINE RETRACTION

**I Bavbek Karakaya**, N Darendeliler, Department of Orthodontics, Faculty of Dentistry, Gazi University, Ankara, Turkey

**AIM:** To investigate the stress distribution in mini-implants inserted in the maxilla and supporting bone using finite element analysis (FEA) during maxillary canine retraction.

**MATERIALS AND METHOD:** The computerized tomography image of a patient was utilized to obtain a three-dimensional model of the maxilla for FEA. In the model, three different mini-implants: Neo-anchor plus (1.6 X 8 mm), Dewimed (1.6 X 8 mm) and Spider (2 X 8 mm) screws and Ormco brackets were employed. Mini-implants were positioned between the roots of the second premolar and first molar and 6 mm gingival to the apical bone crest. The force applied for maxillary canine retraction was taken as 200 g. Von Misses stress distributions in the mini-implants, the surrounding bone and the canine were evaluated to determine the effect of the screws.

**RESULTS:** FEA showed that maximum von Misses stress values obtained in the cortical bone were 2.9, 3.4 and 4.7 MPa for Neo-anchor plus, Dewimed and Spider screws, respectively. The corresponding stress values within the mini-implants were 7.7, 12.0 and 5.3 MPa, respectively.

**CONCLUSION:** Lower stress values occur in the cortical bone when smaller diameter screws are used. However, the stress values decrease further if a screw that has smaller pitch length in the region which is in contact with the cortical bone is inserted.

### SP 280

Friday, June 18 - at 12:15

#### NOSE CHANGES FOLLOWING BIMAXILLARY SURGERY IN TURKISH SUBJECTS WITH CLASS III MALOCCLUSIONS

H Gokalp, **F Aydin Mustafa**, A Bahat Yalvac, Development of Orthodontics, Faculty of Dentistry, University of Ankara, Turkey

**AIM:** Skeletal Class III dentofacial deformities may be the result of mandibular prognathism and/or maxillary deficiency. Severe skeletal Class III deformities are treated by orthognathic surgery in adulthood. Besides hard tissue changes, soft tissue changes are also observed after orthognathic surgery. The purpose of this

study was to determine changes in the soft tissue nose profile in Turkish Class III patients, after maxillary orthognathic surgery.

**SUBJECTS AND METHOD:** Nine Class III patients treated by bimaxillary orthognathic surgery (mean age 19.93, 7 females, 2 males) while the control group consisted of 11 Class III subjects with mandibular prognathism (mean age 19.45, 10 females, 1 male). Conventional lateral cephalometric radiographs were used to evaluate changes after surgery. Steiner and McNamara cephalometric analyses and nose measurements were performed (N = -Pr, Pr-N = Sn, Ac-Pr, hump, NLA, NBA, NMA, SFC, Dconv, Cconv, NboneL, NboneA). The obtained data were analyzed statistically by paired t- and Student's t-tests.

**RESULTS:** Statistically significant differences were found in the treated group before or after surgery for the following measurements: Pr-N = Sn (nasal depth 1), Ac-Pr (nasal depth 2) ( $P < 0.05$ ), NMA (naso-mental angle) ( $P < 0.01$ ), NBA (nasal base angle) and D-conc (lower dorsum convexity) ( $P < 0.05$ ). With regard to SNA, Na Perp.-A ( $P < 0.001$ ), Co-A ( $P < 0.01$ ) ANB and CoGn-CoA, the measurements were statistically different ( $P < 0.01$ ). The remaining parameters were statistically insignificant. Comparison of the treatment and control group showed that Na Perp.-Pog, Hump and SFC (soft tissue facial convexity) measurements were statistically insignificant. D-conv ( $P < 0.01$ ) and all other parameters exhibited statistically significant differences ( $P < 0.001$ ).

**CONCLUSION:** Soft tissue nose profile changes are observed in Turkish Class III subjects after bimaxillary orthognathic surgery.

## SP 281

Friday, June 18 - at 12:15

### PERMANENT TOOTH-CROWN DIMENSIONS IN PREMATURELY BORN CHILDREN

**L Paulsson-Björnsson**, L Bondemark, Department of Orthodontics, Faculty of Odontology, Malmö University, Sweden

**AIM:** According to the literature, the scientific evidence is insufficient to answer the question as to whether premature birth causes altered tooth-crown dimensions. The aim of this study was to evaluate permanent tooth-crown dimensions in prematurely born children and to compare the findings with full-term born controls.

**SUBJECTS AND METHOD:** White pre-term children, 8 to 10 years of age, were selected from the Swedish Medical Birth Register. One group consisted of 36 extremely pre-term children (born before the 29<sup>th</sup> gestational week), and the other 37 very pre-term children (born during gestational weeks 29 to 32). The pre-term children were compared with a control group of 41 full-term children, who were matched for gender, age, nationality and living area. Clinical examinations were undertaken and study casts and panoramic radiographs were obtained for each child. The permanent maxillary and mandibular first molars, central incisors and laterals were measured with a digital sliding calliper on the study casts. The tooth-crowns were measured both mesiodistally and buccolingually. The examiner conducting the measurement analysis was unaware of the subject grouping.

**RESULTS:** The maxillary and mandibular first molars were smaller mesiodistally and buccolingually ( $P \leq 0.01$ ) whereas the central incisors and lower laterals were smaller mesiodistally ( $P \leq 0.02$ ) in the extremely pre-term group compared with the full-term group. A reduction in tooth size of 5-8 per cent was found between the extremely pre-term group and the full-term group. Furthermore, the maxillary first molars and mandibular left first molar were also smaller mesiodistally ( $P \leq 0.035$ ) in the extremely pre-term group compared with the very pre-term group.

**CONCLUSION:** Premature birth is associated with reduced tooth-crown dimensions of the permanent teeth, and the more pre-term the birth the smaller the tooth-crown dimensions.

## SP 282

Friday, June 18 - at 12:15

### DYNAMICS OF THE ERUPTION OF PERMANENT TEETH IN DIABETES TYPE I CHILDREN

**M Koselj**<sup>1</sup>, M Cilenšek<sup>1</sup>, A Meh<sup>2</sup>, T Battelino<sup>3</sup>, M Drevenšek<sup>1</sup>, Departments of Orthodontics <sup>1</sup>Faculty and <sup>2</sup>University Medical Center and <sup>3</sup>Department of Pediatric Endocrinology, Diabetes and Metabolism, University Medical Centre, University Children's Hospital, Ljubljana, Slovenia

**AIM:** Diabetes is one of the most widespread chronic diseases that can also affect oral health. The objective of this study was to determine the effect of diabetes type 1 on the dynamics of the eruption of permanent teeth in children in the mixed dentition period and its effect on the incidence of crowding.

**SUBJECTS AND METHOD:** Seventy-nine children (average age of  $10.8 \pm 1.8$  years): 40 randomly referred children (21 boys, 19 girls, average age  $10.7 \pm 1.8$  years) with diabetes type 1 and a control group of 39 healthy children (19 boys, 20 girls, average age  $10.8 \pm 1.7$  years). Children undergoing orthodontic treatment were excluded. The level of eruption of the teeth was assessed using study models. Each tooth was categorized into one of six levels of eruption, defined according to the data on the standard development of the jaw. Crowding was assessed based on study models measurements using the analysis of Droschl in the mixed dentition period, and the analysis of Lundström in children in the permanent dentition.

**RESULTS:** Children with type I diabetes were found to have accelerated tooth eruption in the period between phases 5 and 6, i.e. during the later eruption phase of teeth into the oral cavity ( $P = 0.049$ ). The dynamics of eruption did not influence crowding.

**CONCLUSION:** Eruption is accelerated in the later phases of eruption of a permanent tooth in the oral cavity in diabetes type I children. There is no effect on the crowding.

## SP 283

Friday, June 18 - at 12:15

### THE EFFECT OF DIABETES TYPE 1 ON ORAL HEALTH FACTORS IN CHILDREN IN THE MIXED DENTITION PERIOD

**M Cilenšek**<sup>1</sup>, M Koselj<sup>1</sup>, A Meh<sup>1</sup>, N Uršič-Bratina<sup>2</sup>, M Drevenšek<sup>1</sup>, <sup>1</sup>Department of Orthodontics, Faculty of Medicine, University of Ljubljana and <sup>2</sup>Department of Pediatric Endocrinology, Diabetes and Metabolic Disease University Children's Hospital, Ljubljana, Slovenia

**AIM:** Diabetes is a chronic disease that can also affect oral health. The objective of this study was to determine the effect of type 1 diabetes on some oral health factors in children in the mixed dentition period.

**SUBJECTS AND METHOD:** Seventy-nine children with an average age of  $10.8 \pm 1.8$  years. The study group included 40 randomly referred children ( $10.7 \pm 1.8$  years) with diabetes type 1. The control group comprised 39 healthy children (average age  $10.8 \pm 1.7$  years), who were comparable by gender and age with the study group. Children who were already undergoing orthodontic treatment were excluded. The quantity of plaque was assessed by drawing a probe on the surface of the tooth next to the gingivae. Plaque was categorized into levels 0 to 3. Bleeding was provoked by drawing the tip of the probe between the free gingiva and the tooth. The relationship of the dental arches was assessed using Angle's classification.

**RESULTS:** Children with type 1 diabetes had a higher incidence of plaque ( $P = 0.000$ ). Second and third degree plaque occurred more often in the study group, whereas zero and first degree plaque were more common in the control group ( $P = 0.000$ ). The children with type 1 diabetes had a higher incidence of provoked gingival bleeding ( $P = 0.000$ ). There was no statistically significant difference between the study and control group in the incidence of malocclusion according to Angle classification ( $P = 0.510$ ).

**CONCLUSION:** Diabetes type 1 affects the incidence of plaque and bleeding. This has to be taken into consideration when planning orthodontic treatment.

### BIOCOMPATIBILITY OF ORTHODONTIC BANDS AFTER EXPOSURE TO DENTAL PLAQUE

**S Hornikel<sup>1</sup>**, C Erbe<sup>1</sup>, I Schmidtman<sup>2</sup>, H Wehrbein<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Medical Biostatistics, Epidemiology and Informatics, Johannes-Gutenberg-University Mainz, Germany

**AIM:** Corrosion parameters of orthodontic alloys and appliances have frequently been assessed in vitro. Biocompatibility of orthodontic bands after in vivo use mimicking different oral hygiene conditions has so far been poorly investigated. The aim of this study was to assess the biocompatibility of orthodontic bands after exposure to human oral conditions, especially plaque accumulation, over the normal period of orthodontic treatment.

**MATERIALS AND METHOD:** Cell adherence and cell morphology of gingival fibroblasts grown on 32 orthodontic bands were tested. The bands had been seated intraorally from 6-37 months.

**RESULTS:** In areas without prior plaque attachment, cell adherence was found for 75.8 per cent of the bands. Cell morphology was evenly spread between spherical and elongated types. Areas with prior plaque attachment showed cell adherence for 83.9 per cent of bands. Cells were of a spherical morphology in 42.2 per cent and of an elongated morphology in 57.7 per cent of cases.

**CONCLUSION:** As no significant difference was observed in either cell adherence or cell morphology in areas with or without prior plaque attachment, it may be assumed that oral conditions during orthodontic treatment do not impair biocompatibility of orthodontic bands.

### RELATIONSHIP BETWEEN DENTAL MALOCCLUSION AND OTITIS MEDIA IN CHILDREN: A DESCRIPTIVE ANALYSIS

**M Fabbrizi**, V Luzzi, C Mirra, G Consoli, Department of Oral Science, Sapienza University of Rome, Italy

**AIM:** To examine the relationship between otitis media and the dentition.

**SUBJECTS AND METHOD:** Sixty-four children, aged 3 to 10 years, selected from those who at their first visit reported suffering from otitis. Their parents completed a questionnaire concerning pacifier use, thumb/finger-sucking, mouth breathing habits, method of feeding, day care attendance, and upper respiratory infections. All patients underwent an orthodontic clinical examination where overbite, crossbite and overjet and the presence of parafunctional oral habits such as atypical swallowing, mouth breathing and sucking of the inner lip were investigated. The collected data were analysed through descriptive analysis. Associations between otitis media and the presence of malocclusion were investigated. The employed tests were chi square and Fisher, with a significance level of 5 per cent.

**RESULTS:** In this study, 82.8 per cent of the children presented some type of malocclusion: anterior open bite (42.2%), increased overjet (42.2%), posterior crossbite (28.1%), deep bite (17.2%). In this group the most common factors were bottle feeding (92.2%), pacifier habit (67.2%), mouth breathing habit (35.9 per cent), finger sucking habit (28.1%), allergic rhinitis (21.8%), day care attendance (21.8%), asthma (3%). No statistically significant association between malocclusion and otitis was observed ( $P > 0.05$ ).

**CONCLUSION:** Although, the results indicate no significant association between malocclusion and otitis media, clinical-descriptive analysis was able to show a high incidence of malocclusion in children with otitis, the presence of risk factors common to both the development of otitis and malocclusion, and an improvement in the symptoms of otitis, as a result of interceptive orthodontic treatment.

### MAXILLARY ARCH WIDTH AND PALATAL DEPTH IN PATIENTS WITH MARFAN SYNDROME

**E Vahle-Hinz<sup>1</sup>**, Y von Kodolitsch<sup>2</sup>, B Kahl-Nieke<sup>1</sup>, H Korbmacher<sup>1</sup>, <sup>1</sup>Department of Orthodontics, Center of Dental and Oral Medicine and <sup>2</sup>Centre of Cardiology and Cardiovascular Surgery, Department of Cardiology, University Hospital Hamburg-Eppendorf, Germany

**AIM:** Marfan syndrome (MFS) is a connective tissue disorder with autosomal dominant inheritance that affects, inter alia, the musculoskeletal apparatus and also the hard palate. The aim of this study was to quantify the morphological changes of the palate in subjects with MFS compared with a control group.

**SUBJECTS AND METHOD:** Thirty-two adult patients (13 males, 19 females) with MFS aged between 18 and 60 years. Thirty-six adults without any syndromic diseases (15 males, 21 females) aged between 25 and 41 years served as the control group. Study casts were obtained for all subjects. The palates were analyzed by applying the following four parameters: maxillary interpremolar and intermolar widths, palatal depth and its ratio to the maxillary molar width according to Hsu. Intergroup differences were tested using the independent t-test.

**RESULTS:** Interpremolar and intermolar width showed highly statistically significant differences between the two groups. Interpremolar (MFS group 32.9 mm, control group 35.8 mm) and intermolar widths (MFS group 41.0 mm, control group 46.1 mm) were significantly reduced in the MFS group. The differences in palatal depth were also statistically significant. Patients with MFS showed an increased palatal depth in comparison with the controls (MFS 18.9 mm, control group 18.0 mm).

**CONCLUSION:** The connective tissue disorder in patients with MFS affects the interpremolar and intermolar width as well as palatal depth. All dimensions were statistically significantly reduced. The high-arched and small palate must be discussed as a co-factor in the increased clockwise rotation of the mandible that has been described in previous studies.

### DOES LOCATION OF CONGENITALLY MISSING TEETH AFFECT DENTOFACIAL MORPHOLOGY?

**A Y Gungor<sup>1</sup>**, H Turkkahraman<sup>2</sup>, Departments of Orthodontics, Faculties of Dentistry, <sup>1</sup>Mustafa Kemal University, Hatay and <sup>2</sup>Suleyman Demirel University, Isparta, Turkey

**AIM:** To find out whether or not the location of congenitally missing teeth affects dentofacial morphology.

**SUBJECTS AND METHOD:** One hundred and fifty four patients with two or more congenitally missing teeth. In the first part of the study, the patients were divided into three groups. Group 1 comprised 37 patients with missing teeth only in the maxilla, group 2, 32 patients with missing teeth only in the mandible and group 3, 85 patients with missing teeth both in the maxilla and mandible. In the second part of the study, patients were again divided into three groups. Group 1 consisted of 45 patients with missing teeth only in anterior region, group 2, 57 patients with missing teeth only in posterior region and group 3, 52 patients with missing teeth both in anterior and posterior regions. Fifty Class I patients without any missing teeth served as the controls in both parts of the study. Eighteen angular and 17 linear measurements were performed on lateral cephalograms. All recorded data were statistically analyzed and compared among the groups.

**RESULTS:** Significant differences were found in anterior and posterior face heights, ramus height and soft tissue measurements, and upper and lower incisor measurements according to localization ( $P < 0.05$ ). Patients with missing teeth only in maxilla had an increased upper lip-E plane length ( $P < 0.05$ ) while those with missing teeth only in mandible or only in anterior region had decreased posterior face heights ( $P < 0.05$ ). Patients with missing

teeth only in posterior region had a shorter ramus and anterior face height ( $P < 0.05$ ).

**CONCLUSION:** Localization of missing teeth significantly affects dentofacial morphology. Missing teeth in the maxilla result in more retrusive lower incisors and upper lip. Anterior and posterior localization of missing teeth has an important impact on face height measurements.

### SP 288

Friday, June 18 - at 13:00

#### DIFFERENCES IN SOFT TISSUE ANALYSIS BETWEEN CONVENTIONAL AND COMPUTERIZED TRACING METHODS

Ö Polat-Özsoy<sup>1</sup>, S Erken<sup>1</sup>, S Özsoy<sup>2</sup>, <sup>1</sup>Department of Orthodontics, Faculty of Dentistry, Baskent University and <sup>2</sup>Ministry of Health, Ankara Topraklik Dental Hospital, Turkey

**AIM:** To evaluate the accuracy of cephalometric soft tissue measurements using computerized tracing of direct digital radiographs in comparison with hand tracing of digital radiographic printouts.

**MATERIALS AND METHOD:** Thirty pre-treatment cephalometric digital radiographs were traced using the Vistadent OC 1.1 computer software program (group 1) and their conventional radiographic printouts were traced manually (group 2) by the same investigator. The soft tissue analyses of Epker and Holdaway were carried out using both methods. Measurement reproducibility was evaluated by calculating intraclass correlation coefficients (ICC), and paired *t*-tests were used to compare differences in individual measurements between the methods. Differences greater than 0.05 were considered to be statistically significant.

**RESULTS:** ICC were higher than 0.85 for both methods. Soft tissue cephalometric measurements using the analyses of Epker and Holdaway were reproducible for both tracing methods ( $P > 0.05$ ).

**CONCLUSION:** Despite minimal discrepancies in measured values between conventional hand-tracing and the computerized method, any differences were minimal and clinically acceptable.

### SP 289

Friday, June 18 - at 13:00

#### EFFECT OF A CHLORHEXIDINE-FLUORIDE GEL ON PERIODONTAL HEALTH IN ORTHODONTIC PATIENTS WITH FIXED APPLIANCES

D İscan<sup>1</sup>, B Ozdemir<sup>2</sup>, B Balos Tuncer<sup>1</sup>, S Bozkurt<sup>2</sup>, H N İscan<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Periodontology, Faculty of Dentistry, Gazi University, Ankara, Turkey

**AIM:** Cervitec Gel (Ivoclar Vivadent AG, Liechtenstein) is an oral health care gel that includes chlorhexidine and fluoride, and is used for the protection of gums and teeth. The aim of this study was to evaluate the effect of Cervitec Gel on periodontal health and levels of *P. gingivalis* and *T. denticola* in patients undergoing fixed orthodontic treatment for at least 4 months.

**SUBJECTS AND METHOD:** Twenty-five patients with fixed orthodontic appliances randomly divided into two groups. Thirteen patients were assigned to the test group (mean age 15.31 years) and given Cervitec Gel, and the remaining 12 patients comprised the control group (mean age 15.17 years) and were given a placebo. All patients used the same toothbrush (Banat, Turkey). After periodontal prophylaxis and oral hygiene motivation, plaque index (PI), gingival index (GI), probing depth (PD) and bleeding on probing (BOP) were recorded and subgingival plaque samples were collected. All measurements were repeated after toothpaste application on day 0, 2 weeks and 1 month. Bacterial identification was performed using RT polymerase chain reaction. Intragroup and between group comparisons were undertaken using the Student's *t*- and Mann Whitney U-tests.

**RESULTS:** Intragroup comparisons revealed that PI and GI significantly decreased, while PD, BOP and bacterial indices did not show significant changes in either group. No significant differences were determined in intergroup comparisons.

**CONCLUSION:** Application of Cervitec Gel did not demonstrate any effect on the clinical indices and bacterial composition of orthodontic patients with fixed appliances.

### SP 290

Friday, June 18 - at 13:00

#### THE WORLD-WIDE-WEB, ORTHODONTICS AND TEMPOROMANDIBULAR DISORDERS: A BALANCED INFLUENCE ON PATIENT DECISION-MAKING?

F Luther, Dental Institute, University of Leeds, England

**AIM:** Controversy over relationships between malocclusion, temporomandibular disorders (TMD) and whether orthodontic treatment cures TMDs remains despite increasing evidence suggesting the aetiology of TMDs has closer links to other chronic pain conditions than to malocclusion. Nevertheless, some clinicians still use orthodontics to treat TMD patients despite instances of litigation. However, 60-80 per cent of world-wide-web (WWW) users have used it to obtain health information (Powell *et al.*, 2003). So what information might TMD patients access in deciding whether to undergo orthodontic treatment? The aim of this research was to assess information on the WWW that could affect patients' decision-making.

**MATERIALS AND METHOD:** Individuals spend approximately 5 minutes searching the WWW when looking for medical information (Eysenbach and Köhler, 2002). Therefore, using a well known search engine and the terms 'TMD' and 'TMJ', the first 10 relevant websites listed (equating to what might be searched in 5 minutes), were examined for advice suggesting malocclusion causes TMD and/or that orthodontics could be used as a TMD treatment.

**RESULTS:** The term 'TMD' found six (of 10) websites which suggested malocclusion causes TMD and/or that orthodontics could be used as a TMD treatment. The term 'TMJ' found five (of 10) websites which suggested the same. The stance of three (of 20) was unclear. The information provided was mostly based on the lowest level of evidence. Some sites provided contact details for patients to access the specific clinic the website was associated with.

**CONCLUSION:** Over half of websites patients are likely to see, suggest malocclusion causes TMD and/or that TMD can be treated with orthodontics but the information may be unbalanced. This suggests that in order to inform patient decision-making appropriately, further education is required amongst clinicians to ensure their understanding of TMD is based on current evidence. Eysenbach G, Köhler C 2002 Does the internet harm health? Database of adverse events related to the internet has been set up. British Medical Journal 324: 573-577 Powell J A, Darvell M, Gray J A M 2003 The doctor, the patient and the world-wide web: how the internet is changing healthcare. Journal of the Royal Society of Medicine 96: 74-76

### SP 291

Friday, June 18 - at 13:00

#### DENTO-SKELETAL EFFECTS OF RAPID MAXILLARY EXPANSION WITH THE FERRO PROTOCOL

M Sorrentino, G Sgritta, P Spinella, F De Gregorio, L Perillo, Department of Orthodontics, Second University of Naples, Italy

**AIM:** Rapid maxillary expansion (RME) is a procedure indicated in growing subjects to correct dentoskeletal transverse discrepancy. Usually skeletal and dental changes are reported. More skeletal changes can be achieved when the expansion is performed after suture opening as recommended by the Ferro Expansion Protocol (FEP). The aim of this study was to compare the transverse skeletal and dental changes in a group of patients with posterior crossbite before and after FEP, with those derived from a normal growth group using postero-anterior (PA) cephalometric radiographs.

**MATERIALS AND METHOD:** Data from PA cephalograms of 21 patients (17 girls, 4 boys) with a posterior crossbite were compared with those from an age-matched normal growth group. The patients were treated with Hyrax-type RME activated until correction of the crossbite and then used as retainer. The mean age at T0 was 8 years 10 months and at T1, 10 years 8 months. Sixteen reference points were used and eight variables were studied consisting of six skeletal and two dental measurements. A Student's *t*-test was used to compare the differences between groups, and the effects of the FEP on skeletal and dental structures.

**RESULTS:** The FEP group showed statistically significant changes in latero- and medial-orbital, latero-nasal, maxillary, and upper and lower molar widths. When compared with other studies, skeletal changes were greater than dental changes.

**CONCLUSION:** There was some evidence that expansion after opening suture produced by the FEP, may result in more significant transverse skeletal increments than dental movements.

## SP 292

Friday, June 18 - at 13:00

### TIME-DEPENDENT BIOMECHANICAL BEHAVIOUR OF THE PERIODONTAL LIGAMENT – AN IN VITRO STUDY

**K Papadopoulou**<sup>1</sup>, T Eliades<sup>2</sup>, L Keilig<sup>1</sup>, A Jäger<sup>3</sup>, C Bouraue<sup>1</sup>,  
<sup>1</sup>Oral Technology and <sup>2</sup>Department of Orthodontics, University of Bonn, Germany and <sup>3</sup>Department of Orthodontics, Aristotle University, Thessaloniki, Greece

**AIM:** Initial tooth displacement, induced by orthodontic forces, can occur rapidly or slowly depending on the physical characteristics of the applied force and response of the periodontal ligament (PDL). Classic theories of tooth displacement do not consider the complex mechanical properties of the PDL, the morphology of the alveolar structure, or the magnitude of the applied force. Current knowledge indicates that there is a need to investigate how changes in stress and strain distribution of the PDL trigger orthodontic tooth movement. The aim of this study was to evaluate the biomechanical time-dependent behaviour of the PDL in relation to force application.

**MATERIALS AND METHOD:** Eighteen pig jaw segments were used. A linear increasing displacement of 0.1 and 0.2 mm in a time span of 5, 10, 20, 30, 60, 120, 300, 450, 600 seconds was applied to the premolar crowns. Resulting forces were measured using a three-dimensional force torque sensor during the loading phase, as well as for a period of 600 seconds after reaching full deflection. Time/force diagrams for each tooth were generated from the obtained data and maximum force as well as forces after relaxation were obtained. Means and standard deviations for each deflection/time combination were calculated.

**RESULTS:** Forces measured for the different specimens varied significantly and decreased with increasing loading time (mean 1.5 and 0.5 N for 5 and 600 seconds, respectively for 0.1 mm displacement), and increased with increasing displacement (4.8 and 2.6 N for 5 and 600 seconds, respectively at 0.2 mm displacement). Forces after relaxation converged to boundary values of 0.4 and 2.0 N for 0.1 and 0.2 mm displacements, respectively. Correlation of forces with root volumes or surfaces was not significant (Spearman rank test).

**CONCLUSION:** Variations in initial forces may be attributed to the biological tissue characteristics of the specimens and probably to the different periodontal status of the teeth rather than root geometry.

## SP 293

Friday, June 18 - at 13:00

### ORTHODONTIC FORCED ERUPTION IN THE COMPROMISED TOOTH: WEEKLY VERSUS SINGLE FIBROTOMY

**I Frezzato**, A Winkler, L Favero, Department of Orthodontics, University of Padua, Italy

**AIM:** When considering treatment of a single tooth affected by a subgingival lesion, orthodontic forced eruption could be the best treatment choice, if the dental root is long enough, the periodontal conditions are good and the periodontal margin is in an appropriate aesthetic position. In forced eruption the tooth is moved out of its alveolus. Fibrotomy is useful to keep the periodontal margin in the existing position. Many authors recommend a weekly fibrotomy. The aim of this presentation is to propose a single fibrotomy performed at the end of the orthodontic forced eruption.

**MATERIALS AND METHOD:** Nine upper second premolars with subgingival lesions. For each tooth forced eruption was per-

formed using brackets and a stiff stainless steel sectional archwire (0.018 X 0.025 inches), with an inset bend to apply direct axial elastic traction to the tooth. Digital photographs, radiographs and periodontal parameters were taken at three stages. Fibrotomy was performed weekly in four patients, and in five others immediately after active eruption (clinical and radiographic evidence of sufficient crown eruption). Data were clinically compared and statistically analysed with Kendall test.

**RESULTS:** Clinical crown lengthening was obtained in all subjects. The periodontal parameters (probing, gingival margin and alveolar bone crest location) demonstrated similar amount of eruption obtained in a similar treatment time in both groups (3 mm and 7 weeks, respectively). Statistical analysis demonstrated different behaviour in the two groups for all parameters during treatment, while post-treatment the results were similar.

**CONCLUSION:** The clinical advantage of the single final fibrotomy is evident, with less discomfort for the patient than the weekly protocol.

## SP 294

Friday, June 18 - at 13:00

### FORCES IN COMPLEX MALALIGNMENT CORRECTION WITH CONVENTIONAL AND SELF-LIGATING BRACKETS

**M Montasser**<sup>1</sup>, T El-Bialy<sup>2</sup>, A Jäger<sup>3</sup>, C Bouraue<sup>3</sup>, Departments of Orthodontics, <sup>1</sup>University of Mansoura, Egypt, <sup>2</sup>University of Alberta, Edmonton, Canada and <sup>3</sup>University of Bonn, Germany

**AIM:** There is general consensus that advocates the use of the lightest force capable for achieving orthodontic tooth movement. As it is difficult to directly measure stresses and strains within the periodontal ligament of loaded teeth, measuring the forces that are applied directly to teeth is a valuable estimation tool. The aim of this research was to measure the forces during complex orthodontic tooth malalignment correction with various archwire-bracket combinations using the Orthodontic Measurement and Simulation System.

**MATERIALS AND METHOD:** Three types of orthodontic brackets (1) conventional ligating brackets (Victory Series and Mini-Taurus), (2) self-ligating brackets (SmartClip, passive self-ligating bracket and Time3, active self-ligating bracket), and (3) a conventional low friction bracket (Synergy). All brackets had a nominal 0.022 inch slot size. The brackets were combined with four archwire types (1) 0.012 inch stainless steel, (2) 0.012 inch Orthonol [nickel titanium (NiTi)], (3) 0.012 inch Thermalloy (NiTi), and (4) 0.0155 inch coaxial. The archwires were tied to the conventional brackets with stainless steel ligatures and elastomeric rings. The malocclusion simulated displacement of a central upper incisor, 2 mm gingivally (X axis) and 2 mm labially (Z axis).

**RESULTS:** All brackets showed the lowest forces when combined with either the coaxial or Thermalloy wires. The forces ranged from  $3.4 \pm 0.2$  N to  $0.7 \pm 0.1$  N in the X axis direction and  $4.5 \pm 0.3$  N to  $0.5 \pm 0.1$  N in the Z axis direction. The highest force was determined when combined with stainless steel wires, ranging from  $6.3 \pm 0.3$  N to  $3.0 \pm 0.1$  N in the X and  $6.3 \pm 0.3$  N to  $1.7 \pm 0.1$  N in the Z direction.

**CONCLUSION:** Small cross-sectional coaxial wires of 0.0155 inches and Thermalloy wires of 0.012 inches are recommended for use in the initial levelling and alignment stage of orthodontic treatment. Elastomeric rings, when used with conventional brackets, increase the force applied to the teeth.

## SP 295

Friday, June 18 - at 13:00

### ANALYSIS OF SUPPORTING AREAS AND ANGLES ON DENTAL PANTOMOGRAMS USING A NEW PROGRAM

**R Reinhardt**<sup>1</sup>, V Kurojedova<sup>2</sup>, H Wehrbein<sup>1</sup>, Departments of Orthodontics, <sup>1</sup>Dental Clinic of Johannes Gutenberg University Mainz, Germany and <sup>2</sup>Dental Clinic of the Medical Academy Poltava, Ukraine

**AIM:** To test a new computer program to determine whether measurements of supporting areas on dental pantomograms (DPT) or on a model, and angles on DPT and on lateral cephalogram (LCG) give identical results.

**SUBJECTS AND METHOD:** Fifty patients (20 males, 30 females). To determine the size of the supporting areas on DPT the new computer program, fr-Win 7.0, was used. Gonion- and basic angles were measured on the DPT and LCG, and the data were compared with each other.

**RESULTS:** The size range of supporting areas measured on DPT and on model was in exact agreement. With a mean of 0.829, Pearson's coefficient of correlation confirmed a significant relationship between the two methods. The size of gonion angle determined on LCG was in exact agreement with that measured on the DPT. Pearson's correlation coefficient, with a mean of 0.8285, showed a significant relationship between the two methods. The 90 per cent confidence interval (CI) of differences in measurements on LCG of the right and left gonion angles on DPT in equivalence region is evidence for equally exact angle measurements on DPT and LCG. The size of basic angle determined on LCG was in exact agreement with that measured on DPT. Pearson's correlation coefficient had a mean of 0.877. The 90 per cent CI of difference of the measurements on LCG to the right and left basic angles on DPT lies in the equivalence region, the angle measurements on the DPT and the LCG agree with each other.

**CONCLUSION:** With the new computer program exact measurement of supporting areas can be made, which are identical with those taken on a model. The DPT can be used for measurements of gonion and basic angles.

## SP 296

Friday, June 18 - at 13:00

### EFFECTIVENESS OF THE TRANSPALATAL ARCH – A COMBINED CLINICAL AND EXPERIMENTAL STUDY

M Tsetsilas<sup>1</sup>, L Keilig<sup>1</sup>, A Jäger<sup>1</sup>, C Bouraue<sup>1,2</sup>, <sup>1</sup>Department of Orthodontics and <sup>2</sup>Oral Technology, University of Bonn, Germany

**AIMS:** Transpalatal arches (tpa) have long been used for orthodontic treatment. They can minimize undesirable side-effects during treatment, add to anchorage, and allow for active tooth movement with reciprocal or asymmetric mechanics. The purpose of this study was to analyse the forces and moments delivered by the TPA of the 'Precision Lingual Arch System' advocated by Burstone (0.032 X 0.032 inch TMA) and to clinically verify the predicted tooth movement.

**MATERIALS AND METHOD:** Two types of activation were investigated bilateral expansion (six cases) and unilateral expansion with torque application (four cases). Forces and moments produced at the attachments by the two different activations were determined using the Orthodontic Measurement and Simulation System (OMSS). The OMSS is based on the principle of the two-tooth model and allows forces and moments acting on two teeth to be measured simultaneously and the tooth movement resulting from this force system to be simulated. Silicon impressions of 10 patients were taken before and 12 weeks after insertion of the TPA. The casts were scanned with a laser scanner and the exact tooth movement was calculated and compared with the simulated tooth movement delivered from the OMSS. For the two types of activation, the TMA TPAs were adjusted to deliver forces of about 4 N.

**RESULTS:** Analysis of the changes pre- to post-treatment showed expansion up to 4 mm during a time period of 12 weeks for the bilateral cases. In all unilateral cases, the achieved tooth movement was in good correlation with the tooth movement predicted in the OMSS simulations. In all cases, vertical and/or sagittal side-effects were negligible.

**CONCLUSION:** The intended tooth movement, as well as the results of the simulation, were in good correlation with the clinically achieved tooth movement. A treatment time of 12 weeks was not sufficient for complete correction of the transverse discrepancy.

## SP 297

Friday, June 18 - at 13:00

### BONE GRAFTING FOR ORTHODONTIC PURPOSES

C Reichert<sup>1</sup>, W Götz<sup>1</sup>, S Reimann<sup>2</sup>, C Bouraue<sup>1,2</sup>, A Jäger<sup>1</sup>, Departments of 1Orthodontics and 2Oral Technology, University of Bonn, Germany

**AIM:** Bone is a dynamic tissue that is resorbed and remodelled by osteoblast-osteoclast interaction. In special cases, e.g. bone grafting of periodontal defects, cysts or alveolar clefts, tooth movement into such a grafted region is desired. Here, bone substitutes (BS) that are resorbed by osteoclastic cells (OC) and subsequently replaced by newly formed bone would be desirable to avoid side effects e.g. root resorption. Historically the resorption capacity of BS was tested in histological animal studies. There are only few approaches to test biodegradation *in vitro*. The aim of this study was to develop a new method to quantify biodegradation of BS by OC.

**MATERIALS AND METHOD:** OC were cultivated in the presence of RANKL and M-CSF on NanoBone<sup>®</sup>, a BS which has been described for complete biodegradation by OC in histological sections. The calcium content of the media was determined by inductively coupled plasma mass spectrometry followed by a visualization of the cells on the BS was analysed by scanning electron microscopy. Hydrochloric acid and basal cell media served as the control.

**RESULTS:** Compared with the control, the calcium concentration decreased as an initial effect after contacting the BS. A strong liberation of calcium from the BS mediated by OC was observed over the next 14 days.

**CONCLUSION:** By developing this method it is possible to give answers to the questions: Is it possible to cultivate OC on respective BS? Are they inert for hydrolytic processes and may they be biodegraded by OC?

## SP 298

Friday, June 18 - at 13:45

### DEFORMATION AND DEACTIVATION CHARACTERISTICS OF NICKEL-TITANIUM WIRES

B Pseiner, J Bantleon, J Freudenthaler, H-P Bantleon, Department of Orthodontics, Medical University of Vienna, Austria

**AIM:** To quantify, *in vitro*, permanent deformation and the force-deflection-rate of preformed nickel-titanium (NiTi) wires during the preliminary alignment phase.

**MATERIAL AND METHOD:** Five specimens of each NiTi archwire [SmartClip<sup>™</sup> Nitinol Classic 0.012 and 0.014 inch and Nitinol heat-activated 0.014 inch (3M Unitek), Damon<sup>™</sup> Align SE200 0.012 and 0.014 inch and CuNiTi 0.014 inch (Ormco), Biostarter<sup>®</sup> 0.012 and 0.014 inch (Forestadent), Sentalloy<sup>®</sup> 0.014 inch (GAC)] were inserted into a static testing device simulating malpositioned teeth. The deflection ranged from 6 to 1 mm. Permanent deformation was studied by scanning the wires after thermocycling of one day, 14 days and 4 weeks. Force measurements of all 0.014 inch wires (three samples each) were performed at 37°C by simulating a three-bracket relationship deflecting the wire 5, 4, 3, 2, and 1 mm. Similar to the clinical situation, the wire was first inserted into the lateral brackets and then activated into the central malpositioned one. During its gradual levelling, the electronic measuring device recorded the force level at every 0.1 mm step. Data were analyzed by one-way ANOVA and Fisher's protected least significant difference interval, calculated at  $\alpha = 0.05$ .

**RESULTS:** Throughout the entire study period, no clinically relevant, permanent deformation was detected for any of the NiTi archwires. However, three of the five Damon CuNiTi specimens were fractured after the 4 week period. Due to the described activation of the wire, the initial vertical force levels seemed very high on the central force sensor (390-520 cN). After 0.3 mm of deactivation (cf. biomechanical properties of the periodontal ligament), all wires reached their martensitic plateau (except Nitinol classic). These force levels ranged from 110 cN (Sentalloy) to 160 cN (Damon Align) at an initial deflection of 3 mm. Regarding the

force levels of the martensitic plateau, Nitinol classic and Sentalloy showed statistically significant differences to the other tested NiTi wires.

### SP 299

Friday, June 18 - at 13:45

#### COMPARISON OF BOND STRENGTH OF EIGHT CONTEMPORARY ORTHODONTIC ADHESIVE SYSTEMS

**C Reicheneder**, P Proff, A Faltermeier, P Römer, Department of Orthodontics, University of Regensburg, Germany

**AIM:** Bond failure of brackets can significantly increase chair-side time, treatment time and efficiency. Therefore it is necessary to find an ideal combination of adhesive and bracket system. The purpose of this study was to compare the shear and tensile bond strengths of eight common adhesive systems for bonding in orthodontics.

**MATERIALS AND METHOD:** One hundred and sixty freshly extracted bovine mandibular permanent incisors randomly divided into eight groups. Self-ligating aesthetic Damon 3 brackets (Ormco, Orange, California, USA) were bonded using the following adhesive systems: Quick Bond (chemically and light-cured; Forestadent, Pforzheim, Germany), Blugloo (Ormco), Enlight LV (Ormco), Kurasper F (Kuraray Dental, Frankfurt, Germany), Transbond LR (3M Unitek, Monrovia, California, USA), Light Bond (Reliance Orthodontic Products, Itasca, Illinois, USA), and Fuji Ortho LC (GC America, Alsip, Illinois, USA). After 24 hours, half of each group was debonded to measure shear bond strength (SBS) and half to measure tensile bond strength.

**RESULTS:** Blugloo showed the highest SBS values, whereas Transbond LR and Quick Bond (chemically and light-cured) had the lowest. Tensile strength was the highest with Fuji Ortho LC and lowest with Quick Bond (chemically and light-cured) and Kurasper F. The tensile strength of light-cured Quick Bond was approximately 47 per cent lower than that of Fuji Ortho LC.

**CONCLUSION:** Blugloo, Fuji Ortho LC, Light Bond, and Enlight LV are among the materials of choice for bonding fixed orthodontic appliances to teeth. All bond strength values were clinically satisfactory except for the tensile strength of chemically and light-cured Quick Bond and Kurasper F.

### SP 300

Friday, June 18 - at 13:45

#### ORAL HYGIENE INDEX IN MALE AND FEMALE PATIENTS BEFORE ORTHODONTIC TREATMENT

**M Albreht Zore**<sup>1</sup>, F Farčnik<sup>2</sup>, M Ovsenik<sup>3</sup>, <sup>1</sup>Health Care Centre Logatec, <sup>2</sup>Orthos, Ljubljana and <sup>3</sup>Department of Orthodontics, University of Ljubljana, Slovenia

**AIM:** Fixed and removable appliances present a risk of plaque accumulation, white spot lesions and gingival irritation. The patients must be able to maintain adequate oral hygiene during the period of orthodontic treatment and retention. The aim of this study was to establish the level of oral hygiene of patients before orthodontic treatment.

**SUBJECTS AND METHOD:** Sixty-two patients (34 males, 28 females), aged 8-37 years, were screened for oral hygiene at three successive controls before orthodontic treatment. Erythrozyne dye was applied to all tooth surfaces and the oral hygiene index (OHI) was then calculated.

**RESULTS:** Total average OHI was 26 per cent (males 28%, females 24%). Average OHI at the first control in males was 43 per cent and in females in 41 per cent. At the second control it was 28 per cent in males and 19 per cent in females. Average OHI at the third control for males and females was 12 and 10 per cent, respectively.

**CONCLUSION:** OHI evaluation is an important part of diagnosis before treatment in orthodontic patients. Although differences in the OHI assessment between genders occasionally reached statistical significance, interpretation of the results should be treated with caution due to the large age differences of the patients.

### SP 301

Friday, June 18 - at 13:45

#### ROOT RESORPTION AFTER LIGHT AND HEAVY ROTATIONAL FORCES FOR 4 WEEKS – A MICRO-COMPUTER TOMOGRAPHIC STUDY

A T Wu<sup>1</sup>, **T Tamer**<sup>2</sup>, C Colak<sup>2</sup>, S Elekdag-Turk<sup>2</sup>, M A Darendeliler<sup>1</sup>, <sup>1</sup>Discipline of Orthodontics, Faculty of Dentistry, University of Sydney, Australia and <sup>2</sup>Department of Orthodontics, Faculty of Dentistry, University of Ondokuz Mayıs, Samsun, Turkey

**AIM:** To quantitatively measure and compare the location(s), dimensions and volume of root resorption craters in human premolars following the application of controlled light and heavy rotational orthodontic forces over a 28 day (4 week) period.

**SUBJECTS AND METHOD:** Fifteen patients requiring bilateral extraction of the maxillary first premolars as part of their orthodontic treatment. Each patient received a heavy rotational force (225 g) on one premolar and a light rotational force (25 g) on the contralateral premolar. Orthodontic rotational forces were applied over 28 days with buccal and palatal cantilever springs; 0.016 inch beta-titanium molybdenum alloys were used to apply the light force and 0.018 inch stainless steel was used for the heavy force. Following the 28 day experimental period, the upper first premolars were extracted under stringent protocols to prevent root surface damage. The samples were then scanned using a microcomputer tomographic X-ray system and analysed using specific software to obtain direct volumetric measurements.

**RESULTS:** The mean volume of resorption craters in the light force group was 0.42 compared with the heavy force group of 0.51 ( $P = 0.013$ ). When separated at the root level, the difference in volume of root resorption craters between the two groups was significantly different only at the mid level ( $P = 0.001$ ). Root resorption craters were consistently detected at the boundaries between the buccal and distal surfaces, and the mesial and lingual surfaces. Positive areas developed significantly more root resorption craters at all three levels, as compared with minimal areas ( $P < 0.001$ ).

**CONCLUSION:** Heavy rotational forces cause more root resorption than light rotational forces. Compression areas (buccal-distal and lingual-mesial surfaces) showed significantly greater root resorption than other areas at all levels of the root.

### SP 302

Friday, June 18 - at 13:45

#### CONE BEAM COMPUTED TOMOGRAPHIC BASED TREATMENT OUTCOME ANALYSES OF ACTIVE AND PASSIVE SELF-LIGATING BRACKETS

**P Cattaneo**, M Treccani, O Annousaki, A Myrda, B Melsen, Department of Orthodontics, School of Dentistry, University of Aarhus, Denmark

**AIMS:** Self-ligating brackets, combined with the prescribed archwires, are reported to lead to a widening of the alveolar process together with good torque control. The aims of this randomized prospective clinical trial were to assess changes in the torque of central incisors, transverse tooth movements and buccal bone modelling in the maxilla achieved with active or passive self-ligating systems.

**SUBJECTS AND METHOD:** Patients with Class I, Class II and mild Class III malocclusions, all characterized by crowding, were randomly allocated for treatment with passive (Damon® 3 MX) or active (In-Ovation) self-ligating brackets. Impressions and cone beam computed tomographic (CBCT) scans were taken before (T0) and after (T1) the completion of treatment, and digital study casts were generated. The overall movements of the central incisors, canines, first and second premolars, and first molars were evaluated. Changes in basal alveolar bone and dental arch shape were assessed using three-dimensional colour-coded maps.

**RESULTS:** Twenty-two patients in the Damon group and 20 in the In-Ovation group completed treatment. For both groups: 1) expansion of the dental arch took place in the canine and even

more in the premolars regions; 2) transverse expansion mainly occurred as a result of tipping; 3) neither bodily movement of the teeth nor augmentation of basal bone could be detected. There was no statistically significant difference in torque-control of the two self-ligating systems, though a tendency for more torque-control was observed with the active brackets.

**CONCLUSION:** In the majority of the cases, the anticipated true-expansion and buccal bone modelling when using self-ligating brackets could not be confirmed using CBCT-based analyses. Active and passive self-ligating brackets showed no difference in torque control. Due to the large inter-individual variation, a patient-specific analysis seems to be mandatory to identify individual factors influencing the final treatment outcome.

### SP 303

Friday, June 18 - at 13:45

#### CEPHALOMETRIC PARAMETERS AND SPEECH ABILITY IN DOWN SYNDROME

**E Pozzani**, G Piana, University of Bologna, Italy

**AIM:** To determine statistically significant correlations among craniofacial characteristics of subjects affected by Down syndrome (DS), which is the most common chromosomal defect, and their oro-facial muscle function and speech ability.

**SUBJECTS AND METHOD:** Thirty-one patients (15 females, 16 males), aged from 15 to 30 years. They were all affected by DS and presented average mental delay (IQ between 35 and 55). Sagittal skull radiographs were obtained and cephalometric analysis was performed by the same person. The following parameters were measured: SNA, SNB, ANB, IMPA, N-S, S-Ar, Go-Me, Ar-Go, N-S-Ar, S-Ar-Go, Ar-Go-Me, Björk  $\Sigma$ , N-Me, S-Go, SN-GoGn, SN-Sna/Snp, Sna/Snp-Go/gn, Ls-L-PgC, Li-L-PgC. In addition, in the study group oro-facial muscle function and speech ability tests were undertaken. Nemoceph software was used for computer cephalometric evaluation. The subjects were divided into two subgroups based on the standard deviation (SD) of the cephalometric landmarks. Those in the range of  $\pm 1$  SD had a value of 1 and outside this range a value of 0. One group had cephalometric values between 5 and 11 and the other group between 12 and 18. A Student's t-test was used for statistical evaluation of quantitative variables.

**RESULTS:** A statistically significant correlation was found between the improved cephalometric parameters of the first group and the numeric values obtained for oro-facial muscle function and speech ability tests ( $P = 0.05$ ). There was also a statistically significant correlation between the poorly defined cephalometric parameters of the second group and the numeric values obtained for oro-facial muscle function and speech ability tests ( $P = 0.05$ ).

**CONCLUSION:** The development of the skull and face of DS patients is highly related to oro-facial muscle function and speech ability. Early physiotherapy, speech therapy, dentofacial orthopaedics and orthodontics have a direct impact on the quality of life of DS patients.

### SP 304

Friday, June 18 - at 13:45

#### mRNA EXPRESSION OF REGULATORY AND GROWTH FACTORS IN OROFACIAL MUSCLES OF DYSTROPHIC MICE

**A Spassov**<sup>1</sup>, T Gredes<sup>1</sup>, T Gedrange<sup>1</sup>, D Pavlovic<sup>2</sup>, C Kunert-Keil<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Anaesthesiology and Intensive Care Medicine, University of Greifswald, Germany

**AIM:** Activity of muscle regulatory factors (MRFs) and mechano growth factors (MGFs) increase in regenerating muscle. Little is known about their precise role and particularly their mRNA expression in the masticatory muscles of mdx mice [an animal model of Duchenne muscular dystrophy (DMD)]. Elucidating some aspects of the process of regeneration in the mdx mice may be relevant for a better understanding of DMD. The aim of this investigation was to study gene expression of MRFs and MGFs in mdx mice masticatory muscles as possible indicators of muscle response to dystrophic changes.

**MATERIALS AND METHOD:** In masticatory muscles of male and female mdx mice the mRNA expression of MRF and MGF was determined by quantitative RT-polymerase chain reaction (mdx versus controls; 100 days old, each group  $n = 10$ ).

**RESULTS:** Quantities of mRNA of insulin-like growth factor-1, MGF and MyoD1 in mdx mice were found to be comparable with the controls. Decreased myostatin mRNA was found in mdx masseter (MAS) and tongue (TON) (3- and 2-fold,  $P < 0.005$  and  $< 0.05$ ). Inversely, the myogenin mRNA was increased in mdx MAS (6-fold) and temporal (TEM) muscles (2.5-fold) (both  $P < 0.05$ ). An unequal expression of MRFs was found between the genders. Myogenin and MyoD1 mRNA was increased in mdx female MAS (2-fold and 11-fold,  $P < 0.01$  and  $P < 0.001$ , respectively). In male mdx mice mRNA myostatin in TEM was decreased (10-fold) as compared with the male control mice.

**CONCLUSION:** Myogenin and myostatin may play an important role in the regeneration process in mdx mouse masticatory muscles that could be regulated by a sex-hormone dependent mechanism.

### SP 305

Friday, June 18 - at 13:45

#### CEPHALOMETRIC MORPHOLOGY OF ADULTS WITH OBSTRUCTIVE SLEEP APNOEA

**A M Enache**<sup>1</sup>, V R Nimigean<sup>2</sup>, A C Didilescu<sup>3</sup>, N Maru<sup>4</sup>, Departments of <sup>1</sup>Orthodontics and Dentofacial Orthopedics, <sup>2</sup>Oral Rehabilitation, <sup>3</sup>Anatomy and Embryology and <sup>4</sup>Topographical and Clinical Anatomy, University of Medicine and Pharmacy 'C.Davila', Bucharest, Romania

**AIM:** Obstructive sleep apnoea (OSA) is a common respiratory disorder, with a frequency that increases with age. The mechanisms contributing to the aetiology of OSA include genetic and environmental changes in craniofacial dimensions. The aim of this study was to assess the cephalometric features of Romanian adults with OSA.

**MATERIALS AND METHOD:** Cephalometric radiograph of 20 randomly selected adult OSA patients (5 females, 15 males, aged 34-76 years) with a mild to moderate apnoea/hypopnoea index (range 5-23) and a mean Body Mass Index of  $28.75 \pm 4.13$ . Linear and angular parameters for dental and skeletal features were assessed.

**RESULTS:** The mean values for SNA and SNB were:  $79.65 \pm 4.89$  and  $76.77 \pm 3.58$ , respectively, with the cranial base angle within the normal range ( $131.2 \pm 4.84$ ). The mean values for the parameters of the vertical skeletal pattern were: NL-NSL  $7.47 \pm 4.07$ , ML-NSL  $29.82 \pm 5.10$ , upper molar-NF  $26.82 \pm 3.81$  and lower molar-MP  $38.62 \pm 3.44$ .

**CONCLUSION:** Romanian adult patients with OSA have a retrognathic anteroposterior skeletal pattern with a severe retroposition of the mandible, but also a retruded maxilla and a tendency to counterclockwise rotation of the mandible.

### SP 306

Friday, June 18 - at 13:45

#### USE OF DELMOPINOL GEL IN PATIENTS UNDERGOING FIXED ORTHODONTIC TREATMENT

**D Dalessandri**, M Dalessandri, A Bordanzi, S Bonadeo, F Zotti, Department of Orthodontics, University of Brescia, Italy

**AIM:** Multibracket orthodontic treatment makes it difficult to maintain correct oral hygiene, thus facilitating plaque accumulation and gingival inflammation. The aim of this study was to verify the effectiveness of Decapinol Gel to reduce plaque formation and gingival inflammation when applied around brackets.

**SUBJECTS AND METHOD:** Thirty subjects randomly separated into two equal groups. Before orthodontic treatment (T0) the plaque (PI) and gingival (GI) indices were measured and professional scaling, instruction and motivation were carried out. After 15 days (T1) the measurement were repeated and the brackets bonded. Group B were trained to apply the gel around each

bracket at home, after brushing and flossing. Measurements were repeated after 15 (T2), 30 (T3) and 45 (T4) days. At T4 treatment was interrupted for 15 days, and then at T5 group A began to use the gel and indices were obtained after 15 (T6), 30 (T7) and 45 (T8) days.

**RESULTS:** At T1 all patients showed both overall PI and GI equal to or lower than 1. At T2 group A showed significantly higher indices than group B. At T3 the difference between the two groups was lower, but still statistically significant ( $P < 0.05$ ). At T4 the indices were still higher in group A than in group B, but the difference was not statistically significant. At T5, no significant difference was observed between the groups. At T6 GI and PI were higher in group B group, but the difference was not statistically significant. At T7 group B showed significantly higher overall indices than group A. At T8 the indices were higher in group A, but the difference was not statistically significant.

**CONCLUSION:** Decapinol Gel can give beneficial effects especially during the early weeks of orthodontic treatment, when patients are not well trained in bracket maintenance.

### SP 307

Friday, June 18 - at 13:45

#### EFFICACY OF AN INDIRECT BONDING TECHNIQUE IN REDUCING PLAQUE RETENTION

**L Laffranchi**, S Bonadeo, A Bordanzi, F Zotti, D Dalessandri, Department of Orthodontics, University of Brescia, Italy

**AIM:** Excessive etching and residual orthodontic adhesive after bonding can facilitate plaque accumulation that causes gingival inflammation. The aim of this research was to establish if an indirect bonding technique with targeted enamel etching and adhesive placement can reduce plaque retention around brackets.

**SUBJECTS AND METHOD:** Thirty patients in whom brackets were bonded using a split mouth design either direct in one quadrant of the upper arch or opposite in the lower quadrant, and indirect in the other two quadrants. With indirect bonding an acrylic custom made tray with holes on the labial tooth surface, exactly corresponding to the bracket base, was prepared for each patient for enamel etching and orthodontic adhesive application. For the first six months of treatment plaque accumulation indices were recorded by one operator who also registered the presence of white spots before and at the end of orthodontic treatment.

**RESULTS:** Plaque accumulation was higher in the first months of treatment for all patients, and then decreased. The mesial and distal brackets edges showed more plaque accumulation than the occlusal edges. A significantly higher presence of plaque deposits during all experimental periods, and of white spots after debonding, was found for those teeth bonded using the direct technique.

**CONCLUSION:** Indirect bonding that involves the use of a custom made tray that allows the clinician to etch dental enamel and apply orthodontic adhesive only where the bracket base is to be bonded, results in a reduction in plaque retention around brackets and a reduction in white spot formation.

### SP 308

Friday, June 18 - at 13:45

#### MUSCULAR EVALUATION IN SKELETAL CLASS II AND III SUBJECTS AT THE END OF GROWTH

**G Galbiati**, L Giannini, S Tronca, F Iazzetti, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** To evaluate neuromuscular functionality and mandibular kinesiology differences between skeletal Class II and III non-growing patients.

**SUBJECTS AND METHOD:** Seventy patients periodically analysed during surgical orthodontic therapy to assess muscular functionality and mandibular kinematic during the different phases of treatment using an electromyographer, Freely (De Gotzen) and the electromyographer and electrognatographer (K6-I, Myotronics). In addition differences between the skeletal Class II and III subjects were analysed to identify different neuromuscular func-

tionality as a consequence of the different alterations of maxillary skeletal relationships. Statistical analysis was undertaken with ANOVA, with significance at the  $P < 0.05$  level.

**RESULTS:** At the beginning of treatment differences existed between the Class II and III patients concerning the IMPACT (expression of muscular activity in real time) and muscular activity during maximal contraction with and without cotton cylinders interposed between the two arches. In the post-operative phases this difference was reduced and at the end of treatment all the patients showed similar muscular activity, indicating the corrective role of surgery. The differences were statistically significant.

**CONCLUSION:** The authenticity of the binomial form-function has been confirmed. Morphology and structure influence the functional capacities of a subject at the end of growth and correction of the dysgnathia.

### SP 309

Friday, June 18 - at 14:30

#### CORRECTION OF ASYMMETRIC TRANSVERSE DISCREPANCIES

**G Galbiati**, L Giannini, S Tronca, F Iazzetti, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** Premature loss of a primary tooth often causes a loss of space and a deviation of the dental midlines. The aim of this presentation is to propose a new application for the rapid palatal expander (RPE) in patients in the mixed dentition that present an asymmetric transverse discrepancy (monolateral crossbite).

**SUBJECTS AND METHOD:** Ten patients, characterized by a maxillary transverse hypoplasia and an interincisive midline misalignment, due to early loss of a primary tooth. A modified Hyrax RPE, made of two bands for the first maxillary permanent molars or for the second primary molars, two palatal arms reaching the mesial planes of the canines and a vestibular arm banded to the central incisor, contralateral to the misalignment was used. The RPE was activated one-quarter turn, twice a day, for 15 days. At the end of the activation the appliance was left *in situ* for 6 months.

**RESULTS:** Maxillary transverse expansion resulting in the creation of a large interincisive diastema was achieved in all patients. The anchorage teeth tipped vestibularly causing a further increase in the transverse arch dimensions. In the sagittal plane rotation of the maxilla in a downward and forward direction occurred due to activation of all circummaxillary sutures with consequent mandibular backward rotation and increase of facial convexity. In the skeletal Class I patients, the change in ANB angle was minimal, while in the skeletal Class II subjects it decreased and in Class III cases it increased, improving in both cases the sagittal relations. Vertically the dimensions increased but this effect relapsed to the original situation during the retention phase.

**CONCLUSION:** A RPE with a vestibular arm is able to restore the correct transverse diameters of the maxilla and at the same time recover interincisive midline symmetry.

### SP 310

Friday, June 18 - at 14:30

#### BOND STRENGTH OF CERAMIC BRACKETS USING DIFFERENT CURING MODALITIES

R Fusco<sup>1</sup>, D Prisco<sup>2</sup>, **V Grassia**<sup>1</sup>, R De Santis<sup>3</sup>, L Perillo<sup>1</sup>, <sup>1</sup>Second University of Naples - Orthodontic Division, <sup>2</sup>University of Naples 'Federico II', <sup>3</sup>IMCB-CNR Institute of Composite and Biomedical Materials, Italy

**AIM:** Ceramic brackets are commonly used in combination with adhesive systems and composite photocured materials in orthodontics. Soft light energy release (SLER<sup>®</sup> Mectron Spa, GE) has recently been introduced in dentistry for photocuring of composite materials. This technology allows thermal control of the curing process by gradually decreasing the light energy in the final step of the irradiation period of the composite material. The aim of this study was to investigate the influence of SLER<sup>®</sup> technology, compared with standard curing modalities, on the bond strength of ceramic brackets applied with a photocured adhesive system.

**MATERIALS AND METHOD:** Forty bovine teeth were used as substrates for bonding ceramic brackets (Clarity APC Plus, 3M). The samples were randomly divided into two groups and photocured with two different modalities (standard and SLER®). After bonding, the samples were stored in water for 24 hours prior to testing (Torsiometer IMAD 5 Nm).

**RESULTS:** Statistical analysis showed a significant difference between the two groups. Samples photocured with SLER® technology demonstrated an increase in mean bond strength values.

**CONCLUSION:** SLER® technology allows thermal control during the final irradiation time, improving the mechanical properties of the composite materials used for bracket bonding.

### SP 311

Friday, June 18 - at 14:30

#### ORTHODONTIC-SURGICAL TREATMENT: ELECTROMYOGRAPHIC AND ELECTROGNATHOGRAPHIC EVALUATION

**L Giannini**, G Galbiati, F Iazzetti, S Tronca, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** Orthodontic surgical treatment has to be planned not only based on clinical and radiographic data, but also by analyzing the neuromuscular function of patients. The aim of this work is the presentation of the evaluation of stomatognathic function through electromyography and electrognathography during orthodontic-surgical therapy.

**SUBJECTS AND METHOD:** Eighty patients who required orthodontic surgical treatment. Two electromyographic analyses and an electrognathograph were used for functional evaluation of the stomatognathic system, a Freely De Gotzen electromyogram and K6-I electromyogram and kinesiograph. Two duo trode silver/silver chloride electrodes were used. The obtained data were evaluated to analyse changes in function and mandibular kinematics over time. The patients were divided into groups based on gender, vertical dimension and skeletal Class in order to evaluate differences between the groups. Statistical analysis was conducted with Levene analysis followed by *t*- and ANOVA tests with Bonferroni *post hoc* correction.

**RESULTS:** At the beginning of the treatment the patients presented compensatory equilibrium to dysgnathia that was altered by pre-surgical orthodontics and by surgical trauma. Functional rehabilitation continued after surgery.

**CONCLUSION:** The diagnostic and prognostic roles of electromyography and kinesiography in surgical orthodontic treatment have been demonstrated. Functional evaluation should be included in diagnostic and therapeutic surgical-orthodontic treatment planning.

### SP 312

Friday, June 18 - at 14:30

#### FUNCTIONAL EVALUATION IN ADULT PATIENTS WITH OPEN AND DEEP SKELETAL BITES

**L Giannini**, G Galbiati, F Iazzetti, S Tronca, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** To compare mandibular kinesiography and neuromuscular function between open and deep skeletal bite adult patients before and after surgical orthodontic treatment.

**SUBJECTS AND METHOD:** Seventy patients (33 skeletal open bite, 37 skeletal deep bite) undergoing orthodontic-surgical treatment who periodically underwent electromyographic evaluation of the masticatory muscles (anterior temporal and masseter) and electrokinesiographic evaluation of mandibular movements. Two electromyographic instruments were used: the Freely electromyograph (De Gotzen, Legnano, Italy) and the electromyograph and electrognathograph K6-I EMG (Myotronics, Tukwila Washington, USA). Statistical evaluation was performed with *t*-tests and ANOVA.

**RESULTS:** A large number of differences were found between the open and deep skeletal bite patients underlined by analysis of the electromyographic data obtained with both electromyographic systems.

**CONCLUSION:** Morphological differences between open and deep bite patients can be demonstrated by instrumental examination and their correction after surgery is observable electromyographically and electrognathographically.

### SP 313

Friday, June 18 - at 14:30

#### THE TRANSVERSE SAGITTAL MAXILLARY EXPANDER: A CEPHALOMETRIC INVESTIGATION IN GROWING PATIENTS

**S Tronca**, F Iazzetti, L Giannini, G Galbiati, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** To cephalometrically evaluate the skeletal and dental effects of the transverse sagittal maxillary expander (TSME) in children with maxillary hypoplasia.

**SUBJECTS AND METHOD:** Fifty patients of both genders, aged from 6 to 15 years, with a maxillary crossbite caused by basal apical narrowness treated with the TSME. For each patient lateral cephalometric analysis was undertaken before treatment and at the end of the retention period. Changes during the observation period were calculated and statistically analyzed.

**RESULTS:** SNA and ANB angles showed a non-significant increase during treatment. SNB angle increased. SN-GoGn increased in most of the patients but no statistically significant modifications were found. SN-SNP.SNA showed a non-statistically significant rotation. The vertical dimensions (S-Go, N-Me) also showed a non-significant modification during treatment. The linear measurement SNP-A showed a statistically significant increase, indicating that the dentoalveolar maxillary process moved anteriorly because of the force delivered by the sagittal screws during active growth. A significant and statistically significant increase in I<sup>1</sup>SN and I<sup>1</sup>FH angles was observed. Overjet changed from negative to positive in all patients.

**CONCLUSION:** The TSME can produce skeletal changes due to the transverse force, and dentoalveolar modification due to the sagittal component that can increase the available arch length resulting in correction of maxillary hypoplasia. The non-significant increase in the anterior vertical dimension may also suggest the use of the TSME in patients with an anterior open bite.

### SP 314

Friday, June 18 - at 14:30

#### COMPARISON OF ARCHAIC AND CONTEMPORARY ITALIAN SKULLS USING THREE-DIMENSIONAL COMPUTED TOMOGRAPHIC VOLUME RENDERING IMAGES

S Cappabianca<sup>1</sup>, **G Tufano**<sup>2</sup>, F Iaselli<sup>1</sup>, V Esposito<sup>3</sup>, L Perillo<sup>2</sup>, <sup>1</sup>Department of Clinical and Experimental Internistics 'F. Magrassi and A. Lanzara' Scientific Section of Radiology, Radiotherapy and Nuclear Medicine, <sup>2</sup>Department of Orthodontic Second University of Naples and Department of Public, Clinic and Preventive Medicine, Section of Human Anatomy, Anatomy Museum, Italy

**AIM:** Variations in linear and angular craniometric measurements correlate with the evolution of the nervous system and gnathic development. The purpose of this research was to assess the evolution of the neuro- and splanchnocranium over two millennia in a southern Italy population through a comparative craniometric analysis of archaic Pompeian and contemporary skulls.

**MATERIALS AND METHOD:** Twenty skulls dating back two millennia, found during excavations of ancient Pompeii were computer tomographically scanned by sequential and contiguous 1 mm slices. On subsequent volume rendering software reconstructions, craniometric landmarks, for linear and angular measurements, were identified. These features were compared with modern skulls on latero-lateral cephalograms.

**RESULTS:** Significant differences of some linear and angular parameters were found between the ancient and contemporary skulls. Major differences concerned prominence of the frontal bone, skull base angle, length of the anterior and posterior skull base, distances from sella to the frontal bone, from basion to

bregma, from sella to lambda, from nasion to lambda and the diameter of the neurocranium from nasion to opisthocranium, transverse dental arch dimension and arch depth. The latter two measurements are important to define the evolution of gnathic development.

**CONCLUSION:** Comparative assessment of linear and angular craniometric measurements of ancient Pompeian and modern skulls clarified the relationship between the neuro- and splanchnocranium and the gnathic development.

### SP 315

Friday, June 18 - at 14:30

#### DIAGNOSTIC SET-UP AND OCCLUSAL SPLINT GUIDE

**F Iazzetti**, L Giannini, G Galbiati, S Tronca, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** Model surgery, typically performed in the week before the surgical procedure, is an integral part of orthognathic surgery treatment protocol. The purpose of this work is to describe an experimental protocol to optimise pre-surgical orthodontic planning by the study of skeletal discrepancies.

**MATERIALS AND METHOD:** The protocol comprises the following sequence of treatment: 1) face-bow record; 2) three-dimensional cephalometric surgical prediction on lateral, postero-anterior and axial radiographs; 3) model surgery based on cephalometric and aesthetic planning simulated on a semi-adjustable articulator. After model surgery, a pre-surgical diagnostic set-up is created; 4) reproduction of the original skeletal situation discrepancies on the articulator – this position represents the final orthodontic preparation to achieve before surgery; 5) an occlusal thermoplastic material splint tightly formed over the teeth is then fabricated representing a guide for the pre-surgical preparation.

**CONCLUSION:** Using measurements provided by the orthodontist and surgeon, the orthodontic laboratory can create the pre-surgical set-up, from which a soft splint can be fabricated for clinical reference. The orthodontic diagnostic set-up is completed to the ideal occlusion, representing the final tooth position after surgery. This type of protocol has the overall purpose of improving the quality of care to the patient and allows the orthodontist to improve dental and skeletal treatment planning leading to better biomechanics and treatment results.

### SP 316

Friday, June 18 - at 14:30

#### IS CARIES IN THE EARLY MIXED DENTITION A PREDICTOR OF CARIES RISK DURING ORTHODONTIC TREATMENT?

**V Soumas**, C B Staudt, S Kiliaridis, Division of Orthodontics, University of Geneva, Switzerland

**AIM:** During orthodontic treatment with fixed appliances oral hygiene is difficult and mineral loss may occur. Patients with a high caries risk should be identified before the start of orthodontic treatment. The objective of the present study was to evaluate if caries in the pre-treatment early mixed dentition is a predictor for the development of caries in the permanent dentition during orthodontic treatment.

**MATERIALS AND METHOD:** The orthodontic database of the University of Geneva was searched for consecutive patients who started treatment in the early mixed dentition and finished treatment in the permanent dentition during 2003-2008. Forty-one patients with pre- and post-treatment radiographs were included. The overall mean treatment duration, generally two-phase, was 46 months. Bitewings and dental pantomograms were evaluated and the DMFT index was calculated.

**RESULTS:** The initial DMFT was 5.5 (SD = 3.7) and final DMFT 3.0 (SD = 3.2). Pre-treatment, 54 per cent of primary molars were affected by caries, followed by 27.4 per cent of the first permanent molars. Post-treatment, 41.4 per cent of the permanent first molars had carious lesions or fillings. A significant correlation was found between caries experience in the early mixed dentition and that during orthodontic treatment ( $r = 0.47$ ,  $P = 0.002$ ).

**CONCLUSION:** Children with high caries activity before treatment are exposed to a higher risk of developing carious lesions during orthodontic treatment. It is advised that these children should undergo an extended prophylaxis programme before and during orthodontic treatment.

### SP 317

Friday, June 18 - at 14:30

#### EVALUATION OF MICROLEAKAGE WHEN BRACKETS ARE CURED FROM DIFFERENT SIDES

**S I Ramoglu**<sup>1</sup>, T Uysal<sup>1</sup>, C Sisman<sup>1</sup>, H Ertas<sup>2</sup>, Departments of 1Orthodontics and 2Conservative Dentistry, Faculty of Dentistry, Erciyes University, Kayseri, Turkey

**AIM:** Metal brackets do not conduct light as well as ceramic brackets during light curing, which may cause incomplete adhesive polymerization. In previous studies, higher microleakage was reported for metal brackets at the gingival sides. The purpose of this *in vitro* study was to investigate if curing the bracket-adhesive complex through different sides; gingival, occlusal and both mesial and distal, will affect the amount of microleakage at the gingival and the occlusal sides of the brackets.

**MATERIALS AND METHOD:** Forty-five freshly-extracted human maxillary premolar teeth divided into three equal groups: group 1 cured at the occlusal side, group 2 cured at the gingival side and group 3 at both the mesial and distal sides. The dye penetration method was used for microleakage evaluation. Microleakage was determined under a stereomicroscope between the enamel-adhesive and bracket-adhesive interfaces. Statistical analysis was performed using Kruskal-Wallis and Wilcoxon tests.

**RESULTS:** When the gingival and occlusal sides were compared, gingival scores were found to be significantly higher for group 1 ( $P < 0.05$ ) for both interfaces. According to multiple group comparisons, statistically significant differences were found at the gingival side for the enamel-adhesive interface ( $P < 0.05$ ) and bracket-adhesive interface ( $P < 0.01$ ).

**CONCLUSION:** Curing the bracket-adhesive complex through different sides affected the amount of microleakage at the gingival side.

### SP 318

Friday, June 18 - at 14:30

#### META-ANALYSIS OF SKELETAL MANDIBULAR CHANGES DURING FRÄNKEL APPLIANCE TREATMENT

**R Cannavale**, M I Castaldo, C I Giordano, G Padricelli, L Perillo, Department of Orthodontics, Second University of Naples, Italy

**AIM:** To perform a meta-analysis to verify mandibular changes produced by the Fränkel-2 (FR-2) appliance during treatment of growing patients with Class II malocclusions when compared with untreated growing Class II subjects.

**MATERIALS AND METHOD:** The literature published from January 1966 to January 2009 was reviewed with search engines, PubMed, the Cochrane Central Register of Controlled Trials (CENTRAL), Scirus, Lilacs, Embase and Scopus. A quality analysis was performed. The effects on primary endpoints were calculated with random-effect models. Heterogeneity was assessed using Q statistic and investigated using study-level meta regression.

**RESULTS:** A total of nine articles were identified. The quality of the studies ranged from low to medium. Meta-analysis showed that THE FR-2 was associated with enhanced mandibular body length [0.4 mm/year 95% confidence interval (CI) 0.182 to 0.618], total mandibular length (1.069 mm/year, 95% CI 0.683 to 1.455) and mandibular ramus height (0.654 mm/year, 95% CI 0.244 to 1.064). A consistent heterogeneity among studies was found for all the considered linear measurements.

**CONCLUSION:** The FR-2 appliance had a statistically significant effect on mandibular growth. Nevertheless, the heterogeneity of the FR-2 effects, the quality of studies, the differences in age, skeletal age, treatment duration and the inconsistent initial diagnosis seem to overstate the benefits of THE FR-2 appliance. An

evidence-based approach to the orthodontic outcomes of FR-2 appliance is needed to select and compare groups of children with the same cephalometric characteristics with and without treatment.

### SP 319

Friday, June 18 - at 14:30

#### COMPARISON IN THE ADHERENCE LEVEL OF STREPTOCOCCUS MUTANS AND STREPTOCOCCUS SOBRINUS ON SELF-LIGATING BRACKETS\*\*\*

**C-J Hwang**, J-Y Cha, J-W Yoon, S-H Choi, College of Dentistry, Yonsei University, Seoul, Korea

**AIM:** To examine the adherence capacity of self-ligating brackets of cariogenic *S. mutans* and *S. sobrinus* that can result in enamel decalcification.

**MATERIALS AND METHOD:** Five types of self-ligating bracket (Clippy-C, Tomy; Mini Clippy; Clarity SL; Speed; Damon 3) for the experimental group and the composite resin bracket (Spirit MB), metal brackets (Victory) and polycrystalline alumina brackets (Clarity) for the control group. To assess the adherence status, the brackets of the experimental and control groups were placed on a 24-well plate containing basal medium with 20 per cent sucrose added and cultured for 3, 6, and 24 hours.

**RESULTS:** For *S. mutans*, there was no difference in the total adhesion amount by bracket type or incubation time ( $P > 0.05$ ). There was a significant difference in the total adhesion amount with the different types of bracket in *S. sobrinus*. Victory showed greater adhesion than Clarity SL and Spirit MB ( $P < 0.05$ ). The adhesion amount for *S. Sobrinus* after 3 hours of incubation was higher than for Clippy-C and Spirit MB. There was no significant difference of adhesion amount following 6 and 24 hours incubation ( $P < 0.05$ ). Scanning electron microscopy examination showed that adhesion amount assumed a high rate for Clippy-C and Victory, and a low rate for Clarity SL.

### SP 320

Friday, June 18 - at 14:30

#### EFFECTS OF AMORPHOUS CALCIUM PHOSPHATE CONTAINING COMPOSITE ON ENAMEL DEMINERALIZATION

T Uysal<sup>1</sup>, **M Amasyali**<sup>2</sup>, S Özcan<sup>3</sup>, A E Koyuturk<sup>4</sup>, D Sağdıç<sup>2</sup>, Departments of Orthodontics, <sup>1</sup>Erciyes University, Kayseri, <sup>2</sup>Gulhane Military Medical Academy, Ankara, <sup>3</sup>Department of Conservative Dentistry, Gazi University, Ankara, <sup>4</sup>Department of Pediatric Dentistry, Ondokuz Mayıs University, Samsun, Turkey

**AIMS:** To evaluate the *in vivo* effects of an amorphous calcium phosphate (ACP) containing orthodontic composite in reducing enamel demineralization around orthodontic brackets, and to compare it with a control.

**SUBJECTS AND METHOD:** Fourteen orthodontic patients divided into two equal groups; they received brackets on all first premolars, bonded with either Aegis Ortho® (Bosworth Co.), an ACP containing orthodontic composite (experimental group), or Concise® (3M Dental Products), a resin-based orthodontic composite (control group). After 30 days, the teeth were extracted, longitudinally sectioned, and evaluated by superficial microhardness analysis. The determinations were made at the bracket edge cementing limits and at occlusal and cervical points 100 and 200 µm from the edge. In all of these positions, indentations were made at depths of 10, 20, 30, 50, 70, and 90 µm from the enamel surface. Analysis of variance (ANOVA) and Tukey *post hoc* test was used. The statistical significance level was set at  $P < 0.05$ .

**RESULTS:** ANOVA showed statistically significant differences for position, material, depth, and their interactions ( $P = 0.000$ ). The multiple comparison test showed that the ACP containing orthodontic composite was significantly more efficient than the control composite, reducing enamel demineralization in almost all evaluations ( $P = 0.000$ ).

**CONCLUSION:** ACP containing orthodontic composite for bonding orthodontic brackets successfully inhibited demineralization *in vivo*. This effect was localized to the area around the brackets and was statistically significant after 30 days.

### SP 321

Friday, June 18 - at 10:00

#### THE IMPORTANCE OF HEATH EDUCATION OF PATIENTS WITH FIXED ORTHODONTIC APPLIANCES

**S Matijević**, Health Center Tivat, Montenegro

The aim of this study is to draw attention to proper and regular oral hygiene in those patients because of possible occurrence of gingivitis and more frequent occurrence of secondary caries and onset of carious lesion.

**METHODS:** The research was conducted on 50 subjects, 14 to 16 years of age, of both sex with fixed orthodontic appliances. The patients were observed from the beginning to the end of the treatment. At the beginning they were given information about proper hygiene and the use of adequate aids (fluorogal sol, special interdental toothbrush) given the importance of even better oral hygiene for those patients in order to prevent the complications.

**RESULTS AND CONCLUSION:** After several monthly check-ups it has been shown that the motivation has weakened in more than 30% of the patients and as the result of that the number of patients with occurrence of signs of catarrhal gingivitis and secondary initial caries has increased. It is necessary to remotivate the patients on a month-to-month basis as well as to provide adequate health education in order to maintain initial results and to prevent the onset of occurrence of complications.

### SP 322

Friday, June 18 - at 12:15

#### EVALUATION OF MAXILLOMANDIBULAR WIDTH USING THREE-DIMENSIONAL CONE BEAM COMPUTED TOMOGRAPHIC IMAGES

**K-M Lee**, M-K Sun, H-S Hwang, J-H Cho, Department of Orthodontics, School of Dentistry, Chonnam National University, Gwangju, Korea

**AIM:** Rapid palatal expansion (RPE) devices are used correct maxillary transverse deficiency. The question that arises is how can maxillary transverse problems be evaluated to enable the use RPE? Traditionally, for analysis of maxillomandibular transverse differences, the two-dimensional (2D) posteroanterior (PA) cephalogram was used because direct measurement of bone width is impossible. Nowadays, three-dimensional (3D) cone-beam (CBCT) which decreases the radiation dose and 3D image software programs make it possible to have 3D measurements on maxillofacial area. The purpose of the present study was to measure maxillary and mandibular bone width on 3D CBCT image and to compare the maxillomandibular width of 3D image with that on a PA cephalogram.

**MATERIALS AND METHOD:** Images were taken of 30 adults with normal occlusion. Reference planes of the 3D images were established to be parallel to the functional occlusal plane at the level of the alveolar crest, furcation, root apex and 2 and 5 mm below root apex of the first molar. On each reference plane, the bone widths were measured at the maxillary and mandibular first molar and first and second premolar.

**RESULTS:** On comparison of the PA cephalograms, the maxillomandibular widths of 3D images of the first and second premolar did not show a statistically significant correlation, while the maxillomandibular widths at the first molar area showed a weak correlation.

**CONCLUSION:** 3D evaluation is mandatory to accurately measure maxillomandibular width.



# CLINICAL POSTERS



## LIST OF CLINICAL POSTERS

### List of Clinical Posters

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
CP 001	THEORETIC COMPARISON OF THREE ORTHODONTIC APPLIANCES <b>M Narimani</b> , Iran	147
CP 002	STATISTICAL ANALYSIS OF THE INDICES OF 500 PATIENTS <b>M Nagai</b> , Japan	147
CP 003	A BRIDGED ANCHOR USING MINI-IMPLANTS WITH A SLOT IN THE HEAD <b>I-G Kim</b> , Korea	147
CP 004	LICHENOID LESION IN AN ORTHODONTIC PATIENT <b>S Bonadeo</b> , Italy	147
CP 005	CLINICAL APPLICATION OF CAD/CAM TECHNOLOGY IN LINGUAL ORTHODONTICS <b>C Szuhaneek</b> , Romania	147
CP 006	RADIOGRAPHIC DIAGNOSIS IN GOLDENHAR SYNDROME – A THREE-DIMENSIONAL COMPUTED TOMOGRAPHIC STUDY <b>S Saccomanno</b> , Italy	148
CP 007	MECHANICAL PROPERTIES OF TITANIUM MOLYBDENUM ORTHODONTIC WIRES <b>T Fleser</b> , Romania	148
CP 008	INFLUENCE OF WELDING METHOD ON MECHANICAL PROPERTIES OF ORTHODONTIC METAL APPLIANCES <b>T Fleser</b> , Romania	148
CP 009	CLINICAL MANAGEMENT OF A PATIENT WITH EXTRUSIVE LUXATION DURING ORTHODONTIC TREATMENT <b>F Tokalak</b> , Turkey	148
CP 010	ORTHODONTIC EXTRUSION OF AN IMPACTED MANDIBULAR THIRD MOLAR WITH TEMPORARY ANCHORAGE DEVICES <b>A Nakhleh</b> , Italy	149
CP 011	LOCAL DISTRACTION OSTEOGENESIS AS THERAPY FOR POST-TRAUMATIC RESTRICTED MAXILLARY GROWTH <b>S Wriedt</b> , Germany	149
CP 012	THE LIP BUMPER AS A MOLAR DISTALIZATION APPLIANCE <b>B Ozel</b> , Turkey	149
CP 013	TWO TYPES OF PALATAL EXPANDERS IN THE TREATMENT OF UNILATERAL MAXILLARY CANINE ECTOPIA <b>D C Bratu</b> , Romania	149
CP 014	SPEECH-RELATED SHAPE CHANGES OF THE TONGUE IN SUBJECTS WITH AND WITHOUT A CLEFT LIP AND PALATE <b>T Ono</b> , Brazil	150
CP 015	MOLAR INTRUSION USING MINISCREWS IN AN ADULT PATIENT WITH A SKELETAL OPEN BITE – A CASE REPORT <b>A Kozanecka</b> , Poland	150
CP 016	SOFT-TISSUE LASERS – A NEW AID IN ORTHODONTICS <b>R Dimovska</b> , Former Yugoslav Republic of Macedonia	150
CP 017	ORTHODONTIC-PROSTHODONTIC TREATMENT OF SEVERE SKELETAL CLASS II PROTRUSION <b>S Rexhepi</b> , Former Yugoslav Republic of Macedonia	150
CP 018	THE VALUE OF BIOLOGICAL AGE FOR DETERMINING DENTOFACIAL ORTHOPAEDIC TREATMENT OPTIONS <b>E Teodorescu</b> , Romania	151
CP 019	OPEN BITE CORRECTION WITH MINISCREWS <b>A Sapač</b> , Slovenia	151
CP 020	EN MASSE RETRACTION WITH ORTHODONTIC MINI-IMPLANTS <b>Z Milosavljević</b> , Serbia	151
CP 021	AN UPRIGHTING AUXILIARY FOR DEEPLY IMPACTED MOLARS <b>M Greco</b> , Italy	151
CP 022	CONE BEAM COMPUTED TOMOGRAPHY AND PYKNODYSTOSIS – A CASE REPORT <b>L Laffranchi</b> , Italy	151
CP 023	ORTHODONTIC-SURGICAL CORRECTION OF A CLASS II DIVISION 1 PATIENT WITH FACIAL NERVE WEAKNESS <b>I Drerup</b> , Germany	152
CP 024	MAXILLARY CANINE-FIRST PREMOLAR TRANSPOSITION IN THE PERMANENT DENTITION. CLINICAL CASES <b>N Vasilakou</b> , Greece	152

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
CP 025	CAUSAL RELATION BETWEEN MALOCCLUSION AND ORAL MUSCLE DYSFUNCTION – A MODEL APPROACH <b>G Antonini</b> , Italy	152
CP 026	CONSIDERATION OF CANINE TEETH: DIAGNOSIS AND TREATMENT <b>S Saccomanno</b> , Italy	152
CP 027	MULTIDISCIPLINARY TREATMENT OF AN ADULT WITH MAXILLARY RETROGNATHISM AND OLIGODONTIA <b>D A Papamanou</b> , Greece	152
CP 028	SPEEDING UP ORTHODONTIC TREATMENT <b>L Barreto</b> , England	153
CP 029	EFFECTIVENESS OF FRAENECTOMY ON TONGUE FUNCTION AND POSTURE – AN ULTRASOUND EVALUATION <b>R Pavlović</b> , Slovenia	153
CP 030	MICROSCOPIC ANALYSIS OF THE ROOT SURFACE OF TEETH <b>C T Preoteasa</b> , Romania	153
CP 031	MODIFICATION OF DENTAL HARD TISSUES DURING ORTHODONTIC TREATMENT – A MICROSCOPIC ANALYSIS <b>C T Preoteasa</b> , Romania	153
CP 032	ORTHODONTIC TREATMENT OF LOWER SECOND MOLAR IMPACTION WITH SKELETAL ANCHORAGE – A CASE REPORT <b>A M Enache</b> , Romania	154
CP 033	SECONDARY CLEFT PALATE DENTAL PULP STEM CELL GRAFTING IN VITRO. A CONE BEAM COMPUTED TOMOGRAPHIC ASSESSMENT OF SCAFFOLD REABSORPTION <b>D Dalessandri</b> , Italy	154
CP 034	AN ORTHODONTIC APPROACH IN TREATING BICONDYLAR FRACTURES <b>E Çerekja</b> , Turkey	154
CP 035	ORTHODONTIC TREATMENT OF A CLASS II MALOCCLUSION IN AN ADULT SUBJECT <b>M Stojanović</b> , Serbia	155
CP 036	THREE-DIMENSIONAL ASSESSMENT OF CLASS III TREATMENT – A CASE REPORT <b>S Lah Kravanja</b> , Slovenia	155
CP 037	EARLY TREATMENT OF A CLASS III MALOCCLUSION WITH A DELAIRE TYPE FACE MASK – A CASE REPORT <b>S Lah Kravanja</b> , Slovenia	155
CP 038	NON-SURGICAL TREATMENT OF A JUVENILE PATIENT WITH PYCNODYSOSTOSIS AND A CLASS III MALOCCLUSION <b>J Berekzai</b> , Germany	155
CP 039	CALIBRATED DOUBLE-SPLIT-CAST MÜNSTERANER MODEL-OPERATION SYSTEM <b>T Ziebur</b> , Germany	156
CP 040	USE OF TEMPORARY ANCHORAGE DEVICES TO CORRECT SEVERE CLASS III MALOCCLUSIONS, WITHOUT ORTHOGNATHIC SURGERY <b>D Mahony</b> , Australia	156
CP 041	ORTHOGNATHIC TREATMENT OF A SEVERE CLASS III MALOCCLUSION: A CASE REPORT <b>E Baharoğlu</b> , Turkey	156
CP 042	DENTO-SKELETAL CHANGES IN A CLASS III MALOCCLUSION SUBJECT WITH AN OPEN BITE – A CASE REPORT <b>A Fekonja</b> , Slovenia	156
CP 043	ORTHODONTIC TREATMENT OF A PATIENT WITH FRONTO NASAL DYSPLASIA <b>H Kan</b> , Turkey	157
CP 044	REVERSE HEADGEAR WITH A DISTALIZATION SCREW – A CASE REPORT <b>E K Atamozlu</b> , Turkey	157
CP 045	MAXILLARY PROTRACTION WITH TWO BONE ANCHORS IN A CLASS III PATIENT WITH OLIGODONTIA <b>E Dumu Dundu Jr.</b> , Belgium	157
CP 046	UPPER MOLAR ALIGNMENT WITH A REVERSE RAPID MAXILLARY EXPANSION APPLIANCE – A THREE-DIMENSIONAL CAST ANALYSIS*** <b>I Drerup</b> , Germany	157
CP 047	ORTHOGNATHIC SURGERY PLANNING ON A THREE-DIMENSIONAL SOLID MODEL <b>M Erkan</b> , Turkey	157
CP 048	ASYMMETRIC PREMAXILLARY DISTRACTION OSTEOGENESIS – A CASE REPORT <b>Ş Karaçay</b> , Turkey	158
CP 049	A MODIFIED CLASS III PROTRACTION MASK ILLUSTRATED BY TWO CLINICAL CASES <b>N Verly</b> , Belgium	158
CP 050	TREATMENT OF A CLASS III MALOCCLUSION WITH THE FRÄNKEL FUNCTIONAL REGULATOR – A CASE REPORT <b>Z Stamenković</b> , Serbia	158
CP 051	EVALUATION OF EARLY CLASS III TREATMENT WITH THE FRÄNKEL APPLIANCE <b>M Lajh</b> , Slovenia	158
CP 052	EARLY TREATMENT OF A TRUE FORCED BITE WITH THE FRÄNKEL APPLIANCE <b>D Kramberger</b> , Slovenia	159
CP 053	EARLY TREATMENT OF A CLASS III MALOCCLUSION WITH A MODIFIED MAXILLARY PROTRACTION FACEMASK <b>M Stojković</b> ,	159
CP 054	SUCCESSFUL ORTHOGNATHIC TREATMENT OF A SKELETAL CLASS III PATIENT USING THREE-DIMENSIONAL SIMULATION <b>I-R Choi</b> , Korea	159
CP 055	USE OF BONE ANCHORS IN A CLASS III SKELETAL PATIENT. A CASE REPORT <b>M Charezinski</b> , Belgium	159

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
CP 056	COMBINED ORTHODONTIC-SURGICAL THERAPY IN A CLASS III PATIENT WITH LATEROGNATHIA <b>M Janošević</b> , Serbia	160
CP 057	TRANSVERSE AND SAGITTAL EFFICACY OF A MODIFIED PALATAL EXPANDER IN SKELETAL CLASS III TREATMENT <b>Rexhepi</b> , Kosova	160
CP 058	MODERATE RELAPSE AFTER MANDIBULAR OSTEOTOMY IN SKELETAL CLASS III SUBJECTS <b>A Ismani</b> , Former Yugoslav Republic of Macedonia	160
CP 059	EARLY TREATMENT OF A CLASS III ANTERIOR SKELETAL CROSSBITE WITH AN ORAL SHIELD <b>M Yanagisawa</b> , Japan	160
CP 060	SOFT TISSUES PROFILE CHANGES DURING TREATMENT OF GROWING CLASS III PATIENTS <b>E Teodorescu</b> , Romania	160
CP 061	TREATMENT OF A CLASS III MALOCCLUSION WITH A COMBINATION OF FRÄNKEL AND FIXED APPLIANCES <b>E Zaletel</b> , Slovenia	161
CP 062	EARLY TREATMENT OF A CLASS III MALOCCLUSION WITH THE MODIFIED FRÄNKEL APPLIANCE <b>S Melink</b> , Slovenia	161
CP 063	TREATMENT OF A CLASS III SUBJECT WITH AN OPEN BITE IN MIXED DENTITION WITH A MODIFIED FRÄNKEL APPLIANCE <b>M Malinger</b> , Slovenia	161
CP 064	TREATMENT OF CLASS III MALOCCLUSIONS – THE KEY ROLE OF THE OCCLUSAL PLANE <b>B Grollemund</b> , France	161
CP 065	A SEVERE CLASS III MALOCCLUSION PATIENT TREATED WITH QUADHELIX <b>R Janev</b> , Former Yugoslav Republic of Macedonia	162
CP 066	THE CLEAR ALIGNER TECHNIQUE IN THE TREATMENT OF PRE-SURGICAL CLASS III MALOCCLUSION SUBJECTS <b>M Greco</b> , Italy	162
CP 067	DESIGN AND DEVELOPMENT OF A FACEMASK FITTED WITH A TIMING DEVICE <b>C Schwartz</b> , France	162
CP 068	EARLY ORTHOPAEDIC TREATMENT OF A CLASS III MALOCCLUSION – A THREE-DIMENSIONAL EVALUATION <b>V Božnik</b> , Slovenia	162
CP 069	PROFESSIONALS' AND LAYPERSONS' APPRECIATION OF CLASS III SURGICAL CORRECTION OPTIONS <b>P Christou</b> , Switzerland	163
CP 070	CONTEMPORARY EVALUATION OF TONGUE FUNCTION AND POSTURE – A CASE REPORT <b>D Kramberger</b> , Slovenia	163
CP 071	EARLY MAXILLARY PROTRACTION WITH A NEW IMPLANT-SUPPORTED NON-COMPLIANCE AND AESTHETIC DEVICE <b>L Favero</b> , Italy	163
CP 072	INTERDISCIPLINARY TREATMENT OF A CLASS III MALOCCLUSION – A CASE REPORT*** <b>G Spaić</b> , Slovenia	163
CP 073	FORWARD TRACTION ON THE MAXILLA IN CLEFT LIP AND PALATE PATIENT*** <b>K Verhovec</b> , Slovenia	163
CP 074	THREE STAGE TREATMENT OF A SKELETAL CLASS III MALOCCLUSION <b>K Arko Kampus</b> , Slovenia	164
CP 075	COMPARISON OF AN UNTREATED CLASS III MALOCCLUSION SUBJECTS COMPARED WITH AN 'AVERAGE' PATIENT– A THREE-DIMENSIONAL EVALUATION <b>A Marion</b> , Slovenia	164
CP 076	ORTHOGNATHIC TREATMENT OF A CLASS III MALOCCLUSION – A CASE REPORT <b>B Mady Maričić</b> , Croatia	164
CP 077	APPLICATION OF ORTHODONTIC MINISCREW-IMPLANTS IN OLIGODONTIA – A CASE REPORT <b>K Raftowicz-Wójcik</b> , Poland	165
CP 078	THE IMPORTANCE OF INTERCEPTIVE ORTHODONTIC TREATMENT <b>J Andjelic</b> , Montenegro	165
CP 079	MODIFIED SLIDING MECHANICS IN EXTRACTION CASES WITH A BIDIMENSIONAL APPROACH <b>P Mozzicato</b> , Italy	165
CP 080	PATHOLOGY IN CRANIOCERVICAL JUNCTION – WHAT DOES IT MEAN? <b>H Arponen</b> , Finland	165
CP 081	IMPROVED SUPERELASTIC NICKEL TITANIUM WIRE FOR TREATMENT OF AN IMPACTED TOOTH WITH AN UNDEVELOPED ROOT <b>C-C Chen</b> , Taiwan	166
CP 082	CLINICAL AND RADIOGRAPHIC IMPROVEMENT DURING ORTHODONTIC TREATMENT IN PERIODONTAL PATIENTS <b>F Aguilar</b> , Mexico	166
CP 083	CEPHALOMETRIC AND DENTAL EVALUATION OF PATIENTS WITH IMPACTED LOWER SECOND MOLARS <b>A Pilotto</b> , Italy	166
CP 084	PROFILE CHANGES IN A CLASS II DIVISION 1 PATIENT AFTER TWO-PHASE ORTHODONTIC TREATMENT <b>E Baytorun</b> , Turkey	166
CP 085	BIOPROGRESSIVE TREATMENT OF AN ANGLE CLASS II DIVISION 2 PATIENT BY MANDIBULAR INCISOR EXTRACTION – CASE REPORT <b>A Aksoy Doğan</b> , Turkey	166
CP 086	CRANIOFACIAL GROWTH AND DEVELOPMENT IN PATIENTS WITH AMELOGENESIS IMPERFECTA <b>I Eleftheriadi</b> , Greece	166

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
CP 087	MICROGNATHIA AS A PROGNOSTIC FACTOR FOR OBSTRUCTIVE SLEEP APNOEA IN PATIENTS WITH SYNDROMES <b>F Synodinos</b> , Greece	167
CP 088	POSTERIOR MAXILLARY OSTEOTOMY TO RE-ESTABLISH INTERMAXILLARY SPACE BEFORE PROSTHODONTICS <b>A Aneiros Ardao</b> , Spain	167
CP 089	INTERDISCIPLINARY TREATMENT OF MULTIPLE AGENESIS – A CASE REPORT <b>A Peniche</b> , México D.F	167
CP 090	HOW TO TORQUE A PALATALLY IMPACTED CANINE <b>M Araghbidi Kashani</b> , Iran	167
CP 091	FORCED ERUPTION – INTRODUCING AN EASY METHOD <b>S Torkan</b> , Iran	168
CP 092	PRE-PROSTHODONTIC ORTHODONTIC TREATMENT IN IMPLANT SUPPORTED RESTORATIONS <b>A Atay</b> , Turkey	168
CP 093	CAMOUFLAGE TREATMENT OF A PATIENT WITH A CLASS II DIVISION 1 MALOCCLUSION – A CASE REPORT <b>Z Stamenković</b> , Serbia	168
CP 094	ORTHODONTIC AND SURGICAL TREATMENT OF IMPACTED MANDIBULAR SECOND PREMOLARS <b>F Sali</b> , Kosovo	168
CP 095	COMBINED ORTHODONTIC-IMPLANT TREATMENT OF CONGENITALLY MISSING LATERAL INCISORS <b>M Stojanović</b> , Serbia	169
CP 096	HERBST APPLIANCE IN THE LATE TREATMENT OF CLASS II DIVISION 1 MALOCCLUSION – A CASE REPORT <b>N Nedeljković</b> , Serbia	169
CP 097	THE ORTHODONTIC MANAGEMENT OF CLEFT LIP AND PALATE PATIENTS <b>E Hussein</b> , Palestinian Territory	169
CP 098	A MODIFIED ELN APPLIANCE IN OPEN BITE TREATMENT OF GROWING PATIENTS – A PRELIMINARY STUDY <b>F Tomassini</b> , Italy	169
CP 099	LONG-TERM RETENTION AFTER TREATMENT OF SKELETAL CLASS II GROWING PATIENTS WITH A DEEP BITE*** <b>J-H Kim</b> , South Korea	169
CP 100	EPIDEMIOLOGY OF CLEFT LIP AND PALATE IN THE FORMER YUGOSLAV REPUBLIC OF MACEDONIA <b>R Dimovska</b> , Former Yugoslav Republic of Macedonia	170
CP 101	A SEVERE COMPLICATION OF BONDED MANDIBULAR LINGUAL RETAINERS <b>P Pazera</b> , Switzerland	170
CP 102	AN ALTERNATIVE METHOD FOR OVEREXTRUDED SECOND MOLARS IN ADULT PATIENTS <b>J E Vlachojanis</b> , USA	170
CP 103	TREATMENT OF A TOOTH WITH A COMPLICATED CROWN-ROOT FRACTURE – A CASE REPORT <b>B Bajraktarova</b> , Former Yugoslav Republic of Macedonia	170
CP 104	DISTORTED SHAPE AND SIZE OF TEETH – A CHALLENGE FOR CONTEMPORARY ORTHODONTIC TREATMENT <b>A Sieja</b> , Poland	171
CP 105	INTERDISCIPLINARY TREATMENT: ORTHODONTIC EXTRUSION WITH MINI-IMPLANTS <b>A Peniche</b> , México D.F	171
CP 106	RELATIONAL DEVELOPMENT IN BABIES WITH CLEFT LIP AND PALATE: A STUDY CARRIED OUT IN FRANCE <b>B Grollemund</b> , France	171
CP 107	USING CASTILLO MORALES STIMULATION PLATE FOR A PATIENT WITH APERT SYNDROME – A CASE REPORT <b>A Ifert</b> , Germany	171
CP 108	LONG-TERM RETENTION AFTER TREATMENT OF A DEEP BITE IN NON-GROWING PATIENTS <b>J-H Kim</b> , South Korea	171
CP 109	TREATMENT OF CLASS II DIVISION 1 MALOCCLUSION, THROUGH COMBINATION OF FIXED AND MOBILE APPLIANCE <b>E Çitaku</b> , Kosova	172
CP 110	GERM EXTRACTION – A THERAPEUTIC APPROACH FOR DENTOFACIAL ORTHOPAEDICS USING MINI-IMPLANTS <b>H Sarabia</b> , Mexico	172
CP 111	SIMULTANEOUS FUNCTIONAL MAGNETIC RESONANCE IMAGING – ELECTROMYOGRAPHIC MEASUREMENT DURING TOOTH TAPPING <b>J Miyamoto</b> , Japan	172
CP 112	RETENTION: A CASE PRESENTATION FOR THE POSTGRADUATE COURSE AT TOKYO DENTAL COLLEGE <b>M Mukai</b> , Japan	172
CP 113	MANDIBULAR SYMPHYSIS ADVANCEMENT WITH TOOTH - AND BONE-BORNE DISTRACTION DEVICES <b>F Çakmak</b> , Turkey	173
CP 114	CONGENITAL ABNORMALITIES OF THE EXTERNAL EAR IN CLINICAL ORTHODONTIC EXAMINATION <b>N Vasilakou</b> , Greece	173
CP 115	PREFORMED DEVICES: TEN-YEAR'S EXPERIENCE <b>D Tarantino</b> , Italy	173
CP 116	ORODENTAL, CRANIOFACIAL AND GENERAL CHARACTERISTICS OF LACRIMO-AURICULO-DENTO-DIGITAL SYNDROME <b>D A Papamanou</b> , Greece	173
CP 117	THE EFFECTS OF THE PENDEX APPLIANCE IN A CLASS II DIVISION 2 MALOCCLUSION SUBJECT <b>T Moreira</b> , Portugal	173

<b>Abstract No</b>	<b>Title, Presentator, Country</b>	<b>Page</b>
CP 118	A DILACERATED UPPER INCISOR IN A YOUNG PATIENT – ORTHODONTIC TREATMENT <b>S Tronca</b> , Italy	174
CP 119	RELATIONSHIP BETWEEN RAPID PALATAL EXPANSION AND SAGITTAL DISCREPANCIES <b>F Iazzetti</b> , Italy	174
CP 120	ORTHODONTIC TREATMENT OF INCOMPLETE TRANSPOSITION IN THE LATE MIXED DENTITION <b>Ş Karaçay</b> , Turkey	174
CP 121	SEGMENTAL MANDIBULAR DENTOALVEOLAR DISTRACTION: AN OPTION TO EXTRACTION THERAPY <b>D Esercizio</b> , Italy	174



### Biotechnology and the Future in Orthodontics

#### CP 001

Wednesday, June 16 - at 15:15

##### THEORETIC COMPARISON OF THREE ORTHODONTIC APPLIANCES

**M Narimani**, Shahed Medical Science, Tehran, Iran

With the extension of different sciences, innovations in orthodontic treatment occur. The importance of such innovations cannot be denied in specialty courses. With the introduction of a new method, questions arise as to the effectiveness compared with conventional ideas,

This presentation will show new appliances; their characteristics from the point of view of bracket forming concept, bracket characteristics, type of required wires and pliers and the treatment stages will be reviewed.

#### CP 002

Wednesday, June 16 - at 15:15

##### STATISTICAL ANALYSIS OF THE INDICES OF 500 PATIENTS

**M Nagai**, Miki Dental Clinic, Nagano, Japan

**AIM:** It has previously been shown that there is no significant difference in the Pont Indices of 21|12 between European and Japanese patients. The aim of this study was to consider the relationship between 21|12 and the ranges of indices of 4|4 and 6|6 corresponding to indexes of 21|12 between 26 and 38 mm at 2 mm intervals, as the Bimler Stomatopedic Treatment Sheet is based on 2 mm increments.

**MATERIALS AND METHOD:** The database of approximately 500 Japanese patients. Statistical processing of 437 patients was carried out after measurement of the patients' original casts. Measurements were carried out of the distal pouch diameter of 4|4 and the fovea centralis diameter of 6|6 on the patients' initial casts after or during treatment. Finally, the average indices of 4|4 and 6|6 per each 2 mm range of 21|12 between 26 and 38 mm were computed.

**RESULTS:** Analysis of this data collected on the Bimler Stomatopedic Treatment Sheets for these patients yielded interesting results. In spite of a variation of 11 mm in the Pont index of the subjects, no significant variance was measured in the indexes of 4|4 or of 6|6.

**CONCLUSION:** The significance of these results needs to be considered further.

#### CP 003

Wednesday, June 16 - at 15:15

##### A BRIDGED ANCHOR USING MINI-IMPLANTS WITH A SLOT IN THE HEAD

**I-G Kim**, Gajirun-e Orthodontic Clinic, Gwangju, Korea

The use of mini-implant anchorage has become more widespread. At the beginning, direct-anchorage biomechanics using a single mini-implant was popular; however it was not possible to change the vector between the mini-implant and the target tooth groups. To overcome this limitation, the mini-implant bridged anchor was invented. For mini-implant bridged anchorage cases, the archwire can be inserted in a slot in the head. This type of anchorage can be used to stabilize a group of teeth or to connect two mini-implants, creating an implant-dental or inter-implant anchorage

unit. A bridged anchor using mini-implants makes it possible to efficiently control molar protraction, incisor intrusion, molar retraction and scissor bite correction.

#### CP 004

Wednesday, June 16 - at 15:15

##### LICHENOID LESION IN AN ORTHODONTIC PATIENT

**S Bonadeo**, C Tian, D Dalessandri, F Zotti, L Laffranchi, Department of Orthodontics, University of Brescia, Italy

**AIM:** The association between oral mucosal lesions and metals has been documented. Mercury, amalgam, nickel, palladium, chromium, cobalt, gold, indium, EGDMA resin and chloroform can cause lichenoid reactions in oral mucosa adjacent to or in contact with prosthetic or orthodontic appliances.

**SUBJECT AND METHOD:** A patient undergoing orthodontic appliance treatment with an asymptomatic reticular white lesion on the left buccal mucosa. The patient did not report any systemic disease in progress. Testing for intolerance to dental metals and dental resins was carried out using the cutaneous patch-test. Subsequently a biopsy was performed on the mucosal lesion.

**RESULTS:** Patch testing showed intolerance to nickel. The biopsy report described a histological appearance with irregular acanthosis of the epithelium with underlying moderate interstitial and perivascular mononucleated inflammation, with lichenoid attitude and minimal vacuoles degeneration of the basal layer epithelium, thickening of the basement membrane, and atrophy of the epithelial ridges. Direct immunofluorescence showed an irregular deposit (non-linear) of fibrinogen at the basement membrane.

**CONCLUSION:** Patients with orthodontic appliances could manifest lichenoid lesion. Tests of intolerance to metals and resins and associated biopsy analysis are required. Therapy is not usually necessary because symptoms are not always present or they disappear once the cause has been removed.

#### CP 005

Wednesday, June 16 - at 15:15

##### CLINICAL APPLICATION OF CAD/CAM TECHNOLOGY IN LINGUAL ORTHODONTICS

**C Szuhaneck**, F Glavan R Jianu, A Muntean, University of Medicine and Pharmacy, Timisoara, Romania

**AIM:** To evaluate the efficacy of customized lingual orthodontics in successful treatment.

**SUBJECTS AND METHOD:** Seven patients with different types of malocclusion all treated with customized lingual orthodontic appliances. The efficiency of this type of treatment was evaluated by means of bracket debonding, patient comfort, oral hygiene and predictability of the results.

**RESULTS:** The treatment objectives were achieved in all cases. The duration of treatment depended on the severity of the malocclusion. The faster results were obtained in subjects with crowding or deep bites. The rate of bracket debonding was very low and patient comfort and oral hygiene showed good results.

**CONCLUSION:** Customized lingual brackets show a high clinical efficiency and represent an important part of the future in aesthetic orthodontics.

**CP 006***Wednesday, June 16 - at 15:15***RADIOGRAPHIC DIAGNOSIS IN GOLDENHAR SYNDROME – A THREE-DIMENSIONAL COMPUTED TOMOGRAPHIC STUDY****S Saccomanno**, F Greco, L Cacucci, C Notaristefano, R Deli, Department of Orthodontics Catholic University of the Sacred Heart, Rome, Italy

**AIM:** Goldenhar syndrome is a congenital condition that includes anomalies of the derivatives of the first and second brachial arches, vertebral defects, and ocular abnormalities, generally epidermoids. The cause of the condition remains unclear but seems to be heterogeneous. Most cases are sporadic. However, familiar cases and concordant monozygotic twins have been reported, suggesting a genetic origin. In addition, several chromosomal abnormalities have been documented in affected individuals. The frequency of this syndrome is estimated as one per 5600 to 26550 births and the male:female ratio is about 3:2 more often affects right hemifacies (right:left-3:2).

**MATERIALS AND METHOD:** Three-dimensional TAC evaluation showed, in young patients, condylar asymmetry. In particular, the right condyle is dimorphic, short and wide; the auricle of the right external ear is ipoplastic, only a dimorphic rough is present; the right middle ear has an ipoplastic tympanic cavity; the right internal ear has an atresico uditive conduct.

**CONCLUSION:** 3D TAC is a powerful diagnostic instrument and offers many advantages including volumetric reproduction of the cranium and soft tissues, no overlap of anatomic parts, high precision of the images, and a constant and easily reproducible reference system.

**CP 007***Wednesday, June 16 - at 15:15***MECHANICAL PROPERTIES OF TITANIUM MOLYBDENUM ORTHODONTIC WIRES****T Fleşer**<sup>1</sup> C Szuhaneş<sup>2</sup>, C Bortun<sup>2</sup>, F Glavan<sup>2</sup>, A Cheveresan<sup>2</sup>, <sup>1</sup>Mechanical Engineering Faculty and <sup>2</sup>Faculty of Dental Medicine, University of Medicine and Pharmacy, Timisoara, Romania

**AIM:** Orthodontic treatment is mainly dependent on the loads developed by different wires. The aim of this study was to compare the mechanical properties of several titanium molybdenum (TMA) archwires now available on the market.

**MATERIALS AND METHOD:** The mechanical properties of the wires were tested during bending and torsion. The results were statistically analyzed.

**RESULTS:** For the same archwire size, the spring back and stiffness rates during bending and torsion varied significantly from one manufacturer to another.

**CONCLUSION:** Optimal use of orthodontic wires can be achieved by careful selection of the appropriate wire type and size to meet the demands of a particular clinical situation.

**CP 008***Wednesday, June 16 - at 15:15***INFLUENCE OF WELDING METHOD ON MECHANICAL PROPERTIES OF ORTHODONTIC METAL APPLIANCES****T Fleşer**<sup>1</sup> C Szuhaneş<sup>2</sup>, F Glavan<sup>2</sup>, A Muntean<sup>2</sup>, C Bortun<sup>2</sup>, <sup>1</sup>Mechanical Engineering Faculty and <sup>2</sup>Faculty of Dental Medicine, University of Medicine and Pharmacy Timisoara, Romania

**AIM:** Joining of metal appliances is frequently necessary to create individual orthodontic appliances and to achieve efficient treatment procedures. The purpose of this study was to evaluate the mechanical strength and the microhardness of joints welded with universal silver solder and with the ultrasonic technique.

**MATERIALS AND METHOD:** The joints were produced by various dental technicians using three different soldering techniques. The tensile strengths were measured with a universal testing machine, while smooth surface and characteristic nugget formation were microscopically evaluated. The results were statistically analysed.

**RESULTS:** The ultrasonic welding technique provided superior strength results compared with the silver solder method.

**CONCLUSION:** Ultrasonic welding is an expensive but efficient technique.

**CP 009***Wednesday, June 16 - at 15:15***CLINICAL MANAGEMENT OF A PATIENT WITH EXTRUSIVE LUXATION DURING ORTHODONTIC TREATMENT****F Tokalak**, M Ozer, Department of Orthodontics, Faculty of Dentistry, University of Ondokuz Mayıs, Samsun, Turkey

**AIM:** Traumatic dental injuries are a serious health problem in orthodontics. Injuries are classified into various fracture and luxation types; one being extrusive luxation which can be described as displacement of the tooth in an extrusive direction.

**SUBJECT AND METHOD:** A 13-year-old girl undergoing fixed orthodontic treatment who requested an emergency appointment. After arrival, she reported falling on her face 14 hours previously. Clinically the upper left central, upper left lateral and upper right central were partially extruded from their sockets. The extruded teeth were elongated from the occlusal plane in association with bleeding from the periodontal ligament, there was pain during occlusion and percussion of the teeth, normal occlusion was prevented because of the premature contacts of the elongated incisors. Radiographically all upper incisors, except the upper right lateral, were diagnosed as having extrusive luxation. All the extruded incisors were replanted with finger pressure, a passive 0.017 × 0.025 beta titanium archwire was placed and no additional orthodontic force applied during the following three months. To prevent occlusal trauma, glass ionomer cement was placed on the lower first molars. A follow-up protocol one year after the end of treatment and six months after the end of active treatment was completed.

**RESULTS:** All the extruded incisors were successfully reimplemented and vitality tests were normal. There was no sign of external or internal root resorption and the periodontal health of the teeth was normal at the 18 month follow-up.

**CONCLUSION:** Emergency replacement, with correct orthodontic management, increases the survival rate of extruded teeth.



## Contemporary Diagnostics and Treatment Approaches

## CP 010

Thursday, June 17 - at 15:15

**ORTHODONTIC EXTRUSION OF AN IMPACTED MANDIBULAR THIRD MOLAR WITH TEMPORARY ANCHORAGE DEVICES****A Nakhleh**, S P Isaza Penco, M Di Lorenzo, Private Practice, Bologna, Italy

**AIM:** The reason for mandibular third molar impaction is due to inadequate space between the distal of the second mandibular molar and the anterior border of the mandibular ascending ramus. The purpose of this study was to evaluate the possibility of success in replacing an impacted mandibular third molar in the extraction site of a second molar.

**SUBJECT AND METHOD:** A 34-year-old female referred due to an iatrogenic fracture of the distal root of the second mandibular molar during a failed attempt at mandibular third molar extraction. The patient was a skeletal and dental Class I normodivergent, no temporomandibular disorder, no myofunctional problems who had undergone previous orthodontic treatment. The mandibular third molar was fully developed and without signs of ankylosis. The treatment plan included removal of the mandibular second molar root and the repositioning of the mandibular third molar. An 8 mm self tapping miniscrew Tomas (Dentaurum, Ispringen, Germany) was inserted in the mandible between the first and the second premolar. A Discovery 0.22 appliances and direct tubes (Ortho Cast mini, Dentaurum) were bonded to the mandibular second premolar and first and third molars. Controls during treatment were made with intraoral radiographs and dental pantomography. After 12 months of treatment the third molar was completely replaced in the position of the second molar.

**RESULTS:** The clinical results were satisfactory with temporary anchorage devices. Patient compliance was not necessary and the treatment was minimally invasive.

**CONCLUSION:** An impacted mandibular third molar could successfully replace a lost mandibular second molar. Such treatment avoids dental implant rehabilitation and surgical third molar extraction.

## CP 011

Thursday, June 17 - at 15:15

**LOCAL DISTRACTION OSTEOGENESIS AS THERAPY FOR POST-TRAUMATIC RESTRICTED MAXILLARY GROWTH****S Wriedt**<sup>1</sup>, B Al-Nawas<sup>2</sup>, S Löblein<sup>1</sup>, H Wehrbein<sup>1</sup>, W Wagner<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Maxillo-Facial-Surgery, J. Gutenberg-University, Mainz, Germany

**AIM:** Post-traumatic ankylosis of upper incisors during childhood often causes long-term restriction of growth in the traumatized area. The aim of this presentation is to show the treatment of such a patient.

**SUBJECT AND METHOD:** A nine-year-old boy with an alveolar bone fracture and intrusion of the upper incisors as a result of trauma; it was not possible to align the traumatized teeth because of post-traumatic ankylosis. The anterior open bite was first masked by lengthening teeth 12 to 21 with composite. Because of further growth the bite re-opened. At 14 years of age a palatal implant was inserted and 6 months later spaces distal to the ankylosed teeth were opened without loss of anchorage using a pala-

tal implant supported transpalatal arch (TPA). The TPA was then extended to the labial and palatal area of the traumatized teeth. A segment of the alveolar process containing the ankylosed teeth and reaching to the nasal floor was osteotomized using the piezosurgical technique. Over a 21 day period the segment was moved 10 mm in an occluso-labial direction using a power of approximately 500 P as defined by bone floating. The direction of force was controlled daily to prevent early closure of the osteotomy line. After achieving the planned position, the segment was fixed with composite and a stiff wire.

**CONCLUSION:** Modified distraction osteogenesis supported on a palatal implant was predictable and free of side-effects to develop segments of alveolar bone, which are restricted in growth because of traumatic injuries.

## CP 012

Thursday, June 17 - at 15:15

**THE LIP BUMPER AS A MOLAR DISTALIZATION APPLIANCE****Z Novruzov**<sup>1</sup>, A Koklu<sup>2</sup>, **B Ozel**<sup>3</sup>, <sup>1</sup>Department of Paediatric Stomatology Azerbaijan Medical University, Baku, Azerbaijan and Departments of Orthodontics, <sup>2</sup>Ankara University and <sup>3</sup>Karadeniz Technical University, Turkey

**AIM:** A Class II division 2 malocclusion is usually characterized by over eruption of the permanent mandibular incisors that occlude to the posterior of the cingulum plateau of the retroclined permanent maxillary central incisors. This results in a reduced or 'hidden' overjet and a pronounced overbite. The aim of this presentation is to describe the molar distalization procedure in a patient with a Class II division 2 malocclusion.

**MATERIALS AND METHOD:** Treatment was initiated with a lip bumper in the upper jaw through the upper first molar teeth and Begg fixed appliances in the lower arch. This was done in order to utilize the tonus of the upper lip, which has been reported to be increased in deep overbite cases. The upper molar teeth were distalized until a bilateral Class I molar relationship was accomplished. Fixed appliance treatment was then initiated in the upper arch and the orthodontic treatment was finished successfully in an overall period of 1 year 6 months.

**CONCLUSION:** The use of increased upper lip tonicity may be an option for upper molar distalization for the treatment of deep bite subjects.

## CP 013

Thursday, June 17 - at 15:15

**TWO TYPES OF PALATAL EXPANDERS IN THE TREATMENT OF UNILATERAL MAXILLARY CANINE ECTOPIA****D C Bratu**, F Agache, A Ogodescu, Department of Paedodontics and Orthodontics, Faculty of Dental Medicine, University 'Victor Babeş', Timişoara, Romania

**AIM:** To present the treatment of unilateral maxillary canine ectopia in two patients with a severe lack of space for correct alignment in the dental arch, using two types of rapid palatal expanders (RPE) with spider screw (Leone, Italy) and two arms (Veltri, Italy).

**SUBJECTS AND METHOD:** An 11-year-old girl in the mixed dentition with unilateral maxillary canine ectopia, a Class I skeletal pattern, brachycephalic, excessive overbite, midline deviation, treat-

ed with a RPE with two arms. The second subject was a 12-year-old girl with the same anomaly treated with a spider screw RPE. At the end of the RPE both patients were treated with fixed appliances, using Roth OMNI brackets, 0.022 inch slot, straightwire technique. The activation scheme was one-quarter turn three times a day, for 9 and 12 days respectively, beginning 24 hours after the device was cemented.

**RESULTS:** After 2 years of orthodontic treatment the canines were successfully aligned in the correct position in the dental arch. The other dentoalveolar aspects were corrected and a neutral relationship in all three reference planes was obtained. Postero-anterior cephalograms analyzed for both patients pre-treatment and immediately after RPE, showed opening of the midpalatal suture.

**CONCLUSION:** The spider screw allows opening of the anterior area and maintains the posterior diameter. The two arm RPE offers increased comfort for the patient, it is easy to construct and to cement, and is also hygienic. Patients treated before the pubertal peak exhibit significantly more effective long-term changes at the skeletal level. If the treatment is finalised after skeletal growth, the gain is dentoalveolar.

## CP 014

Thursday, June 17 - at 15:15

### SPEECH-RELATED SHAPE CHANGES OF THE TONGUE IN SUBJECTS WITH AND WITHOUT A CLEFT LIP AND PALATE

**T Ono**<sup>1,2</sup>, M S Inoue-Arai<sup>1,2,3</sup>, I W Ng<sup>1,2</sup>, J-I Takada<sup>1,2</sup>, K Moriyama<sup>1,2</sup>,  
<sup>1</sup>Maxillofacial Orthognathics, Graduate School, Tokyo Medical and Dental University and <sup>2</sup>GCOE, International Research Center for Molecular Science in Tooth and Bone Diseases, Japan, <sup>3</sup>Universidade Estadual de Maringá, Brazil

**AIM:** To delineate the geometric change in the tongue during plosives with different places of articulation in subjects with and without a cleft lip and palate (CLP) using magnetic resonance imaging (MRI) movie.

**SUBJECTS AND METHOD:** One unilateral (UCLP) and two bilateral (BCLP) CLP subjects and 10 adult subjects without a CLP. One BCLP subject showed a compensatory misarticulation (CMA) during a velar plosive. Custom-made circuitry was connected to a MRI apparatus to enable an external trigger pulse to control the timing of the scanning sequence and to provide an auditory cue for synchronization of the subject's utterances. The subject repeated bilabial and velar plosives, and the run was measured using a gradient echo sequence with a repetition time of 30 ms. Geometric changes in tongue shape were analyzed by measuring distances from the centre of gravity (CoG) and the contours of the tongue. The difference in the trajectory of the CoG between subjects with and without a CLP was statistically compared.

**RESULTS:** The tip of the tongue elongated during a bilabial plosive in all subjects. The posterior dorsum of the tongue elevated during a velar plosive in subjects without a CLP and CLP subjects without CMA, while such geometric changes were not observed in the BCLP subject with CMA. The trajectory of the CoG during both plosives was similar in subjects with and without a CLP and without CMA, but not in BCLP subject with CMA.

**CONCLUSION:** Detailed geometric changes in tongue shape in subjects with and without a CLP were revealed by MRI. Moreover, such changes were different between subjects with and without CMA, irrespective of the presence/absence of a CLP.

## CP 015

Thursday, June 17 - at 15:15

### MOLAR INTRUSION USING MINISCREWS IN AN ADULT PATIENT WITH A SKELETAL OPEN BITE – A CASE REPORT

M Szumielewicz, J Antoszewska, **A Kozanecka**, Department of Maxillofacial Orthopaedics and Orthodontics, Wrocław Medical University, Poland

**AIM:** A severe anterior open bite (AOB) is considered to be one of the most difficult malocclusions to treat. Molar intrusion with

skeletal anchorage has recently been suggested to avoid orthognathic surgery in such cases. The aim of this presentation is to show the effectiveness of miniscrews for intrusion of molars in a skeletal open bite in adult.

**SUBJECT AND METHOD:** A 29-year-old patient with increased lower anterior face height, a convex profile, an AOB of 1.5 mm, a narrowed maxilla, crowding, a Class I molar relationship and a Class II/III canine relationship on the left and right sides, respectively. The treatment plan was extraction of all third molars and second premolars, molar intrusion using miniscrews, retraction of the upper and lower incisors and canines, controlled molar mesialization, and retention. Angular and linear cephalometric measurements before and after treatment were compared.

**RESULTS:** Comparison of pre- and post-treatment angular values revealed a 1.3 degree decrease of ML-NSL angle (from 41.4 to 40.1°), a 1.0 degree decrease of ML-NL angle (from 34.4 to 33.4°) but no significant change in NL-NSL angle. Linear values showed significant intrusion of the second molars (UM2-NL from 21.8 to 20 mm, UM2-NSL from 66.4 to 64.3 mm and LM2-ML from 30.9 to 26 mm). Slight intrusion of the first molars also occurred.

**CONCLUSION:** Miniscrews anchorage is an efficient tool for molar intrusion in the treatment of a skeletal open bite.

## CP 016

Thursday, June 17 - at 15:15

### SOFT-TISSUE LASERS – A NEW AID IN ORTHODONTICS

**R Dimovska**<sup>1</sup>, J Stojanovski<sup>1</sup>, I Kirovski<sup>2</sup>, <sup>1</sup>Private Dental Practice, Kruna MS and <sup>2</sup>Military Hospital, Skopje, Former Yugoslav Republic of Macedonia

Soft tissue lasers have numerous applications in orthodontics, including gingivectomy, fraenctomy, operculectomy, papilla flattening, uncovering temporary anchorage devices, ablation of aphthous ulcerations, exposure of impacted teeth, and even tooth whitening. As an adjunctive procedure, laser surgery has been used to enhance a patient's smile and improve treatment efficacy. Before incorporating soft tissue lasers into clinical practice, the clinician must fully understand the basic science, safety protocol, and associated risks.

Innovative technologies such as the diode laser have provided considerable benefit to dental patients and professionals. Facilitating efficient cutting of tissue and subsequent coagulation, the soft tissue laser enhances tissue healing and can reduce post-surgical complications. Due to the conservative nature of treatment accomplished with the laser, this technology is useful in orthodontic procedures.

The diode laser is utilized in both aesthetic enhancement of the smile, and treatment management of soft tissue issues that impede efficient orthodontic treatment. Whether orthodontists are correcting excessive gingival display or aiding crown lengthening, laser technology is increasing the level of care to patients. Not only are lasers making a practice more effective and efficient, but they are also providing a more comfortable patient experience.

## CP 017

Thursday, June 17 - at 15:15

### ORTHODONTIC-PROSTHODONTIC TREATMENT OF SEVERE SKELETAL CLASS II PROTRUSION

**S Rexhepi**<sup>1</sup>, A Ismani<sup>2</sup>, A Rexhepi<sup>3</sup>, B Ahmeti<sup>3</sup>, L Mustafa<sup>4</sup>,  
<sup>1</sup>Private practice Struga, Clinics of <sup>2</sup>Maxillofacial Surgery and <sup>3</sup>Orthodontics, Skopje and <sup>4</sup>Private practice, Gostivar, Former Yugoslav Republic of Macedonia

**AIM:** To show the orthodontic-prosthodontic treatment of a subject with a severe skeletal Class II malocclusion

**SUBJECT AND METHOD:** A 14-year 1 one month old boy with severe upper anterior protrusion, an overjet of 11 mm, slight anterior crowding and hypodontia of two lower first premolars. Posterior key relationship seemed to be a Class II. Clockwise mandibular rotation was seen on the lateral cephalogram. Because of the severe skeletal measurements, orthognathic treatment was recom-

mended. As the patient refused surgery orthodontic treatment was started using the straightwire technique. Both maxillary first premolars were extracted and the incisors retracted. Active treatment was completed in 25 months. At the end of therapy a fixed denture was inserted in the lower arch between the first premolar.

#### CP 018

Thursday, June 17 - at 15:15

##### THE VALUE OF BIOLOGICAL AGE FOR DETERMINING DENTOFACIAL ORTHOPAEDIC TREATMENT OPTIONS

**E Teodorescu**<sup>1</sup>, E Galan<sup>2</sup>, V-M Milicescu<sup>1</sup>, S Milicescu Jr.<sup>3</sup>, A Bencze<sup>4</sup>, Departments of <sup>1</sup>Orthodontics, <sup>3</sup>Fixed Prosthodontics and <sup>4</sup>Oral Rehabilitation, 'Carol Davila' University of Medicine and Pharmacy and <sup>2</sup>Private Practice, Bucharest, Romania

**AIM:** To re-evaluate the agreement between chronological and biological age.

**SUBJECTS AND METHOD:** Fifteen growing children in the mixed dentition period. Each patient presented a type of skeletal anomaly and a difference between chronological and biological age. For biological age, the synergy or lack of concordance between dental age (determined clinically and radiographically), skeletal maturity (hand-wrist radiographs) and endocrinological age were investigated.

**RESULTS:** All patients exhibited a biological precocity correlated with chronological age that seemed to be related to environmental factors, and in particular dietary factors.

**CONCLUSION:** In subjects with skeletal anomalies, the biological resources of the patient should be investigated in order to manage the remaining potential growth; the hand-wrist radiograph is a valuable tool, used both at the beginning, and during the treatment.

#### CP 019

Thursday, June 17 - at 15:15

##### OPEN BITE CORRECTION WITH MINISCREWS

**A Sapač**<sup>1</sup>, M Drevenšek<sup>2</sup>, M Rejc Novak<sup>3</sup>, <sup>1</sup>Orthas - Private Practice, Murska Sobota, <sup>2</sup>Department of Orthodontics, Medical Faculty, University of Ljubljana and <sup>3</sup>Zavod Ort-Line, Ljubljana, Slovenia

**AIM:** To show the treatment of a subject with an anterior open bite (AOB) using miniscrew anchorage.

**SUBJECT AND METHOD:** A 23-year old male patient with an AOB. He had a tongue thrust habit, disturbed chewing and disturbed speech. Intraoral examination showed a Class I canine and molar relationship on both sides with a 3 mm AOB, a crossbite of the premolars and first molars, and dental erosion. The teeth were sensitive to heat and chemical stimulus. Absolute intrusion of the posterior teeth was planned to autorotate the mandible in a counterclockwise direction, close the open bite, reduce anterior face height and improve facial profile without the need for surgical intervention. The teeth were protected with composite, the upper first molars were splinted with a transpalatal bar placed 4 mm away from the palatal vault and an intrusive force was then applied to the miniscrews (Orlus, Ortholution, Korea) on the buccal side.

**RESULT:** The 3 mm AOB was corrected after 7 months off intrusion. Treatment resulted in a normal overbite (2 mm) and overjet (2.5 mm), a Class I canine and molar relationship, as well as a consonant midline. Cephalometric analysis showed a reduced interbasal angle, ANB and face height.

**CONCLUSION:** Intrusion of the posterior teeth is a good treatment for patients with an AOB.

#### CP 020

Thursday, June 17 - at 15:15

##### EN MASSE RETRACTION WITH ORTHODONTIC MINI-IMPLANTS

**Z Milosavljević**, T Vulicević, P Nikolić, S Colić, M Andrić, School of Dentistry, University of Belgrade, Serbia

**AIM:** To present the use of orthodontic mini-implants in *en masse* retrusion of canines and incisors in Class II division 1 patients.

**SUBJECTS AND METHOD:** Adult patients, with no growth potential and with a severe Class II division 1 malocclusion (SNA angle >82°, ANB angle >6°, I/SpP angle < 65°, overjet more than 6 mm). Decisions concerning extraction of the upper first premolars, implantation of miniscrews and retraction of the anterior teeth were made after analysis of study models and radiographs. When contact was achieved between the upper and lower incisors, the implants were removed, and any persisting extraction spaces, were closed.

**CONCLUSION:** Orthodontic mini-implants are effective as absolute anchorage because there is no mesial movement of the posterior teeth allowing the extraction space to be used only for retraction of the anterior teeth.

#### CP 021

Thursday, June 17 - at 15:15

##### AN UPRIGHTING AUXILIARY FOR DEEPLY IMPACTED MOLARS

**M Greco**, A Maselli. A Giacotti, University of Rome Tor Vergata Fatebenefratelli Hospital, Italy

**AIM:** Traditional orthodontic treatment of impacted second molars is challenging. The aim of this presentation is to illustrate the use of a new auxiliary device for uprighting impacted second molars.

**SUBJECTS AND METHOD:** Twelve patients presenting (mono or bilateral) second molar impaction treated by means of tooth uprighting following non-conventional mechanics based on skeletal anchorage. The auxiliary device comprised a piece of mesh (Leone<sup>®</sup>, a NiTi coil spring exerting force of 150 g and a miniscrew (SIA Excalibur<sup>®</sup>). The piece of mesh was cut and ligated to the NiTi coil spring, then a miniscrew (length: 10 mm, diameter: 1.8 mm) was inserted in the retromolar area. After surgical exposure of the impacted molar, the G-Molar-Up was bonded to the occlusal-distal surface and immediately tied to the miniscrew.

**RESULTS:** All patients were successfully treated and the second molars were uprighted.

**CONCLUSION:** The G-Molar-Up appears to be an effective option due to the simple bonding procedure, favourable biomechanics, and the continuous force exerted.

#### CP 022

Thursday, June 17 - at 16:00

##### CONE BEAM COMPUTED TOMOGRAPHY AND PYKNODYSTOSIS – A CASE REPORT

**L Laffranchi**<sup>1</sup>, P Fontana<sup>2</sup>, S Bonadeo<sup>2</sup>, F Zotti<sup>2</sup>, D Dalessandri<sup>1</sup>, Departments of Orthodontics, <sup>1</sup>University of Turin and <sup>2</sup>University of Brescia, Italy

**AIM:** Pyknodysostosis is an autosomal-recessive disorder of osteoclast dysfunction causing osteosclerosis, due to deficiency of cathepsin K. It is characterized by short stature, a large skull and short terminal phalanges. Oral and maxillofacial features include an obtuse mandibular gonial angle, hypoplastic paranasal sinuses, primary and permanent teeth with frequent crowding in the lower or upper dental arch. During childhood radiographic images of osteosclerosis, often show open fontanelles and sutures and the presence of Wormian bones, pathological fractures and obstructive sleep apnoea become apparent. The aim of this study was to analyse the features of a patient with pyknodysostosis using computed tomography.

**SUBJECT AND METHOD:** A 9 year 11 month old male with pyknodysostosis caused by an heterozygote mutation in the gene p.R241X. For surgical reasons, a cone beam computed tomograph was required which was obtained with a NewTom 3G.

**RESULTS:** There was a relationship between the dental elements themselves and surrounding anatomical structures.

**CONCLUSION:** Cone beam computed tomography and three-dimensional reconstructions of patients with maxillo-facial dysmorphic disorder allow the best diagnosis and prognosis and accurate planning of surgical treatment. It is possible to obtain images of the whole skull that allow complete orthodontic diagnosis.

**CP 023**

Thursday, June 17 - at 16:00

**ORTHODONTIC-SURGICAL CORRECTION OF A CLASS II DIVISION 1 PATIENT WITH FACIAL NERVE WEAKNESS****I Drerup**, K Heckmann, J Lisson, Department of Orthodontics, University of Saarland, Homburg/Saar; Germany

**AIM:** To present the combined orthodontic-surgical treatment of a patient with additional aggregating factors. The patient showed an excessive Class II division 1 malocclusion (ANB 10.3°), weakness of the facial nerve on the right side and cerebral paresis. The weakness of the facial nerve and the lathernognathia contributed to an asymmetric facial structure.

**SUBJECT AND METHOD:** The interarch relationship exhibited mandibular retrognathia and laterognathia with limited mouth opening (30 mm). The overjet was 18 mm and the overbite 4 mm. Analysis of the plaster casts showed 9 mm crowding in the upper and 9 mm in the lower arch. The upper incisors presented excessive protrusion (Ils/NA 31.3°). Initial treatment began with the extraction of the first premolars followed by insertion of a transpalatal and lingual arch for anchorage. At the same time, upper and lower preadjusted straightwire appliances were fitted. After several visits the patient underwent mandibular advancement using an Obwegeser/Dal Pont bilateral sagittal split osteotomy.

**RESULTS:** Combined orthodontic-surgical treatment resulted in greatly improved occlusal function and considerable aesthetic correction. Both weakness of the facial nerve and limited mouth opening did not significantly improve. After surgical intervention the facial nerve appeared less affected and the asymmetry was less significant. Overjet and overbite were reduced to 2 mm.

**CONCLUSION:** Given that the patient showed a clearly expressed Class II malocclusion and several other features, combined orthodontic-surgical treatment was the therapy of choice.

**CP 024**

Thursday, June 17 - at 16:00

**MAXILLARY CANINE-FIRST PREMOLAR TRANSPOSITION IN THE PERMANENT DENTITION. CLINICAL CASES****N Vasilakou**, P Synodinos, Department of Orthodontics, University of Athens, Greece

**AIM:** To review orthodontic treatment considerations of three patients presenting with maxillary canine-first premolar transposition in the permanent dentition.

**SUBJECTS AND METHOD:** Three clinical cases selected from the files of an orthodontic practice where the transposed canines and first premolars were orthodontically moved to their normal places in the maxillary arch.

**RESULTS:** Treatment alternatives included the maintenance and alignment of the teeth in their transposed position (with or without orthodontic treatment) and orthodontically facilitated correction of the natural order of the teeth. Early diagnosis of the transposed eruption pathway of the upper canine was the most decisive factor facilitating the timely application of orthodontic treatment towards re-establishing a normal occlusion, thus avoiding any occlusal and restorative compromises.

**CONCLUSION:** Treatment of choice is selected for each case separately after thoroughly considering overall facial and dental characteristics, treatment timing, duration and cost, patient's preference and orthodontist's experience.

**CP 025**

Thursday, June 17 - at 16:00

**CAUSAL RELATION BETWEEN MALOCCLUSION AND ORAL MUSCLE DYSFUNCTION – A MODEL APPROACH****G Antonini<sup>1</sup>**, S Saccomanno<sup>1</sup>, L D'Alatri<sup>2</sup>, D Scarpa<sup>1</sup>, R Deli<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Otorhinolaryngology Catholic University of the Sacred Heart, Rome, Italy

**AIM:** Deleterious habits alter functions that can cause anomalies in orofacial morphology. To correct these problems, orthodontic treatment can be supported by myofunctional therapy to recover

the normal function of the oral muscles. The aim of this study was the creation of a diagnostic and therapeutic model, with an interdisciplinary approach, that allows avoidance of any relapse that can occur after orthodontic treatment.

**SUBJECTS AND METHOD:** Twenty-three patients with atypical swallowing together with, in some subjects, thumb sucking and oral breathing. A clinical and instrumental case history was drawn up for each case, in order to give a correct orthodontic and functional diagnosis. Initially, anomalies, which if left untreated, could compromise the success of therapy (maxillary contraction, oral breathing, short lingual fraenum) were corrected. A different therapeutic approach was then applied on the basis of skeletal and dental features, using different therapies both for duration and orthodontic appliances.

**RESULTS:** Both from a diagnostic and therapeutic point of view, important results have been achieved especially through muscle analysis with a dynamometer and surface electromyography.

**CONCLUSION:** Orthodontic therapy, in the presence of deleterious habits, is insufficient to resolve the orthodontic issues. It is necessary to combine orthodontic treatment with myofunctional therapy.

**CP 026**

Thursday, June 17 - at 16:00

**CONSIDERATION OF CANINE TEETH: DIAGNOSIS AND TREATMENT****S Saccomanno**, A Oliverio, C Notaristefano, G Antonini, R Deli, Department of Orthodontics Catholic University of the Sacred Heart, Rome, Italy

**AIM:** Tooth inclusion is the non-eruption of a dental element in the mouth when physiological time limit for the element to erupt has passed. An element is then 'inclusive' when it is not erupted and has no eruptive potential. Predisposing factors to consider are hardness of the palatal bone and mucoperiosteal tissue, a longer course to cover for the canine compared with other teeth, early loss of the primary canine or its late root resorption after loss, and lack of space. Diagnosis is made through anamnesis and clinical and radiographic examination. Therapy depends on the type of inclusion.

**MATERIALS AND METHOD:** Ten clinical cases have been studied. **RESULTS:** Treatment gave excellent results with eruption of all teeth. The correct therapeutic programme for recovering the included canines must consider problems such as ankylosis and unfavourable positions before any treatment.

**CONCLUSION:** Correct planning and therapeutics are basic requirements for the alignment of the tooth when an included canine is diagnosed.

**CP 027**

Thursday, June 17 - at 16:00

**MULTIDISCIPLINARY TREATMENT OF AN ADULT WITH MAXILLARY RETROGNATHISM AND OLIGODONTIA****D A Papamanou<sup>1</sup>**, O N Haralabakis<sup>2</sup>, A K Tsourakis<sup>3</sup>, <sup>1</sup>Department of Orthodontics and <sup>2</sup>Dental School, University of Athens and <sup>3</sup>Private Practice, Athens, Greece

**AIM:** Oligodontia is defined as the congenital absence of six or more permanent teeth, excluding the third molar. The aetiology of congenital absence of teeth is believed to be due to heredity or developmental anomalies. The aim of this presentation is to show the treatment of a patient with a skeletal anteroposterior discrepancy and multiple congenitally missing teeth through orthodontic treatment, and emphasize the importance of collaboration with other dental disciplines to obtain optimal aesthetic and functional results.

**SUBJECT AND METHOD:** An 18-year-old male with 22 congenitally missing permanent teeth. The only permanent teeth present were all first molars, the upper right canine, the upper central incisors, the lower right canine and the lower first premolars. Twelve primary teeth were also present. The patient had a severe maxil-

lary anteroposterior discrepancy (ANB:  $-8^\circ$ , SNA:  $73.1^\circ$ , Lande's angle  $78.8^\circ$ ) and maxillary vertical deficiency with decreased maxillary incisor exposure. Angle classification was Class III on the right and Class I on the left.

**RESULTS:** All present teeth were bonded and banded. After 36 months of active treatment the remaining teeth were correctly positioned for implants to be placed. All primary teeth were extracted, maxillary surgical advancement was performed and 12 implants were placed (6 in the maxilla and 6 in the mandible) in order to support temporary restorations. After completion of orthodontic treatment (6 months later), permanent prosthodontic restorations were placed. One month before the end of orthodontic treatment the upper right canine became secondarily ankylosed.

**CONCLUSION:** A combination of orthodontic, orthognathic and prosthodontic treatment can contribute to a satisfying outcome in such a complicated and demanding case.

## CP 028

Thursday, June 17 - at 16:00

### SPEEDING UP ORTHODONTIC TREATMENT

L Barreto, D Spary, Birmingham Dental Hospital, England

If orthodontic treatment can be completed in a shorter time, whilst still achieving the same standard of finish, it would benefit the patient and the orthodontist. A disciplined approach and some improved mechanics might deliver this desirable outcome. Examples of methods of shortening treatment time will be given. The following mechanics are considered:

1. Logical bracket selection and accurate bracket placement: e.g. using the MBT gauge to avoid excess space in the upper lateral incisor region
2. Reduction of friction: using double tubes to allow the lower first molars to slide forwards
3. Multi-tasking: using stabilising arches to allow the use of Class II elastics
4. Use of elastics: the modified Parkhouse arch for rapid space opening and the use of Swiss twists for closing space in the canine regions
5. Use of auxiliaries such as the single tooth torquing auxiliary.

## CP 029

Thursday, June 17 - at 16:00

### EFFECTIVENESS OF FRAENECTOMY ON TONGUE FUNCTION AND POSTURE – AN ULTRASOUND EVALUATION

R Pavlovič<sup>1</sup>, J Volk<sup>1</sup>, M Mušič<sup>2</sup>, B Gašpirč<sup>3</sup>, M Ovsenič<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>3</sup>Parodontology, University of Ljubljana and <sup>2</sup>Institute of Oncology Ljubljana, Slovenia

**AIM:** The length of lingual fraenum plays an important role in tongue function and posture and in the functional diagnosis of orthodontic patients. It is difficult to clinically evaluate tongue function and posture in patients with a short lingual fraenum. The aim of this presentation is to show the effectiveness of laser fraenulotomy on tongue function and posture.

**SUBJECT AND METHOD:** An 8-year-old boy with speech problems. During clinical examination a short lingual fraenum and tongue posture on the mouth floor were diagnosed. Assessment of tongue function and posture was evaluated using ultrasonography before and one year after fraenectomy. The difference between infantile and somatic swallowing pattern and tongue posture was evaluated using ultrasonography (ultrasound system Voluson 730 Expert and a 3D convex transducer RAB 2-5 MHz). An Erbium laser was used for the surgical fraenectomy.

**RESULTS:** The irregular swallowing pattern and tongue posture did not change after surgical intervention.

**CONCLUSION:** Ultrasonography is a non-invasive, reliable, and objective diagnostic tool in the assessment of tongue posture and function and may be useful in evaluation of different treatment modalities of functional irregularities. A fraenectomy was not successful in improving tongue function and posture. How-

ever, the method should further be evaluated on a larger sample in order to confirm the hypothesis of epigenetic regulation (fraenulotomy) of tongue function and posture.

## CP 030

Thursday, June 17 - at 16:00

### MICROSCOPIC ANALYSIS OF THE ROOT SURFACE OF TEETH

CT Preoteasa, E Ionescu, E Preoteasa, A C Didilescu, A M Bencze, Faculty of Dental Medicine, University 'Carol Davila', Bucharest, Romania

**AM:** External root resorption (ERR) is an irreversible phenomenon associated with orthodontic treatment, which can have medical, ethical and legal implications. The aim of this comparative in vitro study was to assess dental morphology at the apical root surface between orthodontically treated teeth.

**MATERIALS AND METHOD:** Nine extracted teeth, four with a history of fixed orthodontic treatment. The five teeth without a history of orthodontic treatment included one tooth, two recently erupted teeth and two teeth with severe dental wear (correlated with excessive occlusal loading). Microscopic examination was undertaken with a Zeiss Stemi 2000C (Carl Zeiss GmbH) to determine whether changes in surface morphology and the apical contour (that can be correlated with a lack of substance) were present.

**RESULTS:** The root surface of the recently erupted teeth had a smooth appearance. Teeth with dental wear had a generally uniform surface appearance, but with a greater roughness. All teeth with a history of orthodontic treatment presented a very different root surface morphology. Lacunae were present, in different numbers, size and arrangement. These aspects were seen as being characteristic of ERR.

**CONCLUSION:** Microscopic analysis is a useful method for identifying and analysing ERR, being able to provide important details for a better understanding of this phenomenon.

## CP 031

Thursday, June 17 - at 16:00

### MODIFICATION OF DENTAL HARD TISSUES DURING ORTHODONTIC TREATMENT – A MICROSCOPIC ANALYSIS

CT Preoteasa, E Ionescu, E Preoteasa, A C Didilescu, A D Ganea, Faculty of Dental Medicine, University 'Carol Davila', Bucharest, Romania

**AIM:** Enamel alterations (discolouration, decay) and root resorption are two undesirable side-effects of orthodontic treatment. The aim of this research was to analyse the dental hard tissues at the coronal and apical root level in order to identify morphological changes, possibly related to fixed orthodontic treatment.

**MATERIALS AND METHOD:** Four extracted maxillary premolars, all with a history of fixed orthodontic treatment (for a period between 11 and 13 months). Orthodontic treatment was carried out with a conventional vestibular appliance and self-curing composite for bracket bonding. For all teeth a microscopic examination was carried out with a Zeiss Stemi 2000C (Carl Zeiss GmbH) to determine whether there were any changes in surface morphology at the coronal and apical levels.

**RESULTS:** On the vestibular surfaces of the teeth, the area where the bracket was bonded was clearly delineated. The tooth surface under the bracket presented a normal aspect. In contrast, the adjacent area showed modifications regarding colour and roughness. Regarding the apical root surfaces, lacunae were identified in all teeth. These were different in number, size and arrangement. These aspects were seen as being characteristic of external root resorption associated with orthodontic treatment.

**CONCLUSION:** The use of an orthodontic appliance can be correlated with changes in the morphology of the dental hard tissues. Further investigations with a larger sample size would be useful to analyse both the incidence and severity of these alterations.

**CP 032**

Thursday, June 17 - at 16:00

**ORTHODONTIC TREATMENT OF LOWER SECOND MOLAR IMPACTION WITH SKELETAL ANCHORAGE – A CASE REPORT**

**A M Enache**<sup>1</sup>, I Nicolescu<sup>2</sup>, <sup>1</sup>Department of Orthodontics and Dentofacial Orthopaedics, Faculty of Dental Medicine, University of Medicine and Pharmacy 'C. Davila', Bucharest and <sup>2</sup>St. Mary Hospital, Private Practice, Romania

**AIM:** To present the treatment of lower second molar impaction using miniplates for maximum anchorage.

**SUBJECT AND METHOD:** A 14 year 7 month-old female patient with a Class II dental and skeletal malocclusion, high angle, second lower molar impaction and upper and lower anterior crowding. The parents refused orthognathic surgery and orthodontic treatment required extractions. In order to decide which teeth to extract, the lower second molars were aligned using miniplates for anchorage placed in the vertical ramus of the mandible.

**CONCLUSION:** Skeletal anchorage is a useful method for orthodontic treatment

**CP 033**

Thursday, June 17 - at 16:00

**SECONDARY CLEFT PALATE DENTAL PULP STEM CELL GRAFTING IN VITRO. A CONE BEAM COMPUTED TOMOGRAPHIC ASSESSMENT OF SCAFFOLD REABSORPTION**

**D Dalessandri**<sup>1</sup>, L Laffranchi<sup>1</sup>, M G Piacino<sup>1</sup>, C Paganelli<sup>2</sup>, P Bracco<sup>1</sup>, Departments of Orthodontics, <sup>1</sup>University of Turin and <sup>2</sup>University of Brescia, Italy

**AIM:** In complete cleft lip and palate patients, in order to obtain palatal closure, dental pulp stem cells (DPSCs) obtained from primary teeth can be used during secondary grafting. It is important to monitor scaffold reabsorption and bone formation during the healing process. Spongostan® has been suggested as scaffold material in order to convey DPSCs into the bone defect. The aim of this *in vitro* study was to investigate the possibility of evaluating, using cone beam computed tomography (CBCT), the presence of this scaffold material under different conditions dry, wetted with a saline solution and with blood.

**MATERIALS AND METHOD:** A test phantom was constructed utilizing a dry skull inserted into a plastic sheath lined with foam rubber and covered by wet sheet. The scaffold material was wetted with a syringe by inserting the needle through the plastic package and it into the extraction socket of tooth 3.6 and into the left nasal cavity. Several CBCT examinations were utilized using the Newton 3G® scanner (FOV: 6 inches).

**RESULTS:** It was possible to evaluate all the phantom bone structures, to identify the tested material and to determine differences between the blood and the saline solution. It was not possible to localize the presence of the dry material.

**CONCLUSION:** CBCT results in a lower X-ray dose compared with traditional multislice CT. This is important in young patients with pathologies, such as those with a cleft lip and palate (CLP), who require several radiographic examinations during treatment. The *in vitro* results suggests that CBCT can be useful in the assessment of scaffold reabsorption and bone formation during the bone healing process after secondary grafting in complete CLP patients.

**CP 034**

Thursday, June 17 - at 16:45

**AN ORTHODONTIC APPROACH IN TREATING BICONDYLAR FRACTURES**

**E Çerekja**<sup>1</sup>, A-A Demirkaya<sup>1</sup>, K Uğurlu<sup>2</sup>, <sup>1</sup>Marmara University and <sup>2</sup>Division of Plastic and Reconstructive Surgery, Sisli Etfal Hospital, Istanbul, Turkey

**AIM:** To show the orthodontic treatment approach in a patient with bicondylar fractures.

**SUBJECT AND METHOD:** A 48-year-old male with bicondylar fractures. The patient had difficulty in closing the mouth and pain. An anterior open bite and disocclusion was present. The

shortening of the two ramii was found to be 3.5 mm from the posteroanterior radiograph. To restore ramus height reduced by the contraction of the muscles and to distract the segments to the normal height, a pivot splint, 3.5 mm thick on the molar area in the ideal occlusion, was used with intermaxillary fixation. Stainless steel 0.017 X 0.025 inch wires were passively bent on the brackets which were indirectly bonded to the dental arches by transfer trays. Intermaxillary fixation was used for 6 weeks.

**RESULTS:** After treatment, no asymmetry was present and the patient could perform all the movements without pain.

**CONCLUSION:** Intermaxillary fixation with pivot splints are a good treatment option to surgery in subjects with condylar fractures.



### Class III Malocclusion

#### CP 035

Wednesday, June 16 - at 15:15

##### ORTHODONTIC TREATMENT OF A CLASS II MALOCCLUSION IN AN ADULT SUBJECT

**M Stojanović**, <sup>1</sup>Private practice, Paraćin, Serbia

**AIM:** Correction of a Class II division 2 malocclusion and unilateral reverse-bite of the permanent canines in order to relieve pain in the patient's temporomandibular joints (TMJ), and achieve correct occlusion and facial symmetry.

**SUBJECT AND METHOD:** A 32-year old female patient with a Class II division 2 malocclusion (characterised by three-dimensional disharmony, particularly in the vertical dimension), a reverse-bite of the right permanent canines and TMJ pain. Disocclusion of the permanent canines (keys of occlusion) was used to bring them from a reverse into the normal position. This was accompanied by correction of the pronounced curve of Spee, which was performed by levelling and application of mild continuous forces, matching the dental arches and control of the vertical positions of the maxillary and mandibular incisors. Fixed appliance therapy was directed towards functional adjustment of the joint to the new situation, thus relieving the patient of pain.

**RESULT:** Reversal of the Class II division 2 malocclusion into a normal occlusion, and correction of the reverse-bite of the permanent canines resulted in the elimination of TMJ pain and correct facial symmetry.

**CONCLUSION:** The above condition should be recognized much earlier in life, sometimes even in early childhood.

#### CP 036

Thursday, June 17 - at 10:45

##### THREE-DIMENSIONAL ASSESSMENT OF CLASS III TREATMENT – A CASE REPORT

**S Lah Kravanja**<sup>1</sup>, J Primožič<sup>2</sup>, M Ovesnik<sup>1</sup>, <sup>1</sup>Department of Orthodontics, University, Medical Centre and <sup>2</sup>Orthos, Ljubljana, Slovenia

**AIM:** The morphological characteristics and facial appearance of a Class III malocclusion subject can develop because of a functional dynamic imbalance of soft tissues. The aim of this research was to evaluate the orthodontic treatment of a 6 year-old girl with a Class III malocclusion and facial appearance with asymmetry and chin deviation to the right.

**MATERIALS AND METHOD:** Clinical examination, facial and intraoral photographs, dental casts, dental pantomogram, lateral cephalogram and a three-dimensional (3D) evaluation of the girl's face were taken at baseline (T0). The girl was treated with a modified Fränkel appliance type III. After 24 months (T2) the facial appearance was assessed using a 3D laser scanning technique in order to evaluate treatment success. The scan of the girl was compared with a facial template of an average face, constructed using facial scans of 60 children. Differences in facial morphology were assessed quantitatively and qualitatively. Palatal volume was measured at T0 and T2.

**RESULTS:** At T0 3D analysis showed a retrognathic midface and a prognathic chin with a slight deviation to the right. The girl had a more prominent chin and less prominent midface area compared with the average face. After 24 months of treatment, 3D soft tis-

sue analysis showed a less retrognathic midface and correction of the chin deviation. A correct lip seal, nasal breathing, and tongue function and posture were established and a good occlusion was obtained.

**CONCLUSION:** Early treatment of Class III malocclusions with a modified Fränkel appliance was able to improve facial appearance as well as the functional and morphological irregularities together with space conditions of the oral cavity.

#### CP 037

Thursday, June 17 - at 10:45

##### EARLY TREATMENT OF A CLASS III MALOCCLUSION WITH A DELAIRE TYPE FACE MASK – A CASE REPORT

**S Lah Kravanja**<sup>1</sup>, M Drevšek<sup>2</sup> <sup>1</sup>Departments of Orthodontics, <sup>1</sup>University Medical Centre and <sup>2</sup>University of Ljubljana, Faculty of Medicine, Slovenia

**AIM:** Children with Class III malocclusions are difficult to treat due to maxillary deficiency and excessive growth of the mandible. For children with an antero-posterior and vertical maxillary deficiency a reverse-pull headgear-facemask is the preferred treatment. The aim of this presentation is to show treatment with a Delaire type facemask for maxillary protraction following palatal expansion in a growing child with a Class III malocclusion and functional disorders.

**SUBJECT AND METHOD:** A 7-year-old girl in the mixed dentition. Clinical examination showed a bilateral crossbite, negative overjet and overbite, prominent chin, and morphological and jaw discrepancies. Facial and intraoral photographs, dental casts, a dental pantomogram and a lateral cephalogram were taken at baseline. The girl was treated with an acrylic plate for palatal expansion and upper incisor correction. Treatment continued with a quadhelix and rigid arch for attachment to the facemask. The hooks were located in the upper incisor and canine area, gingivally placed and above the occlusal plane, which places the force vector near the centre of resistance of the maxilla. After eruption of the upper incisors, straightwire brackets were bonded and Kobayashi ligatures for the facemask elastics. The direction of pull for protraction was more forward than down. The forehead and chin pads enabled the maxilla to be moved down as well as forward and the mandible to rotate downward and backward. A force of approximately 400 g per side was applied for up to 14 hours a day.

**RESULTS:** The retrognathic maxilla, prognathic mandible, negative overjet and overbite were corrected after 18 months of treatment. There was a general improvement of the facial profile and dental appearance, and the anterior teeth were aligned.

**CONCLUSION:** Whether orthognathic surgery will be required will be determined by mandibular growth during and after adolescence.

#### CP 038

Thursday, June 17 - at 17:30

##### NON-SURGICAL TREATMENT OF A JUVENILE PATIENT WITH PYCNODYSTOSIS AND A CLASS III MALOCCLUSION

**J Barekzai**, M Kotthaus, B Kahl-Nieke, Department of Orthodontics, Centre for Dental and Oral Medicine, University Medical Centre Hamburg-Eppendorf, Germany

**AIM:** Pycnodysostosis is a rare autosomal recessive inherited disease belonging to the group of craniotubular bone dysplasias.

Mandibular osteomyelitis is common due to impaired vascularity of the sclerotic bone. Successful surgical correction has been mentioned in case reports, but the quality and vascularity of the bone limits the possibilities for surgical correction, which include infection risks for patients with pycnodysostosis. This presentation describes a non-surgical orthodontic treatment option.

**SUBJECT AND METHOD:** A 14-year-old girl with pycnodysostosis with a Class III malocclusion and vertical growth pattern. After extraction of two lower first premolars, an expansion plate was used to expand the maxillary dentoalveolar arch, followed by a full-bracket appliance, combined with Class III elastics to correct the anterior crossbite and eliminate the mandibular displacement. After debanding, an active plate was used to finish treatment. All appliances were removed 34 months after initiating treatment.

**RESULTS:** A successful outcome was achieved by correcting the crossbite and eliminating the mandibular displacement. The final occlusion remained stable. Expected problems with tooth movement because of sclerotic complications in the bone did not occur.

**CONCLUSION:** Non-surgical orthodontic treatment permitted acceptable results. Surgical treatment with risks of infection and inadequate bone adaptation was not necessary.

### CP 039

Thursday, June 17 - at 17:30

#### CALIBRATED DOUBLE-SPLIT-CAST MÜNSTERANER MODEL-OPERATION SYSTEM

**T Ziebur**, T Stamm, U Ehmer, Westfalian Wilhelms University, Münster, Germany

Orthognathic surgery is elective and therefore the whole concept of integrated orthodontic and surgical correction of dentofacial deformities requires a predictable quality procedure. The calibrated double-splint Münsteraner model surgery provides a standardized interactive solution to model surgery and splint construction. The articulator-based hardware system is capable of performing three-dimensional simulations for all types of surgical treatment, including combined distraction osteogenesis. It also allows for fabrication of differentiated intermediate and final splints. Case reports featuring Class III malocclusion subjects demonstrate the system's clinical concept.

### CP 040

Friday, June 18 - at 15:15

#### USE OF TEMPORARY ANCHORAGE DEVICES TO CORRECT SEVERE CLASS III MALOCCLUSIONS, WITHOUT ORTHOGNATHIC SURGERY

**D Mahony**, Private Specialist Orthodontic practice, Australia and Brazil

**AIM:** Many Class III patients are unable to afford orthognathic surgery, or are happy with their facial appearance, but wish to improve their dental alignment to prevent further wear to the anterior teeth. The aim of this study was a cephalometric review of 20 consecutive cases, via a clinical audit, successfully treated via upper arch development and full fixed appliances, with the use of temporary anchorage devices (TADs) in the lower jaw

**MATERIALS AND METHOD:** The patients had initially signed a surgical waiver form understanding that their ideal treatment would involve a combination of orthodontics and orthognathic surgery, but they were happy to accept a compromised result via dental compensation of their Class III jaw pattern. Some had the lower third molars extracted to create space for *en masse* retraction of the entire mandibular dentition. All had 12 mm TADs placed vertically in the superior oblique ridge of the mandible, in the region between the lower first and second molars. A 9 mm NiTi closing coil spring was used from the TADs to posted hooks on a lower 0.019 X 0.025 stainless steel archwire, in a 0.022 edgewise bracket slot.

**RESULTS:** Cephalometric evaluation showed a forward rotation of the occlusal plane with minimal retroclination of the lower incisors, and an ideal Class I buccal segment relationship. All patients were happy with the changes to their facial profile.

**CONCLUSION:** This non-surgical technique should be offered to Class III patients seeking a non-surgical approach for the correction of their negative overjet.

### CP 041

Friday, June 18 - at 15:15

#### ORTHOGNATHIC TREATMENT OF A SEVERE CLASS III MALOCCLUSION: A CASE REPORT

**E Baharoğlu**<sup>1</sup>, İ Kocadereli<sup>1</sup>, G Tunçbilek<sup>2</sup>, E Mavili<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Plastic and Reconstructive Surgery, Hacettepe University, Ankara, Turkey

**AIM:** To present the orthodontic and orthognathic treatment of a patient with a Class III malocclusion with severe mandibular prognathism.

**SUBJECT AND METHOD:** A 24-year-old female diagnosed with skeletal discrepancy, with the chief complaint of poor aesthetics. Clinically she had severe mandibular prognathism and moderate maxillary retrognathism. In addition she had a facial asymmetry with the chin deviated towards the right side. The canine and molar relationship was Class III with a 5 mm reverse overjet. Diagnostic records included cephalometric, postero-anterior (PA) and panoramic radiographs, intra- and extraoral photographs and dental casts. Cephalometric analysis showed skeletal a Class III relationship (ANB: -7°) with a mandibular excess (SNB: 84.0°) and maxillary deficiency (SNA: 77°). The mandibular incisors were lingually inclined (L1-MP: 72°). PA analysis showed facial asymmetry with a 4 mm mandibular deviation to the right side. Roth 0.018 inch slot type brackets were used. The patient was ready for bimaxillary orthognathic surgery in 16 months. The maxilla was advanced 3 mm sagittally with a Le Fort I osteotomy and the mandible was set back 5 mm, allowing autorotation with a modified bilateral sagittal split osteotomy.

**RESULTS:** A Class I occlusion with normal overjet and overbite was achieved in 20 months. Post-treatment cephalometric evaluation showed a skeletal Class I jaw base relationship (ANB: 0°) with the maxillary deficiency corrected (SNA: 87°).

**CONCLUSION:** Post-treatment the facial appearance showed a dramatic change. The protrusive chin and concave profile were improved by orthodontic/orthognathic treatment, resulting in a straight profile.

### CP 042

Friday, June 18 - at 15:15

#### DENTO-SKELETAL CHANGES IN A CLASS III MALOCCLUSION SUBJECT WITH AN OPEN BITE – A CASE REPORT

**A Fekonja**, Orthodontic Department, Health Centre, Maribor, Slovenia

**AIM:** To describe and illustrate the therapeutic possibilities for correction of a Class III open bite malocclusion using a fixed straightwire appliance and a facemask.

**SUBJECT AND METHOD:** A patient with skeletal and dentoalveolar components of a Class III malocclusion with an open bite. Clinical examination revealed lip incompetence, mouth breathing, and abnormal tongue posture and function. Dentoalveolar examination showed protrusion of the upper and lower incisors. Skeletal examination showed a downward and backward rotated mandible, maxillary retrusion and increased gonial and interbasal angles, Wits amount, and lower face height. The skeletal and dentoalveolar component of the Class III open bite malocclusion were evaluated by analyzing orofacial function, dental casts and lateral cephalometric radiographs before and after treatment.

**RESULTS:** Angular as well as linear horizontal and vertical changes were found after a treatment period of 26 months.

**CONCLUSION:** A fixed appliance in association with extraoral force (facemask) is an elective mechanism for correction of this malocclusion. The treatment changes that occurred were mainly dentoalveolar.

**CP 043**

Friday, June 18 - at 15:15

**ORTHODONTIC TREATMENT OF A PATIENT WITH FRONTONASAL DYSPLASIA****H Kan**, T Taner, Department of Orthodontics, Faculty of Dentistry, Hacettepe University, Ankara, Turkey

**AIM:** Frontonasal dysplasia is a rare congenital syndrome characterized by orbital hypertelorbitism, a high frontal hairline, mid-facial hypoplasia, nasal hypoplasia, cleft lip and palate, defect of alae nasi, and bifid nose or grooved nasal tip. The aim of this case report is to present the effects of combined rapid palatal expansion (RPE) and facemask treatment on dentofacial structures and upper airway dimensions in a patient with frontonasal dysplasia and maxillary retrusion.

**SUBJECT AND METHOD:** A 9 year 6 month-old Caucasian male with frontonasal dysplasia and a skeletal Class III malocclusion. The patient had undergone surgery for cranial synostosis at 18 months of age. Clinical examination showed that the patient was a mouth breather who snored. A bilateral posterior crossbite, anterior open bite and reverse overjet were observed. Radiographic examination showed that the two mandibular central incisors were congenitally absent. Lateral cephalometric measurements indicated maxillary retrusion and mild mandibular protrusion. Orthodontic treatment started with combined bonded RPE and facemask therapy. Expansion was obtained by turning the screw once a day for one month. When the maxilla was positioned anteriorly, the facemask and bonded RPE were removed and a quad-helix appliance was inserted for retention.

**RESULTS:** At the end of the treatment, mouth breathing and snoring were eliminated. Cephalometric measurements were changed as follows: overjet increased from -3 to +7 mm, convexity value increased from -2 to +5 mm, maxillary depth angle increased from 92 to 94 degrees, SNA angle increased from 79 to 83 degrees, SNB angle reduced from 79 to 77 degrees, and facial depth angle decreased from 93 to 90 degrees. Upper airway dimensions and palatal, middle and inferior pharyngeal spaces significantly increased.

**CP 044**

Friday, June 18 - at 15:15

**REVERSE HEADGEAR WITH A DISTALIZATION SCREW – A CASE REPORT****E K Atamozlu**, Y B Goyenc, Department of Orthodontics, Faculty of Dentistry, Selcuk University, Konya, Turkey

**AIM:** With reverse headgear in patients with a Class I molar relationship, after anterior movement of the maxilla, the relationship becomes Class II. The aim of this research was to eliminate this effect and gain space using a screw through the sagittal plane for the distalization of the first molars.

**SUBJECT AND METHOD:** An 8-year 10-month-old female patient with a Class III malocclusion. Reverse headgear and removable appliance to the maxilla were used for 5 months. A screw was placed in the appliance between the upper molars to provide movement in the distal direction. After correction of the Class III relationship, a chin cap used for mandibular retention.

**CONCLUSION:** With this appliance a dental Class I relationship is avoided and space is gained easily, without the use of fixed appliances. The treatment period was shortened because molar movement posteriorly and maxillary movement anteriorly were achieved at the same time.

**CP 045**

Friday, June 18 - at 15:15

**MAXILLARY PROTRACTION WITH TWO BONE ANCHORS IN A CLASS III PATIENT WITH OLIODONTIA****E Dumu Dundu Jr.**, C Boucher, A Balon-Perin, R Glineur, Department of Orthodontics and Dentofacial Orthopaedics, Free University of Brussels, Erasmus Hospital, Belgium

**AIM:** Facemask therapy is a common way to treat Class III patients in the mixed dentition. The principle is based on traction between

a maxillary appliance and a facemask. The aim of this presentation is to describe the treatment of a patient with a Class III skeletal relationship with agenesis.

**SUBJECT AND METHOD:** An 11-year-old girl with a Class III skeletal relationship and seven missing teeth (excluding the third molars). The maxillary arch was in a complete crossbite relationship with the mandibular arch. The treatment plan was to protract the maxilla anteriorly with a facemask. Because of the lack of posterior maxillary teeth, two bone anchors were placed in the infra-zygomatic crest of the maxilla. Three weeks later, a force of approximately 500 g was applied to both sides.

**RESULT:** A significant anterior maxillary displacement, a normalization of the anterior crossbite and a Class III dental relationship to a Class I dental relationship were achieved.

**CONCLUSION:** Advancement of the maxilla and improvement of the soft tissue profile can be achieved using skeletal anchorage in combination with facemask therapy in mixed dentition patients.

**CP 046**

Friday, June 18 - at 15:15

**UPPER MOLAR ALIGNMENT WITH A REVERSE RAPID MAXILLARY EXPANSION APPLIANCE – A THREE-DIMENSIONAL CAST ANALYSIS\*\*\*****M Stec-Slonicz**, **I Drerup**, J Lisson, Department of Orthodontics, University of Saarland, Homburg/Saar, Germany

**AIM:** Pre-surgical orthodontic treatment of maxillary deficiency aims at congruent dental arches. In many Class III malocclusions the transverse discrepancy between the maxilla and mandible makes this difficult. The aim of this presentation is to demonstrate correction of buccally tipped molars with a reverse rapid maxillary expansion (RME) appliance. To visualize the dental and palatal changes three-dimensional cast analysis was performed.

**SUBJECT AND METHOD:** A 17-year-old female with severe maxillary deficiency combined with a tendency towards mandibular prognathism. Cast analysis revealed protrusion of the upper incisors and significant buccal tipping of the upper first and second molars, resulting from previously inefficient orthodontic treatment. The overjet was negative (-2 mm) with a Class III molar relationship of over one premolar width with a mandibular shift of 4 mm to the left. As the treatment plan included a Le Fort I osteotomy for maxillary advancement, initial tooth alignment was necessary. The greatest problem was buccal tipping of the upper molars. For appropriate pre-surgical arch alignment, horizontal correction of these teeth was necessary. To achieve palatal movement of the molars a reverse RME appliance with an open jack screw (Forestadent, Pforzheim, Germany; maximum expansion 8 mm) was inserted, fixed to the first and second upper molars. The screw was adjusted twice a day (0.5 mm daily), resulting in 0.25 mm gap closure with every turn.

**RESULTS:** After 16 days, the horizontal molar position was corrected and the now closed jackscrew was immobilized with dental cement. Total palatal movement of the first molars of 6 mm and second molars of 7 mm was achieved.

**CONCLUSION:** The reverse RME appliance seems to be effective for alignment of buccally tipped upper molars.

**CP 047**

Friday, June 18 - at 15:15

**ORTHOGNATHIC SURGERY PLANNING ON A THREE-DIMENSIONAL SOLID MODEL****M Erkan**<sup>1</sup>, Ş Karaçay<sup>1</sup>, E Ulker<sup>2</sup>, H Karagöz<sup>2</sup>, G Sönmez<sup>3</sup>, Departments of <sup>1</sup>Orthodontics, <sup>2</sup>Plastic and Reconstructive Surgery and <sup>3</sup>Radiology, GMMA Haydarpasha Education Hospital, İstanbul, Turkey

**AIM:** Three-dimensional (3D) volumetric imaging allows improved visualization of the morphologic deformity, planning the surgical approach, and evaluating the response to treatment. In future, this technique will replace conventional two-dimensional cephalometric planning and plaster cast model surgery. The aim

of this presentation is to show the use of a 3D solid model constructed through cone-beam computed tomography in order to plan orthognathic surgery of a patient with a Class III malocclusion and facial asymmetry.

**SUBJECT AND METHOD:** Clinical examination of a 21-year-old male patient revealed a concave profile, facial asymmetry, negative overjet, and a mandibular midline shift of 3 mm to the left. Cephalometric and frontal radiographs were taken to determine the skeletal deformity. A 3D solid model of the patient was constructed and a Le Fort I and bilateral sagittal split ramus osteotomy were performed on the model. Segments on the model were stabilized to their new locations. Triangular axial gaps between the proximal and distal osteotomized mandibular segments, which occurred depending on the rotational movement of the distal mandibular segment, were filled with dental modelling wax. Templates of the gaps were created to determine the size of the bone grafts required during surgery.

**RESULTS:** A Class I molar relationship was obtained; the midline shift, concave profile and facial asymmetry were corrected. Model surgery performed on the 3D solid model reduced the duration of surgery since the bone grafts, constructed according to the templates manufactured before surgery, had adequate dimensions to fill the gap.

**CONCLUSION:** Construction of 3D solid models through contemporary digital design and manufacturing techniques can be a useful method for surgical planning, improving communication between medical experts and patients, and facilitating customization of treatment devices. Although producing the models is expensive, using them for pre-operative planning reduces the time required and the difficulty of surgery.

#### CP 048

Friday, June 18 - at 15:15

##### ASYMMETRIC PREMAXILLARY DISTRACTION OSTEOGENESIS – A CASE REPORT

Ş Karaçay<sup>1</sup>, O Bengi<sup>2</sup>, K M Okçu<sup>3</sup>, M Erkan<sup>1</sup>, <sup>1</sup>Department of Orthodontics, GMMMA Haydarpasha Education Hospital, Istanbul and Departments of <sup>2</sup>Orthodontics and <sup>3</sup>Maxillofacial Surgery, GMMMA Center of Dental Sciences, Ankara, Turkey

**AIM:** Distraction osteogenesis (DO) is the process of bone and soft tissue regeneration through the application of graduated tensile stress. Premaxillary DO is a new method to correct mild midface deficiency while gaining space by increasing the arch parameter. Asymmetric application of premaxillary DO may be an effective method to correct midline shift, while treating an anterior crossbite. The purpose of this report is to introduce asymmetric premaxillary DO.

**SUBJECT AND METHOD:** A 13-year-old female patient with a concave profile, negative overjet and a maxillary midline shift of 3 mm to left. Intraorally the molar and canine relationship was Class I on right and Class III on the left. The midline shift was due to congenital absence of the left maxillary first premolar. The patient was treated with asymmetric premaxillary DO. A maxillary anterior segmental osteotomy was performed under general anaesthesia to allow movement of the anterior segment. The distractor was cemented on the teeth and the screw was activated by the patient at a rate of 0.75 mm/day. The anterior and right posterior segments of the distractor were ligatured to each other and the anterior arms of the hyrax screw were soldered closer to the left side. Due to this mechanism, the distractor worked like a fan type screw and distracted one side more by forcing the premaxilla to rotate on its axis thus correcting the midline shift and anterior crossbite. Following the distraction and consolidation periods, treatment continued with fixed orthodontic appliances.

**RESULTS:** At the end of treatment, the overjet was corrected, space was obtained for the missing left premolar, the midline shift was corrected, the premaxillary segment and upper lip were protruded, and a linear profile was obtained.

#### CP 049

Friday, June 18 - at 15:15

##### A MODIFIED CLASS III PROTRACTION MASK ILLUSTRATED BY TWO CLINICAL CASES

N Verly, A Balon-Perin, M Charezinski, R Glineur, Department of Orthodontics and Dentofacial Orthopedics, Free University of Brussels, Erasmus Hospital, Belgium

**AIM:** To present the treatment of two Class III malocclusion patients in whom a modified protraction facemask was used. The two patients had maxillary hypoplasia in the sagittal and also in the transverse dimension.

**SUBJECTS AND METHOD:** The first case was a young girl aged 8 years 7 months who presented with a Class III malocclusion with an anterior crossbite. She was treated with a modified protraction mask (including a palatal expansion device) because of her severe long face profile. Orthopaedic treatment started with palatal expansion for 8 days. The facemask procedure was then initiated with a force of 350 g for 3 weeks and then a force of 550 g for 12/14 hours per day for six months. At the end of orthopaedic treatment, the anterior crossbite was corrected and a Class I molar relationship was achieved without increasing the long face profile and without the development of an open bite. The second case was a young boy aged 8 years 5 months with a Class III malocclusion with an anterior open bite (AOB) and a lateral left crossbite. He was treated with a modified protraction mask (including a palatal expansion device) because of his AOB. Orthopaedic treatment started with palatal expansion for 10 days. Maxillary protraction was then commenced with a force of 350 g for 2 weeks and then 550 g 12/14 hours per day for 6 months. At the end of treatment, the lateral crossbite was corrected and the open bite was closed.

**CONCLUSION:** Compared with conventional protraction mask, the unwanted side-effect, i.e. counter clockwise rotation of the maxilla was prevented. This was achieved by force application at a point near the centre of resistance of the maxillary bone.

#### CP 050

Friday, June 18 - at 15:15

##### TREATMENT OF A CLASS III MALOCCLUSION WITH THE FRÄNKEL FUNCTIONAL REGULATOR – A CASE REPORT

Z Stamenković, N Randjelović, N Nedeljković, B Novakov, S Novakov, Department of Orthodontics, University of Belgrade, Serbia

**AIM:** To present the treatment of a male patient aged 7 years 10 months at the beginning of treatment with a Fränkel appliance type III for correction of a Class III malocclusion caused by maxillary retrognathism.

**MATERIALS AND METHOD:** Study casts, dental pantomographs and lateral cephalograms were analysed before and after treatment. ANB angle was –2 degrees (SNA 78°, SNB 80°. The patient had a negative overjet of –3 mm. He was treated with a functional appliance for 19 months for correction of the malocclusion, following which for the next 12 months it was used only at night for retention.

**RESULTS:** After nine months of treatment the overjet was corrected. The patient had a correct Class I sagittal relationship, ANB angle 2 degrees, caused by sagittal growth of upper jaw (SNA 82°, SNB 80°).

**RESULTS:** The Fränkel functional regulator III was successful in improving growth of the upper arch during early treatment of a Class III malocclusion.

#### CP 051

Friday, June 18 - at 15:15

##### EVALUATION OF EARLY CLASS III TREATMENT WITH THE FRÄNKEL APPLIANCE

M Lajh, Medical Health Center Dr. Adolf Drolc, Maribor, Slovenia

**AIM:** To evaluate the effectiveness of treatment with a Fränkel functional appliance on growing children with Class III malocclusions.

**SUBJECTS AND METHOD:** Ten children (4 boys, 6 girls) with a Class III malocclusion, aged from 4.0 to 7.9 years. Clinical examination, facial and intraoral photographs, study cast analysis, panoramic radiographs and lateral cephalometric analysis were obtained at the beginning and after 18 months of treatment with a Fränkel III appliance. During intraoral examination, overjet, sagittal occlusal relationship of the canines and first molars and anterior and posterior arch width were recorded. Orofacial functions such as lip seal, mode of breathing, swallowing pattern and tongue posture were examined. For all subjects a negative overjet, mesiocclusion, immature swallowing pattern and incorrect tongue posture on the mouth floor were diagnosed at baseline. A Class III facial appearance was also observed.

**RESULTS:** After 18 months of treatment a Class I occlusion with a positive overjet was achieved. The swallowing pattern was improved and correct tongue posture was established.

**CONCLUSION:** As Class III subjects can be successfully treated by modifying the growth pattern, early treatment with a Fränkel appliance is recommended. The Fränkel functional regulator alters both mandibular posture and the contour of the facial soft tissues. It serves as an upper arch expansion appliance and improves irregular orofacial functions.

### CP 052

Friday, June 18 - at 16:00

#### EARLY TREATMENT OF A TRUE FORCED BITE WITH THE FRÄNKEL APPLIANCE

**D Kramberger**, A Fekonja, Orthodontic Department, Health Centre, Maribor, Slovenia

**AIM:** A true force bite (TFB) has a favourable prognosis and should be differentiated from a pseudo force bite (PFB) in which the prognosis of treatment is unfavourable. Patients with a TFB shift the mandible forward during the articular phase and into a prognathic forced bite. The aim of this presentation is to show the importance of treating a TFB during eruption of the permanent incisors.

**SUBJECT AND METHOD:** A male subject in which early treatment of TFB was undertaken. He had a dentoalveolar Class III relationship on the right side, an anterior crossbite, a shift of the maxillary midline to the left, a retrognathic midface profile and functional disturbances related to speech problems. He wore a Fränkel type III appliance for 16 hours a day. The cephalogram showed a normal axial position of the incisors. Using a functional appliance the mandible is held in a posterior position. The construction bite was taken in a tête-à-tête position, which allowed the upper incisors to erupt in a proper axial position with correct overbite.

**RESULTS:** A correct overbite was obtained in nine months that resulted in occlusal stability.

**CONCLUSION:** Early treatment with the Fränkel appliance has many advantages. Treatment time is relatively short, it is pleasant for the patient and contributes to profile aesthetics as soon as the incisors are in a correct position.

### CP 053

Friday, June 18 - at 16:00

#### EARLY TREATMENT OF A CLASS III MALOCCLUSION WITH A MODIFIED MAXILLARY PROTRACTION FACEMASK

**M Stojković**, B Zečević, I Stojković, Private Practice

**AIM:** To evaluate the clinical effects of a modified maxillary protraction facemask. Specially designed intraoral vestibulo-palatal arches were used that carry the point of force application from the centre of resistance of the maxillary mixed dentition.

**SUBJECTS AND METHOD:** Ten patients (mean age: 8.5 years). Modified intraoral vestibulo-palatal removable arches were constructed. Protraction therapy using a combination of the specially designed intraoral arches and a facemask was started. In a young child facemask treatment can produce some forward displacement of the maxilla. The point of force application has to pass through the centre of resistance of the maxilla, and to have paral-

lel application of the elastic force on both sides. A unilateral force of 500 g was applied and the patients were instructed to wear the facemask for 16 hours a day. The average treatment time was 9.5 months. Cephalometric lateral radiographs were taken before and immediately after removal of the intraoral arches, superimposed on the N-S line, were analysed the point A.

**RESULTS:** The modified intraoral arches were effective in protracting the maxilla. Point A moved anteriorly by 2.91 mm. All three effects; forward movement of the maxilla, forward displacement of the maxillary teeth and downward-backward rotation of the mandible were typical responses to facemask treatment. The maxillary occlusal plane showed significant clockwise rotation.

**CONCLUSION:** In young children, the magnitude of skeletal change is usually greater than the dental change. In adolescents, the change is almost entirely dental.

### CP 054

Friday, June 18 - at 16:00

#### SUCCESSFUL ORTHOGNATHIC TREATMENT OF A SKELETAL CLASS III PATIENT USING THREE-DIMENSIONAL SIMULATION

**I-R Choi**, National Health Insurance Corporation, Ilsan Hospital, Yonsei University College of Medicine, Seoul, Korea

**AIM:** To date, orthognathic surgery planning depends on 'paper surgery' using postero-anterior and lateral cephalometric radiographs. Although two-dimensional morphing programs for predicting the results of orthognathic surgery have been developed, they have several limitations. Three-dimensional (3D) simulation will be a good substitute, but techniques for improvement of accuracy are needed. The aim of this presentation is to show a patient with a skeletal Class III malocclusion and facial asymmetry who successfully underwent orthognathic surgery using a 3D stimulation program.

**SUBJECT AND METHOD:** After pre-operative orthodontic treatment, 3D images were constructed using 3D computed tomographs (CT) and an impression was taken for a post-operative occlusal cast. The cast was scanned with a 3D laser scanner. The scanned cast and CT images were overlapped at the occlusal surface of the teeth and then the results of the two-jaw surgery were simulated using the Rapidform™ 2006 program.

**RESULTS:** When the 3D images were simulated with paper surgery data (1.5 mm canting correction, 4 mm posterior impaction, 1.5 mm rotation of the maxilla, impaction and vertical reduction osteotomy [intraoral vertical ramus osteotomy (IVRO) set back 8 mm], mandibular menton point still remained in an asymmetric position. By changing the treatment plan (4 mm posterior impaction, 2 mm rotation and 2 mm total shift of the maxilla, IVRO set back 7 mm), a successful outcome was obtained.

**CONCLUSION:** Post-operative planned occlusion images were integrated directly on to the CT images to improve the accuracy of the 3D simulations thus allowing precise diagnosis of orthognathic surgery.

### CP 055

Friday, June 18 - at 16:00

#### USE OF BONE ANCHORS IN A CLASS III SKELETAL PATIENT. A CASE REPORT

**M Charezinski**<sup>1</sup>, A Balon-Perin<sup>1</sup>, S Giot-Wirgot<sup>2</sup>, R Glineur<sup>1,3</sup>, <sup>1</sup>Clinique D'Orthodontie and <sup>3</sup>Service de Chirurgie Maxillo-Faciale - Hopital Erasme and <sup>2</sup>Service de Stomatologie - Hopital des Enfants Huderf and Chu Saint-Pierre, Universite Libre de Bruxelles, Belgium

**AIM:** Controlled distalization of a complete maxillary arch represents a challenge in orthodontic treatment. Nowadays bone anchors are suggested in such anchorage issues. The aim of this presentation is to show the orthodontic and surgical treatment using two bone anchors, Bollard miniplates.

**SUBJECT AND METHOD:** A young female aged 16 years 3 months. The main clinical and radiographic features before treatment were a hyperdivergent skeletal and dental Class III (ANB: -8°;

AO/BO: -15 mm), extracted four first premolars without residual space, proclined central upper incisors (I/F: 130°), upper incisor root resorption, anterior crossbite and anterior open bite. The treatment plan consisted of an orthodontic dental decompensation phase, a surgical phase and a finishing orthodontic phase. Bollard miniplates were successfully used in the first phase with two segmental wires (stainless steel 0.016 X 0.022 inch).

**RESULT AND CONCLUSION:** Treatment resulted in molar distalization of 5 mm. Interestingly, the mechanics also had a distalizing effect on all mesial teeth (premolar, canine and incisors). The use of bone anchors allowed controlled distalization of a complete maxillary arch and successful orthodontic and surgical Class III treatment.

#### CP 056

Friday, June 18 - at 16:00

##### COMBINED ORTHODONTIC-SURGICAL THERAPY IN A CLASS III PATIENT WITH LATEROGNATHIA

**M Janošević,** G Filipović, Department of Orthodontics, Medical Faculty, University of Niš, Serbia

**AIM:** To show the results of interdisciplinary treatment of a patient with a Class III malocclusion and laterognathia.

**SUBJECT AND METHOD:** A 19-year-old patient with dominant mandibular prognathism and laterognathia of the mandible to the left, with aesthetic and functional problems, treated orthodontic-surgically. Cephalometric analysis showed a skeletal Class III malocclusion with dominant mandibular prognathism, an anterior type of face, longer mandibular corpus, upper incisor protrusion and lower incisor retroclination. Pre-surgical orthodontic treatment was carried out by modelling of the transverse relationship and levelling of the dental arches. Surgical correction included a mandibular bilateral sagittal split osteotomy with rotation of the mandible to the right. Post-operative orthodontic treatment included fine detailing of the occlusion with the aim of achieving maximum intercuspation.

**RESULTS:** Satisfactory aesthetic and functional results were achieved by combined orthodontic-surgical treatment.

#### CP 057

Friday, June 18 - at 16:00

##### TRANSVERSE AND SAGITTAL EFFICACY OF A MODIFIED PALATAL EXPANDER IN SKELETAL CLASS III TREATMENT

**A Rexhepi<sup>1</sup>,** S Rexhepi<sup>2</sup>, A Ismani<sup>3</sup>, F Sali<sup>1</sup>, B Mehmeti<sup>4</sup>, <sup>1</sup>Clinic of Orthodontics, University Dental Clinical Centre 'St Panntelejmon' Skopje, <sup>2</sup>Private practice Struga, <sup>3</sup>Clinic of Maxillofacial Surgery Skopje, Former Yugoslav Republic of Macedonia and <sup>4</sup>Clinic of Orthodontics, Prishtina, Kosovo

**AIM:** To present the clinical dentofacial effects of a modified palatal expander in skeletal Class III subjects with maxillary sagittal and transverse insufficiency.

**SUBJECT AND METHOD:** A male patient 14 years 3 months old with a Class III malocclusion, posterior crossbite and ectopically positioned maxillary canines. A modified Bertoni palatal expander was used for transverse expansion and sagittal protraction. Cephalometric radiographs and dental casts were obtained at the beginning and end of expansion. Total screw activation was 7.5 mm within 3 weeks.

**RESULTS:** The mean maxillary arch expansion in the molar region was 7 mm (measured comparing before and after screw activation). Sagittal growth of the maxilla increased; the mean value of SNA to SNB changed from 45.5 to 52.5 mm.

**CONCLUSION:** Maxillary volume changes in the sagittal and transverse plane, also free space formed for aligning both the maxillary canines into the dental arch to correct without surgery this skeletal Class III.

#### CP 058

Friday, June 18 - at 16:00

##### MODERATE RELAPSE AFTER MANDIBULAR OSTEOTOMY IN SKELETAL CLASS III SUBJECTS

**A Ismani<sup>1</sup>,** S Rexhepi<sup>2</sup>, L Mustafa<sup>3</sup>, A Rexhepi<sup>4</sup>, <sup>1</sup>Maxillofacial Surgery Skopje, <sup>2</sup>Private practice Struga, <sup>3</sup>Private practice Gostivar and <sup>4</sup>Clinic of Orthodontics, University Dental Clinical Centre 'St Panntelejmon', Skopje, Former Yugoslav Republic of Macedonia

**AIM:** To evaluate lower arch stability after surgical correction in skeletal Class III subjects.

**SUBJECTS AND METHOD:** Ten young patients (6 females, 4 male mean age 18.6 years) with a skeletal Class III deformity who had undergone a mandibular setback with bilateral sagittal split osteotomy. Lateral cephalograms were taken obtained before (T0) and 1.5 (T1) and 3.5 years post-surgery (T2).

**RESULTS:** There was a significant decrease of the mandibular and occlusal plane angles from T1 to T2, and retraction of points B, Me, and Go. Moderate vertical relapse was seen at points B, Me and Go from T1 to T2. The mandibular setback at T1 of 2.9 mm relapsed to 0.8 mm at T2.

**CONCLUSION:** A moderate relapse was observed in mandibular movements 3.5 years after osteotomy.

#### CP 059

Friday, June 18 - at 16:00

##### EARLY TREATMENT OF A CLASS III ANTERIOR SKELETAL CROSSBITE WITH AN ORAL SHIELD

**M Yanagisawa<sup>1</sup>,** W Machata<sup>2</sup>, K Sato<sup>3</sup>, M Yanagisawa<sup>4</sup>, <sup>1</sup>Chofu Orthodontic Clinic, Tokyo Japan, <sup>2</sup>(closed private clinic) Sheboygan Wisconsin, USA <sup>3</sup>Department of Pediatric Dentistry and <sup>4</sup>Nihon University School of Dentistry, Tokyo Japan

**AIM:** To demonstrate the effectiveness of the YC3 shield appliance (a muscle-training appliances) used in the early treatment of skeletal anterior crossbites.

**SUBJECT AND METHOD:** A 6 year 3 month old female patients. At initial examination she presented with a Hellman's dental age of IIC with an overjet of -1.9 mm, overbite of +2.4 mm, and a first molar mesial step terminal plane. The facial profile was straight. The patient was diagnosed with a skeletal anterior crossbite. Treatment changes were observed with lateral cephalometric films taken at 11 months, 4 years 3 months, and 9 years 9 months after the initial examination.

**RESULTS:** Clinically, the overjet showed improvement within 4 months after initial placement of the appliance. The lateral cephalometric film, taken 9 years 9 months after the first examination, showed positive increases in SNA, ANB, and U1 to FH. SNA was 1.5 degrees, ANB was 2.7 degrees, and U1 to FH increased by 13.2 degrees. ANB angle improved from minus to plus. The occlusal plane relationship to FH angle decreased by 9 degrees, a result of a flattening of the occlusal plane. Flattening of the occlusal plane contributed to the improvement of AB to the occlusal plane.

**CONCLUSION:** Treatment with the YC3 shield improved the reverse overjet (crossbite correction) in a short period and produced positive changes in craniofacial morphology. The YC3 shield is effective in the early treatment of skeletal anterior crossbite cases.

#### CP 060

Friday, June 18 - at 16:00

##### SOFT TISSUES PROFILE CHANGES DURING TREATMENT OF GROWING CLASS III PATIENTS

**E Teodorescu<sup>1</sup>** V-M Milicescu<sup>1</sup>, E Ionescu<sup>1</sup>, S Milicescu Jr.<sup>2</sup>, M Crisan<sup>1</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Fixed Prosthodontics, 'Carol Davila' University of Medicine and Pharmacy, Bucharest, Romania

**AIM:** Compared with other skeletal anomalies, Class III malocclusions are frequently accompanied by facial changes.

**MATERIALS AND METHOD:** Comparative analysis, before and after treatment of profile photographs, the mesh-diagram and the

distances in soft tissues profile changes between nasion-soft tissue nasion, hyperhinion-soft tissue hyperhinion, subnasale-soft tissue subnasale, point A-soft tissue point A, labrale superior-upper incisor tip, stomion-upper incisor tip, labrale inferior-lower incisor tip, labiomentale sulcus-point B, pogonion-soft tissue pogonion, menton-soft tissue menton. The data were also compared normal values for age and gender.

**RESULTS:** Treatment of moderate Class III skeletal anomalies during the mixed dentition period allowed the shaping of the facial profile into the variability norms. The treatment of severe Class III patients frequently offers satisfactory aesthetic results; in some cases the soft tissues compensate the bony lack of balance.

## CP 061

Friday, June 18 - at 16:00

### TREATMENT OF A CLASS III MALOCCLUSION WITH A COMBINATION OF FRÄNKEL AND FIXED APPLIANCES

**E Zaletel**, Orthos, Ljubljana, Slovenia

**AIM:** Modification of growth might be possible in the period of maturation with an orthopaedic force or with a functional appliance. Dentoalveolar malformations can be corrected with the assistance of neuromuscular activity induced by a functional appliance. The purpose of this study was to evaluate skeletal changes, orofacial functions and facial appearance in a growing child, treated with the Fränkel type III appliance, a protraction mask and fixed appliances.

**SUBJECT AND METHOD:** A 4-year-old female patient with a Class R III malocclusion, crossbite and deviated orofacial functions. Clinical examination, dental casts, facial and intraoral photographs and lateral cephalograms were taken after every therapeutic phase. After 16 months of treatment and excellent cooperation, correction of the crossbite, open bite, negative overjet and sagittal and transverse relationships were obtained. During the mixed dentition, treatment was focused on deviated orofacial functions using a modified Fränkel appliance. In order to achieve correction of the occlusion, a fixed appliance in combination with a protraction mask was used.

**RESULTS:** After eight years of therapy, the relationships between the upper and lower jaws in the sagittal, vertical and transverse plane were improved. Moreover, nasal breathing, correct lip seal and improvement of the facial appearance were attained. However, incorrect tongue posture and function remained. According to the analysis of the dental casts and lateral cephalograms, the effectiveness of treatment was evaluated. Cephalometric superimpositions showed more downward and forward growth of the maxilla than the mandible

**CONCLUSION:** Early orthopaedic treatment with the protraction mask and with a Fränkel appliance improved the skeletal relationship, facial appearance and some of the functional irregularities.

## CP 062

Friday, June 18 - at 16:45

### EARLY TREATMENT OF A CLASS III MALOCCLUSION WITH THE MODIFIED FRÄNKEL APPLIANCE

**S Melink**<sup>1</sup>, S Lah Kravanja<sup>1</sup>, M Ovsenik<sup>2</sup>, F Farčnik<sup>3</sup>, Departments of Orthodontics, <sup>1</sup>University Medical Centre and <sup>2</sup>University of Ljubljana, and <sup>3</sup>Zavod Orthos, Ljubljana, Slovenia

**AIM:** The morphological characteristics of Class III malocclusions are usually accompanied by functional problems of lip seal, incorrect breathing and incorrect tongue posture and function. The aim of this study was to evaluate treatment success in a patient with a Class III malocclusion using the modified Fränkel type III (FR-II) appliance.

**SUBJECT AND METHOD:** A 7 year 11 month old girl with a Class III malocclusion. Functional diagnosis revealed incorrect breathing through the mouth, incorrect tongue posture and an infantile swallowing pattern. Study cast analysis showed a Class III molar relationship with a reverse overjet (-4 mm) and a midline shift to the left (2.5 mm). A lateral cephalogram indicated a retrognathic

maxilla and mandible, a Class III skeletal relationship and a vertical growth pattern. The modified FR-III was designed by Farčnik (2008). An acrylic plate was added in the middle of the transpalatal arch of the FR-III for chewing gum placement, to stimulate a correct tongue posture on the palate. A lingual shield was designed to obtain a correct position of the tongue in the oral cavity (on the palate). The patient was instructed to wear the appliance full-time.

**RESULTS:** After one year of treatment, there was a significant improvement of orofacial functions. Nose breathing, a correct tongue posture, and a somatic swallowing pattern were registered. Facial appearance as well as the morphological irregularities of the occlusion significantly improved.

**CONCLUSION:** According to Fränkel's belief, the dynamic play of forces was able, through the epigenetic path, to influence genetically coded osteogenetic growth areas of the orofacial complex.

## CP 063

Friday, June 18 - at 16:45

### TREATMENT OF A CLASS III SUBJECT WITH AN OPEN BITE IN MIXED DENTITION WITH A MODIFIED FRÄNKEL APPLIANCE

**M Malinger**, F Farčnik, J Primožič, Orthos, Ljubljana, Slovenia

**AIM:** Long-term observation shows that mandibular prognathism with a hereditary background cannot always be successfully treated with the conventional Fränkel type 3 appliance. Usually it is not possible to intercept lateral widening of the mandible caused by unfavourable tongue position on the mouth floor. To overcome this incorrect tongue position, additional elements were incorporated into the conventional Fränkel appliance. The aim of this study is to present the use of the modified Fränkel type 3 appliance in the early mixed dentition period.

**SUBJECT AND METHOD:** A 7-year-old girl with a Class III malocclusion and an open bite who started treatment with a modified Fränkel appliance in March 2008. The appliance had two additional elements – a lingual shield that eliminated the pressure of the tongue on the lingual surface of the lower jaw and forced the tongue to a more cranial position, and a small concave plate in the middle of the palatal arch in which a source of pleasant taste and aroma could be placed, to stimulate the tip of the tongue to an upright position.

**RESULTS:** After one year of treatment, a Class I molar relationship and an overjet and overbite of 1 mm were obtained. SNA and SNB angles did not change, while Wits appraisal improved from -4 to -1.5 mm. Treatment resulted in a correct lip seal and the profile greatly improved.

**CONCLUSION:** Treatment of a Class III malocclusion with an open bite could be more successful with a modified functional regulator than with the conventional Fränkel appliance.

## CP 064

Friday, June 18 - at 16:45

### TREATMENT OF CLASS III MALOCCLUSIONS – THE KEY ROLE OF THE OCCLUSAL PLANE

**B Grollemund**, O Matern, C Schwartz, W Bacon, J-L Raymond, Functional Unit of Dentofacial Orthopaedics, University Hospital, Strasbourg, France

**AIM:** Class III malocclusions are associated with various dysfunctions including chewing disorders. The restoration of a physiological normal mastication constitutes an important aim of treatment. The aim of this presentation is to show a method of Class III malocclusion treatment based on a correct re-orientation of the occlusal plane (OP) in order to achieve an optimal masticatory mechanism that is essential for long-term stability. Class III patients present an OP that is more or less inclined backward and downward. Orthopaedic correction should not only address protraction of the maxilla but also correct the orientation of the OP.

**MATERIALS AND METHOD:** Maxillary protraction with a face-mask and extrusion of the lower molars allow reorientation of the OP. This repositioning of the lower molars determines the fi-

nal three-dimensional (3D) orientation of the OP and is obtained using a bonded acrylic-splint that does not cover the maxillary incisors, with a rapid palatal expander and facemask treatment. During six to twelve months of treatment, several 3D grindings are made in order to achieve sagittal and vertical orientation of the desired OP.

**RESULTS:** Early normalization of mastication via reorientation of the OP has an immediate consequence as it breaks the 'dysmorpho-functional' spiral (described by Delaire). Correction of the Class III skeletal dysmorphism also allows normalization of muscle posture (labio-mental, lingual, velo-pharyngeal) and of orofacial functions (nasal ventilation, deglutition, mastication). Thus rehabilitation of physiological mastication results in a decreased likelihood of relapse.

**CONCLUSION:** The reorientation of the OP is essential for the rehabilitation of mastication, which is the ultimate goal of orthopaedic treatment for Class III malocclusions, and is probably the best option for overall long-term stability. It is easier to achieve when the OP is short; i.e. in young patients.

#### CP 065

Friday, June 18 - at 16:45

##### A SEVERE CLASS III MALOCCLUSION PATIENT TREATED WITH QUADHELIX

**R Janev**, L Jakupi, E Arsova, S Bekir, MACEDONIA, THE FORMER YUGOSLAV REPUBLIC

**AIM:** To present the treatment of a 13-year-old patient with a constricted maxilla, impacted maxillary canines and a lateral open bite. In addition the upper left first premolar had been extracted.

**MATERIALS AND METHOD:** A fixed orthodontic appliance with a quadhelix was used at the beginning of treatment. The primary maxillary canines were extracted and then a corticotomy was performed of both maxillary permanent canines with a vestibular approach. In the lower jaw, the first premolars were extracted.

**RESULTS:** After 32 months of treatment the posterior crossbite, and open bite were corrected, the upper canines were placed in their correct position in the dental arch and the lower incisors were retracted into the premolar extraction spaces. Facial appearance was satisfactory after treatment.

**CONCLUSION:** A comprehensive approach to a complex case is needed for successful orthodontic treatment.

#### CP 066

Friday, June 18 - at 16:45

##### THE CLEAR ALIGNER TECHNIQUE IN THE TREATMENT OF PRE-SURGICAL CLASS III MALOCCLUSION SUBJECTS

**M Greco**<sup>1</sup>, S Marrocco<sup>2</sup>, D Scopelliti<sup>2</sup>, A Giancotti<sup>1</sup>, <sup>1</sup>University of Rome Tor Vergata Fatebenefratelli Hospital and <sup>2</sup>Department of Maxillo-Facial Surgery, 'Santo Spirito' Hospital, Rome, Italy

**AIM:** Invisalign treatment is gaining wider applications in the treatment of different malocclusions, not only for simple arch alignment. The purpose of this study was to evaluate the efficacy of the Invisalign system in the treatment of anterior crossbite.

**SUBJECTS AND METHOD:** Nine adult patients (mean age 26.5 years) with a skeletal Class III malocclusion, molar and canine Class III relationship, hyperdivergent skeletal pattern, negative overjet (-4 mm) and negative overbite (-3 mm). The treatment plan focused on complete correction of the skeletal malocclusion by means of pre-surgical orthodontic treatment followed by a maxillary Le Fort 1 intervention associated with a mandibular osteotomy. All patients were treated in the first pre-surgical phase by means of Invisalign aligners in order to achieve arch levelling and alignment. Radiographs were obtained at the beginning T1, pre-surgical (T2) and end of treatment (T3), and the clincheck software was assessed by orthodontists and the surgeon in order to achieve arch coordination. A few days before the planned intervention, fixed appliance were positioned. Orthodontic finishing was achieved with fixed appliances.

**RESULTS:** A normal overjet and overbite and a Class I molar and canine relationship was achieved in all patients. The skeletal Class III malocclusion was corrected to a Class I occlusion. Cephalometric superimposition (T2-T1) showed a satisfactory change in maxillary and mandibular incisor inclination in the initial phase of Invisalign treatment.

**CONCLUSION:** The Invisalign technique proved to be an efficient option for pre-surgical treatment of skeletal Class III malocclusions.

#### CP 067

Friday, June 18 - at 16:45

##### DESIGN AND DEVELOPMENT OF A FACEMASK FITTED WITH A TIMING DEVICE

**C Schwartz**, C Freyss, O Matern, R Mathis, J-L Raymond, Department of Orthodontics, University Hospital, Strasbourg, France

Orthopaedic treatment of Class III malocclusions usually includes a fixed intra-oral device (splint, double dental-arch, traction support or even a maxillary expander, and an extra-oral device, generally a facial mask (Delaire, Petit). The actual time the facemask is worn is a downside to this treatment because it relies entirely on the patient's and their parents' cooperation. To date the actual time the mask is worn cannot be objectively measured. Failure and success of such treatment is, therefore, difficult to assess. Through collaboration with an engineering school (INSA, Strasbourg, France) a timer was developed that can be fitted to the masks currently on the market. After clinical experimentation, this device appeared efficient.

This device aims to assess the actual wear time of the mask with the purpose of controlling not only the efficiency and correct functioning of the treatment, but also to assist communication with parents concerning the importance of rigorous wearing of the mask. In the long-term this could also allow better adjustment of the recommendations for orthopaedic treatment of Class III malocclusions and, through retrospective statistical studies, to identify the best conditions ensuring successful correction.

#### CP 068

Friday, June 18 - at 16:45

##### EARLY ORTHOPAEDIC TREATMENT OF A CLASS III MALOCCLUSION – A THREE-DIMENSIONAL EVALUATION

**V Božnik**<sup>1</sup>, M Verdenik<sup>1</sup>, J Volk<sup>1</sup>, A Eigon<sup>2</sup>, M Ovsenik<sup>1</sup>,

<sup>1</sup>Department of Orthodontics, University of Ljubljana and

<sup>2</sup>Stomatološka ordinacija, Ljubljana, Slovenia

**AIM:** To evaluate the effectiveness of treatment of Class III malocclusion in the early mixed dentition period with a facemask using three-dimensional (3D) soft tissue analysis and 3D ultrasonography.

**SUBJECT AND METHOD:** An 8 year-old girl with a Class III malocclusion. Clinical examination, facial and intraoral photographs, dental casts, a dental pantomogram, lateral cephalogram, ultrasound evaluation of tongue posture, and soft tissue analysis of the girl's face were undertaken at baseline (T0), and after orthopaedic treatment with a facemask for 6 months and a modified Fränkel appliance for 5 months (T1). The facial scans (T0-T1) were superimposed over the average facial template, constructed using facial scans of 21 girls, aged 8.5 years. Differences in facial morphology were assessed quantitatively and qualitatively.

**RESULTS:** At T0, 3D analysis showed a retrognathic midface and a prognathic chin. The girl had a more prominent chin and a less prominent midface area compared with the average face. At T1, 3D soft tissue analysis showed a less retrognathic midface, the facial appearance had become almost similar to the average face. Ultrasound at T0 showed an irregular tongue posture. At T1 the tongue still remained on the floor of the mouth.

**CONCLUSION:** Early orthopaedic treatment of the Class III malocclusion improved facial appearance, but functional irregularities still remained.

**CP 069**

Friday, June 18 - at 16:45

**PROFESSIONALS' AND LAYPERSONS' APPRECIATION OF CLASS III SURGICAL CORRECTION OPTIONS**

**P Christou**<sup>1</sup>, M Fabr  <sup>2</sup>, C Mossaz<sup>3</sup>, S Kiliari  s<sup>3</sup>, <sup>1</sup>Athens University, Greece and <sup>2</sup>Private practice, Gland and <sup>3</sup>Geneva University, Switzerland

**AIMS:** To compare laypersons' and professionals' perception of the soft tissue profiles of Class III adults, and to evaluate which cephalometric variables are likely to influence the profile assessment score (PAS).

**MATERIALS AND METHOD:** Lateral headfilms and coloured profile photographs of 18 skeletal Class III (10 males, 8 female) adult Caucasian patients (mean age of 24.5 years) prior to surgery, and four male and five female Caucasian patients (mean age of 27.4 years) with a dental Class I occlusion and no major skeletal discrepancy. The aesthetics of each profile photograph was evaluated by 18 laypersons and 18 orthodontists using a 10-graded visual analogue scale (VAS) and a standard profile for calibration. Intra-observer reliability was tested and no significant error was found. Paired and unpaired t-tests were used to compare the scores. The association between various cephalometric variables and the PAS was tested.

**RESULTS:** Laypersons were less critical compared with orthodontists in their evaluation of the Class III profiles (+0.3 grade on the VAS) as well as the reference profiles (+0.7). The reference profiles were found to be more attractive than the Class III profiles by both laypersons and orthodontists (+2.3 grades). The degree of facial concavity had a negative predictive value for the orthodontists' and laypersons' evaluations.

**CONCLUSION:** The degree of facial concavity together with the steepness of the mandibular plane were negatively predictive factors for the PAS given by the orthodontists.

**CP 070**

Friday, June 18 - at 16:45

**CONTEMPORARY EVALUATION OF TONGUE FUNCTION AND POSTURE – A CASE REPORT**

**D Kramberger**<sup>1</sup>, J Volk<sup>2</sup>, M Lajh<sup>1</sup>, M Mu  i  <sup>3</sup>, M Ovsenik<sup>4</sup>, <sup>1</sup>Health Centre Maribor, <sup>2</sup>Department of Orthodontics, University Medical Centre, <sup>3</sup>Institute of Oncology, and <sup>4</sup>Department of Orthodontics, Medical Faculty, University of Ljubljana, Slovenia

**AIM:** To evaluate tongue posture using three-dimensional (3D) ultrasonography.

**SUBJECT AND METHOD:** A 6 year-old girl with a Class III malocclusion. During clinical examination an irregular tongue function was registered. The facial appearance was Class III as well as the occlusion of the first permanent molars. Study cast analysis revealed a reverse overjet, a crossbite of 52-63 and a midline deviation. A transverse discrepancy was measured between the primary canines (0.5 mm) and between the molars (1 mm). Swallow pattern and tongue posture were evaluated using ultrasonography. Voluson 730 Expert with a 3D convex transducer was used to obtain two and 3D ultrasound images of the tongue.

**RESULTS:** The duration of the swallowing cycle was prolonged. A visceral swallowing pattern was diagnosed and the tongue posture was registered on the mouth floor.

**CONCLUSION:** Ultrasonography is a non-invasive, simple diagnostic tool for assessment of tongue function and posture. It enables objective evaluation of tongue function and posture in young growing children with a Class III malocclusion.

**CP 071**

Friday, June 18 - at 16:45

**EARLY MAXILLARY PROTRACTION WITH A NEW IMPLANT-SUPPORTED NON-COMPLIANCE AND AESTHETIC DEVICE**

**L Favero**, A Winkler, E Stellini, University of Padua, Italy

**AIM:** Hypomaxillary non-surgical Class III malocclusions in paediatric patients are treated by means of transverse maxillary expansion (if needed), combined or followed by a sagittal protraction with a Delaire mask. Micro-implant orthodontics is a widespread and useful choice, even in early treatments, allowing improved anchorage without the need for patient compliance. The aim of this research is to propose a new intraoral, maxillary fixed appliance, anchored to palatal micro-implants, for aesthetic and non-compliance maxillary protraction.

**SUBJECT AND METHOD:** After biomechanical *in vitro* testing of the device, a young Class III patient in the complete primary dentition with an anterior crossbite, non-surgical prognosis, and very low compliance profile was selected. The device was prepared with TMA mesializing springs protruding from a Nance type palatal button, anchored with two micro-implants to the palatal bone. The exerted force was applied to a steel lingual splint, bonded and banded to the maxillary teeth. A security device was added in the anterior part of the splint to avoid the risk of appliance detachment.

**RESULTS:** After 9 months of therapy, the anterior crossbite was resolved, and the sagittal discrepancy improved. Tracings confirm the Delaire-like results of the device, with lower skeletal effects and higher dentoalveolar movements. Micro-implants allowed the use of heavy active forces without the need for patient compliance, and without an encumbering or visible appliance.

**CONCLUSION:** The proposed appliance allows early non-compliance treatment of a Class III hypomaxillary non-surgical malocclusion, in a fast and aesthetic way. Since the anchorage is maxillary, the skeletal effects are limited to the anterior part of the palate, while the dental effects are higher, but sufficient to achieve a normal Class I dental relationship.

**CP 072**

Friday, June 18 - at 17:30

**INTERDISCIPLINARY TREATMENT OF A CLASS III MALOCCLUSION – A CASE REPORT\*\*\***

**G Spai  **<sup>1</sup>, M Mila  i  <sup>1</sup>, S Juri  in<sup>1</sup>, M Ovsenik<sup>1</sup>, N Ihan Hren<sup>2</sup>, Departments of <sup>1</sup>Orthodontics and <sup>2</sup>Maxillofacial and Oral Surgery, Faculty of Medicine, Ljubljana, Slovenia

**AIM:** Treatment of Class III malocclusion demands multidisciplinary approach.

**SUBJECT AND METHOD:** A 24 year-old female with a Class III malocclusion. Clinical examination, facial and intraoral photographs, dental casts, dental pantomogram, lateral cephalogram and soft tissue analysis of the girl's face were taken at baseline (T0), pre-surgery (T1) and post-treatment (T2). The superimposition technique was used in the analysis, along the S (sella)–N (nasion) line to follow treatment changes occurring in T0, T1, T2.

**RESULTS:** The results after surgery were stable following which post-surgical orthodontic treatment was undertaken. After multidisciplinary treatment, the function and facial appearance was improved.

**CONCLUSION:** The presented case demonstrates the effectiveness of combined multidisciplinary treatment and its capability to improve jaw and dental relationships as well as soft tissue and facial appearance.

**CP 073**

Friday, June 18 - at 17:30

**FORWARD TRACTION ON THE MAXILLA IN CLEFT LIP AND PALATE PATIENT\*\*\***

**K Verhovec**<sup>1</sup>, M Dreven  ek<sup>2</sup>, Departments of Orthodontics, <sup>1</sup>University Medical Centre and <sup>2</sup>Faculty of Medicine, University of Ljubljana, Slovenia

**AIM:** To demonstrate a conventional orthodontic approach to correct an anterior crossbite in a unilateral cleft lip and palate (CLP) patient.

**SUBJECT AND METHOD:** A 10-year-old boy with a Class III malocclusion with a complete left-sided CLP. An upper quadhelix appliance was used in conjunction with a facemask. Through early intervention, a more basal response was obtained. When an

expanding device such as the upper quadhelix was incorporated, the palate was more susceptible to lateral expansion. A facemask produces the most dramatic results in the shortest period of time. This system has three basic components a facemask, Kobayashi hooks fixed on bonded brackets, and elastics. The facemask is composed of a forehead pad and chin pad that are connected by two heavy steel support rods. Elastics are attached to the facemask to produce a forward and downward traction on the maxilla. Forward traction results in separation of the maxillary sutures and should stimulate growth. Normally, the facemask is worn full time (about 20 hours per day) for 4 to 6 months, and then at nighttime only for an additional period of time. After 5 months of facemask treatment, appropriate incisal overjet was obtained.

**RESULTS:** Improved maxillary arch and corrected anterior cross-bite through maxillary expansion and the sagittal skeletal relation are determined.

**CONCLUSION:** Forward maxillary traction was found to be an effective phase I method of treatment for orthodontic patients with a CLP characterized by maxillary skeletal growth retardation in the period of the mixed dentition.

#### CP 074

Friday, June 18 - at 17:30

##### THREE STAGE TREATMENT OF A SKELETAL CLASS III MALOCCLUSION

**K Arko Kampus**, ZD Ormoz, Slovenia

**AIM:** To demonstrate the importance of early orthodontic treatment in a patient with a skeletal Class III malocclusion.

**SUBJECT AND METHOD:** A 7-year old girl, with a Class III malocclusion. There was no familiar background of the orthodontic malformation, but several functional disorders such as an incompetent lip seal, mouth breathing and incorrect tongue posture were observed. Study cast analysis showed a Class I molar relationship and Class III canine relationship. There was midline shift to the left side (1.8 mm) and a reverse overjet (-2 mm). The maxilla was retrognathic and the skeletal relationship Class III, with a horizontal growth pattern. Treatment commenced with a Fränkel 3 appliance in February 2004. The patient was instructed to wear the appliance full time. After 10 months the overjet and overbite were normal and the molar relationship was Class II. In view of the lack of space for the upper canines, treatment was continued with the pendulum appliance followed by fixed appliance therapy. Treatment was completed in May 2007.

**RESULTS:** Treatment resulted in a Class I molar and canine relationship, correct overbite and overjet and appropriate dental arches. Sufficient space was created for the tongue and correct tongue posture, a good lip seal and mostly nose breathing were observed.

**CONCLUSION:** Early, carefully planned, treatment, during the most critical years, resulted in a stable jaw relationship.

#### CP 075

Friday, June 18 - at 17:30

##### COMPARISON OF AN UNTREATED CLASS III MALOCCLUSION SUBJECTS COMPARED WITH AN 'AVERAGE' PATIENT - A THREE-DIMENSIONAL EVALUATION

**A Marion**, D Selan, M Verdenik, M Ovsenik, Department of Orthodontics, University of Ljubljana, Slovenia

**AIM:** The Fränkel type III appliance is used for early orthopaedic treatment in Class III growing children to improve facial appearance as well as space conditions of the oral cavity. The aim of this study was to evaluate growth and developmental characteristics in a Class III untreated patient.

**SUBJECT AND METHOD:** A boy with a Class III malocclusion (8 years) and 26 boys (mean age = 8.4 years). Facial images were taken using a three-dimensional (3D) imaging system, consisting of two high resolution laser cameras (Minolta Vivid VI910 3D). Clinical examination, facial and intraoral photographs, dental casts, a dental pantomogram, a lateral cephalogram and a 3D evaluation

of the boy's face were obtained at baseline. The patient's image was superimposed on the average facial template constructed from children with no morphological malocclusions in the mixed dentition period. A colour map was used to highlight differences between the average male shell and the boy's facial shell.

**RESULTS:** Analysis of the superimposed image showed that most of the face matched the average. A minimal difference was seen in the temporal area and nose (around 2 mm) but a greater difference was obvious at the chin. The subject has a prognathic chin of 4 mm.

**CONCLUSION:** Assessment of facial morphology is important in orthodontic diagnosis. 3D imaging enables assessment of facial morphology in a new, more objective, non-invasive and friendly way. It provides an opportunity to compare different growing samples, to study the difference between facial morphology of normocclusion on the one hand and malocclusion and facial pathology on the other.

#### CP 076

Friday, June 18 - at 17:30

##### ORTHOGNATHIC TREATMENT OF A CLASS III MALOCCLUSION - A CASE REPORT

**B Mady Maričić<sup>1</sup>**, R Cerović<sup>2</sup>, R Vidaković, M Slaj<sup>4</sup>, <sup>1</sup>Department of Paediatric Dentistry and Orthodontics, University of Rijeka, <sup>2</sup>Department of Maxillofacial and Oral Surgery, Clinical Hospital Center, <sup>3</sup>Dental Polichlinic Slaj-Anic, <sup>4</sup>Department of Orthodontics, University of Zagreb, Croatia

**AIM:** Treatment planning and treatment evaluation of Class III malocclusions is always a challenge for an interdisciplinary approach. This study reports a case treated by orthognathic surgery.

**SUBJECT AND METHOD:** A 20-year-old female with a skeletal Class III malocclusion. The clinical findings were: ANB angle, -3 degrees; overbite, -2 mm, overjet -2 mm; with a distance between the maxillary and mandibular midlines of 7 mm. Orthodontic treatment with fixed orthodontic appliances corrected the relationship between the dentoalveolar and bone structures, following which surgical correction, that included a bimaxillary osteotomy, was undertaken. To provide good and stable intercuspation vertical settling was undertaken.

**RESULTS:** Combined orthodontic and surgical treatment resulted in a Class I occlusion with aesthetically pleasing results.



## CLINICAL POSTERS

### Free Topics

CP 077

Wednesday, June 16 - at 15:15

#### APPLICATION OF ORTHODONTIC MINISCREW-IMPLANTS IN OLIGODONTIA – A CASE REPORT

**K Raftowicz-Wójcik**, J Antoszewska, B Kawala, Department of Orthodontics and Dentofacial Orthopedics, Wrocław Medical University, Poland

**AIM:** Congenital tooth agenesis can be divided into hypodontia, oligodontia and anodontia. The term hypodontia is used, when one to six teeth are missing, oligodontia when the number of missing teeth exceeds six; third molars not included and anodontia when there is complete absence of teeth. The pathogenesis of oligodontia is largely unknown it may be caused by local trauma, chemo- or radiotherapy and mutations of several genes; it may also be associated with a syndrome such as ectodermal dysplasia.

**SUBJECT AND METHOD:** A 16-year-old adolescent boy with congenital agenesis of 12 teeth: 13, 14, 15, 17, 24, 25, 27, 34, 35, 37, 45 and 47. He also had persistent primary teeth 53, 54, 55, 65, 73, 74 and 85. His medical history was had no abnormality. Clinical examination revealed a reduced lower face height and increased overbite. The treatment plan included intrusion of the upper incisors with an orthodontic miniscrew-implant inserted below the nasal spine for uprighting of the lower molars, utilizing mini-implants in the retromolar area. The primary teeth (except for 73) were extracted before alveolar augmentation, about 6 months prior to the end of active orthodontic treatment. After orthodontic and surgical treatment, implants and prosthetic replacements were planned.

**CONCLUSION:** The treatment of oligodontia is complex and requires cooperation of orthodontists, prosthodontists and oral surgeons. It is a biomechanical challenge and miniscrew implants are helpful in such cases, serving as temporary anchorage devices.

CP 078

Wednesday, June 16 - at 15:15

#### THE IMPORTANCE OF INTERCEPTIVE ORTHODONTIC TREATMENT

**J Andjelić**, S Matijević, Health Centre, Tivat, Montenegro

**AIM:** To demonstrate that adequate interceptive orthodontic treatment at an early age can increase the possibility of tooth alignment and reduce treatment time and the severity of malocclusion.

**SUBJECTS AND METHOD:** The study was carried out on pre-school children. Anamnesis data and clinical examination data were used in the research.

**RESULTS:** Thirty per cent of respondents required treatment, mostly with space maintainers, vestibular plates, chin cups or myofunctional therapy.

**CONCLUSION:** The factors that can cause malocclusion should be detected as early as possible. The teamwork of paediatric dentists and orthodontists, as well as interceptive orthodontic treatment in the primary and early mixed dentition, can eliminate aetiological factors, utilize growth power and prevent occurrence of severe malocclusions.

CP 079

Wednesday, June 16 - at 16:00

#### MODIFIED SLIDING MECHANICS IN EXTRACTION CASES WITH A BIDIMENSIONAL APPROACH

**P Mozzicato**, M Greco, A Giaccotti, University of Rome Tor Vergata-Fatebenefratelli Hospital, Italy

**AIM:** Orthodontic tooth movement is influenced by the characteristics of the applied force and the condition of the periodontal tissues of individual patients. Generally, the side-effects observed during tooth movement with friction mechanics are tipping, binding, and lack of vertical control, as well as the risk of anchorage loss and incisor extrusion. Sliding mechanics is the term used in orthodontics for controlled movement of teeth along archwires. The ideal use of sliding mechanics would be the best combination of archwire size, bracket slot and force which would translate teeth along the archwire with minimal friction, without excessive tipping, and without anchorage loss. The aim of this presentation is to illustrate modified sliding mechanics in extraction cases, using an undersized archwire during space closure with anterior anchorage in the bidimensional technique.

**SUBJECTS AND METHOD:** Ten patients with a Class I malocclusion, bimaxillary protrusion, dental midlines non-coincident, crowding in the lower arch and increased overjet and overbite. Four first premolars were extracted and during space closure a modified archwire was tested. The archwire used was a 0.018 X 0.025 stainless steel on the anterior teeth and a 0.018 inch stainless steel on the lateral and posterior teeth in order to maintain anterior anchorage. Different to the standard bidimensional procedure, NiTi coils with a force of 150 g were used instead of 300 g.

**RESULTS:** All clinical cases were successfully treated. The incisors were retracted, the axial inclinations of the teeth were corrected, and the facial profile was preserved.

**CONCLUSION:** The double section archwire is an effective alternative to optimize lateral and posterior sliding, reduce frictional resistance, improve the teeth sliding along the archwire and allow the use of lighter forces.

CP 080

Wednesday, June 16 - at 16:00

#### PATHOLOGY IN CRANIOCERVICAL JUNCTION – WHAT DOES IT MEAN?

**H Arponen**, J Waltimo-Sirén, Institute of Dentistry, University of Helsinki, Finland

The aim of this presentation is to inform and draw attention of clinicians to pathological entities affecting the craniocervical junction. The craniocervical junction comprises the anterior and posterior skull base, and atlantoaxial complex. It forms a vital passage for the nervous system. An abnormal relationship of the structures can result in compression of the brainstem when the bony portions of the junction dislocate into the posterior cranial fossa. This rare condition may be asymptomatic or lead to disturbed circulation of the cerebro-spinal fluid, vascular compromise, and neurological symptoms such as sensory abnormalities, motor myelopathy or paresis, and eventual death. Symptoms generally arise when the head is flexed. Pathology can present as posterior skull pain or a C2 sensory deficit.

Craniocervical junction pathology is termed 'basilar invagination' when the odontoid process protrudes into the foramen magnum, and 'basilar impression' when the odontoid process is positioned far above the caudal borders of the skull. Platybasia i.e. flattening of the skull base, is often associated with the above-mentioned phenomena. Pathology is congenital or progressively expressed, as in syndromes characterised by abnormal connective tissue. Diagnosis of pathology can be made from lateral radiographs, magnetic resonance or computed tomographic images. Neurosurgery is recommended when neurological symptoms are present, and cord compression is confirmed. When these features are absent, a conservative treatment approach may be applied, such as a collar. Craniocervical junction pathology can be an incidental finding in an asymptomatic patient. Thus, orthodontists, who analyse lateral skull radiographs, can have a significant role in detecting pathology, as early intervention is important before symptoms arise.

### CP 081

Wednesday, June 16 - at 16:00

#### IMPROVED SUPERELASTIC NICKEL TITANIUM WIRE FOR TREATMENT OF AN IMPACTED TOOTH WITH AN UNDEVELOPED ROOT

**C-C Chen**, Department of Dentistry, Tzu Chi General Hospital Taipei Branch, Taiwan

**AIM:** Incomplete root development sometimes occurs when teeth are impacted due to the path of eruption being blocked by other teeth or insufficient space for eruption. The aim of this research was to determine whether root development would continue following a change in direction of the eruption path using an improved superelastic Ni-Ti alloy wire (ISW), which can be adjusted by special heat treatment allowing simplified force systems in many tooth movements.

**SUBJECT AND METHOD:** A 13-year-old female patient, whose panoramic radiograph showed left mandibular second premolar impaction with incomplete root development. Surgical exposure was carried out and forced eruption was started using the ISW.

**RESULTS:** Forced eruption was successful. However root development was not complete until approximately 6 months after tooth eruption.

**CONCLUSION:** ISW could be a suitable material for forced eruption of impacted teeth. There may, however, be some delay in root development.

### CP 082

Wednesday, June 16 - at 16:00

#### CLINICAL AND RADIOGRAPHIC IMPROVEMENT DURING ORTHODONTIC TREATMENT IN PERIODONTAL PATIENTS

**F Aguilar**, S Tavera, Universidad Nacional Autonoma, Mexico

**AIM:** To report the outcome of orthodontic-periodontic therapy that included initial periodontal therapy, followed by orthodontic treatment with fixed preadjusted appliances.

**SUBJECTS AND METHOD:** Five randomly assigned patients. The patients were diagnosed with advanced stage periodontal disease. Periapical radiographs and oral photographs were taken for initial records, and every two months thereafter until the end of treatment.

**RESULTS:** Orthodontic-periodontic treatment increased the bone level and improved the clinical conditions.

**CONCLUSION:** Orthodontic treatment should be considered in patients with advanced periodontal disease, once initial periodontal therapy has been concluded.

### CP 083

Wednesday, June 16 - at 16:00

#### CEPHALOMETRIC AND DENTAL EVALUATION OF PATIENTS WITH IMPACTED LOWER SECOND MOLARS

**A Pilotto**, G Galluccio, E Barbato, Department of Orthodontics, University of Rome Sapienza, Italy

**AIM:** To determine if patients with lower second molar impaction exhibit common dental and skeletal characteristics.

**SUBJECTS AND METHOD:** Seven patients aged 13–21 years with at least one impacted lower second molar. Analysis of lateral cephalometric head films was used to determine SNA, SNB, FMA, FMIA, IMPA, ArGogMe, and  $\Sigma$  (NSAr+SArGog+ArGogMe) angles, and study casts to determine overjet, overbite, first molar and canine relationships. The analyses were based on the casts and diagnoses made at 13–14 years of age, and the measurements were compared with normal values.

**RESULTS:** The patients had a Class II malocclusion (ANB = 5.5°), counterclockwise growth pattern ( $\Sigma$  = 392.5°), smaller gonial angle (ArGogMe = 124.57°), smaller FMA angle (21.85°), and a larger IMPA and overjet.

**CONCLUSION:** Patients with impacted lower second molars have a short mandible with vertical condylar growth.

### CP 084

Wednesday, June 16 - at 16:00

#### PROFILE CHANGES IN A CLASS II DIVISION 1 PATIENT AFTER TWO-PHASE ORTHODONTIC TREATMENT

**E Baytorun**, İ Kocadereli, Department of Orthodontics, Faculty of Dentistry, Hacettepe University, Ankara, Turkey

**AIM:** To present the two-phase treatment of a Class II division 1 patient with mandibular retrusion, severe overjet and convex profile.

**SUBJECT AND METHOD:** A 12.4 year-old boy with the chief complaint of a poor aesthetic facial appearance. Cephalometric evaluation showed a skeletal and dental Class II division 1 malocclusion with a severe overjet (10 mm), a convex profile and reduced lower anterior face height angle (35°). The lower lip was trapped behind the upper incisors with a prominent labiomental fold. During phase 1, he was treated with a Fränkel-2 appliance. After achieving good facial and skeletal changes, phase 2 was commenced. The upper and lower arches were aligned and levelled with 0.018 inch slot size Roth appliances.

**RESULTS:** A Class I occlusion with a normal overjet and overbite was achieved in 14 months. The overjet was reduced from 10 to 2.5 mm and lower face height was increased from 35 to 40 degrees.

**CONCLUSION:** Functional appliance application with appropriate treatment time and good patient cooperation can result in significant changes in the facial profile and dentition.

### CP 085

Wednesday, June 16 - at 16:00

#### BIOPROGRESSIVE TREATMENT OF AN ANGLE CLASS II DIVISION 2 PATIENT BY MANDIBULAR INCISOR EXTRACTION – CASE REPORT

**A Aksoy Doğan**, H Alkış, Department of Orthodontics, Faculty of Dentistry University of Süleyman Demirel, Isparta, Turkey

**AIM:** To present the orthodontic treatment, using the bioprogressive technique, of a patient with a Class II division 2 malocclusion and mandibular canine-lateral transposition.

**SUBJECT AND METHOD:** A 12-year-old boy whose main complaint was the appearance of his anterior teeth. He had a Class II division 2 malocclusion with retroclined central maxillary incisors and a deep overbite of 5 mm. Intraoral examination showed mandibular canine-lateral incisor transposition in the right quadrant of the dental arch. The lateral incisor was extracted as the panoramic radiograph revealed the presence of lateral incisor root resorption. Cervical headgear was used for molar distalization. After

levelling of the maxillary and mandibular arch, 0.016 X 0.022 inch reverse curve archwires were used for three months to intrude the maxillary and mandibular anterior teeth and open the bite.

**RESULTS:** The teeth were successfully aligned in their correct positions and an Angle Class I canine and molar relationship was achieved after 1 year 9 months of active treatment. Cephalometric analysis showed that ANB, 1-1(°) angle had decreased whilst Co-Gn, Pg-N, ANS-Me measurements and GoGn-SN (°) angle increased. The mandibular plane to FH, upper molar position and lower face height increased.

**CONCLUSION:** Cervical headgear in combination with reverse curve archwires is a treatment alternative for correction of a Class II malocclusion and deep bite opening.

## CP 086

Wednesday, June 16 - at 16:00

### CRANIOFACIAL GROWTH AND DEVELOPMENT IN PATIENTS WITH AMELOGENESIS IMPERFECTA

**I Eleftheriadi**, F Synodinos, E Inglezos, M Papagrigrorakis, Department of Orthodontics, Dental School, University of Athens, Greece

**AIM:** To review current data on issues pertaining to growth and development of the craniofacial structures of patients with amelogenesis imperfecta (AI)

**MATERIALS AND METHOD:** Characteristic clinical cases from the files of the Orthodontic Clinic of the Dental School, University of Athens, and a private orthodontic practice of Athens. For all patients included in the study a series of (a minimum two) lateral cephalometric radiographs was available, ranging from their initial admission to the clinic to their early adulthood.

**RESULTS:** AI represents a relative rare (prevalence ranging 1/700-1/14000) pathological condition, characterized by the hypoplastic/hypomineralised enamel of almost all teeth. It may present itself in a solitary or syndromic form. In all AI cases some kind of inheritance pattern was identified. Several genes have been shown to be implicated in the pathogenesis of AI, including AMELX, ENAM, KLK4 and MMP20. In most cases of AI, craniofacial structures are also affected including their retrognathic mandibular Class II malocclusion, in addition to an increased lower face height associated with a skeletal and dental open bite.

**CONCLUSION:** Restorative dentistry and orthodontics are equally involved in the treatment of AI, from the initial stages of the development of the primary dentition until early adulthood. Timely application of preventive and restorative methods to facilitate dental health, early orthodontic intervention in all cases in order to minimize the skeletal components of orthodontic problems, are the most important factors safeguarding the quality of life of AI patients.

## CP 087

Wednesday, June 16 - at 16:00

### MICROGNATHIA AS A PROGNOSTIC FACTOR FOR OBSTRUCTIVE SLEEP APNOEA IN PATIENTS WITH SYNDROMES

**F Synodinos**<sup>1</sup>, I Eleftheriadi<sup>1</sup>, A Amfiloxiou<sup>2</sup>, M Papagrigrorakis<sup>1</sup>,  
<sup>1</sup>Department of Orthodontics, Dental School, University of Athens and <sup>2</sup>Sismanoglio Hospital, Athens, Greece

**AIM:** To review current data on issues pertaining to micrognathia as a risk factor involved in the aetiology of developing obstructive sleep apnoea (OSA), in subjects presenting craniofacial syndromes.

**MATERIALS AND METHOD:** Characteristic clinical cases from the files of the Clinic of Sleep Apnoea of the Dental School, University of Athens and a private orthodontic practice of Athens. For all patients a minimum two lateral cephalometric radiographs were available, ranging from their initial admission to the clinic to early adulthood.

**RESULTS:** A severely hypoplastic/micrognathic mandible and/or maxilla is typically observed in subjects presenting craniofacial syndromes such as Pierre-Robin, Down, Apert, Crouzon, Treacher-Collins and Parrott. In such cases, the development of clinically

significant signs and symptoms of OSA is strongly associated with the severity of the micrognathia of the jaw(s).

**CONCLUSION:** Micrognathia is typically identified as an aetiological factor predisposing to OSA in patients with craniofacial syndromes.

## CP 088

Wednesday, June 16 - at 16:00

### POSTERIOR MAXILLARY OSTEOTOMY TO RE-ESTABLISH INTERMAXILLARY SPACE BEFORE PROSTHODONTICS

**A Aneiros Ardao**, University of Santiago de Compostela, Spain

**AIM:** Severely extruded maxillary molars can interfere with prosthodontic rehabilitation of the edentulous mandibular molar region. Previous treatment options have included restoration of the mandible in an abnormal occlusal plane by shortening the teeth; extensive foreshortening of the extruded antagonists, necessitating possible endodontics; periodontal crown-lengthening surgery and crowns; a combination of the two previous options; extracting the extruded teeth; or no treatment. Surgical treatment is an alternative for these situations. The posterior maxillary segmental osteotomy (PMSO) is a simple but precise technique to manage this problem. The purpose of this presentation is to show a case of PMSO for mandibular implant placement

**SUBJECT AND METHOD:** A 49-year-old female with severe dentoalveolar extrusion of the posterior maxillary region. As the maxillary teeth were nearly touching the mandibular gingival mucosa, a PMSO was performed to increase the freeway space. The treatment plan was developed after diagnostic casts were mounted in a semi-adjustable articulator. An acrylic splint was fabricated for use during and after surgery. The PMSO was performed under general anaesthesia. The mobilized segment was placed in the correct position using the splint. The segment was fixed with titanium miniplates. To maintain the placement of the superiorly positioned segment, the patient wore the splint before placement of the mandibular implants.

**CONCLUSION:** PMSO resulted in successful treatment for a situation demonstrating posterior maxillary segmental collapse.

## CP 089

Wednesday, June 16 - at 16:00

### INTERDISCIPLINARY TREATMENT OF MULTIPLE AGENESIS – A CASE REPORT

**A Peniche**, P Lara, Orthodontic Department, UNAM, DEPeI FO, México D.F

**AIM:** To show the treatment of a patient with multiple agenesis.

**SUBJECT AND METHOD:** A 35 year-old female patient with multiple tooth agenesis with occlusal implications such as instability, ectopic permanent maxillary canines, primary upper canines, increased overbite, extrusion of teeth 11, 21, multiple interdental spaces and several rotations. The agenesis was not related to any syndrome. The aim of treatment was to rehabilitate the patient functionally and aesthetically, to create sufficient space for the implants, to close unwanted spaces to achieve symmetry, and finally to correct the overbite, with the MBT 0.022 inch technique.

**RESULTS:** Favourable changes were noted in the occlusion including good alignment of the teeth and reduction of overjet and overbite. This was achieved by intrusion of the upper incisors and extrusion of the mandibular molars.

**CONCLUSION:** Treatment resulted in an improvement in occlusal stability and aesthetics.

## CP 090

Wednesday, June 16 - at 16:45

### HOW TO TORQUE A PALATALLY IMPACTED CANINE

**M Araghbidi Kashani**, Department of Orthodontics, School of Dentistry, Shahed University, Tehran, Iran

Orthodontic treatment of palatally impacted maxillary canine teeth is usually time-consuming and difficult. The distant location of the palatally impacted canine tooth from the main archwire

makes it difficult to exert the desired force system to the impacted tooth. In most cases the tooth can be brought into the line of occlusion only by tipping movement, but as a result of new imaging tools that provide a better view of the relationship of the roots of the impacted tooth and neighbouring teeth, in a few cases the roots of the impacted maxillary canine are in such a situation that the root of the impacted tooth must be moved before it can be pulled into its correct position in the dental arch. The aim of this presentation is to introduce a torquing lasso that can torque the impacted tooth so that its root is moved away from the neighbouring tooth.

## CP 091

Wednesday, June 16 - at 16:45

### FORCED ERUPTION – INTRODUCING AN EASY METHOD

H R Pakshir, A Karamifar, H R Fattahi, M Davari, **S Torkan**,  
Orthodontic Research Center, Shiraz University of Medical  
Sciences, Iran

**AIM:** Many clinical situations exist in which a tooth fractured down to the cemento-enamel junction is observed which at first appears cannot be saved without a prosthetic bridge or implant treatment. The aim of this presentation is to introduce a simple appliance that can be used for forced eruption.

**MATERIALS AND METHOD:** A silicone impression of the area is taken and poured. After endodontic treatment, a space is prepared in the canal and a post made using 0.036 inch stainless steel wire is cemented to the prepared post space. After pouring the impression, two teeth adjacent to the fractured area on either side should be surveyed to define the height of their contours. The designed appliance, which has a horizontal bar and two circumferential clasps and is interconnected between at least two adjacent teeth on either side, is then waxed-up and moulded. An elastic chain is applied to the post to extrude the teeth. For determination of the extent of tooth extrusion a periapical radiograph is taken every 2 weeks. In the present study this appliance was used for 15 teeth (eight upper lateral and seven upper central incisors).

**RESULT:** Since the appliance takes its retention from the adjacent teeth and because the reactive force of eruption is in a gingival direction, usually there is no dislodgement. The mean extrusion observed was  $2.7 \pm 0.3$  mm on the fractured tooth and  $0.2 \pm 0.08$  mm on the adjacent teeth.

**CONCLUSION:** This appliance can be used to save teeth that seem to be unrestorable. Considering the periodontal situation, the prognosis of teeth and patient motivation, forced eruption can be beneficial.

## CP 092

Wednesday, June 16 - at 16:45

### PRE-PROSTHODONTIC ORTHODONTIC TREATMENT IN IMPLANT SUPPORTED RESTORATIONS

Ş Karaçay<sup>1</sup> M Erkan<sup>1</sup>, **A Atay<sup>2</sup>**, <sup>1</sup>Department of Orthodontics, GMMMA Haydarpasha Education Hospital, İstanbul and <sup>2</sup>Department of Prosthodontics, GMMMA Center of Dental Sciences, Ankara, Turkey

**AIM:** Early loss of permanent teeth may cause various malocclusions due to migration of neighbouring teeth to the extraction side. In some cases, prior to implant supported prosthetic restorations, space for the lost tooth needs to be obtained. The aim of this presentation is to show the orthodontic treatment of two patients in whom space was required for implant supported restorations.

**SUBJECTS AND METHOD:** A 16-year-old female who had lost her right central incisor in childhood due to a trauma. Her right lateral incisor had migrated, completely occupying the space of the lost tooth. In the mandible, the canines were in a vestibular position and the incisors were misaligned because of the insufficient arch perimeter. The first premolars were extracted and the maxillary lateral incisor was repositioned, space for the missing upper maxillary central incisor was opened, the misaligned lower incisors were corrected, and a Class I molar relationship was achieved with

fixed orthodontic appliances. Implant supported prosthetic restoration was performed for the rehabilitation of the missing incisor. The second patient was a 20-year-old female whose mandibular first molar had been extracted 2 years previously. The second molar was tipped and mesialized due to early loss of the first molar was uprighted and distalized with orthodontic treatment. Prosthetic restoration was undertaken with two implants inserted in the region of the missing tooth.

**RESULTS:** Appropriate aesthetic and function was achieved in both patients using a multidisciplinary approach.

**CONCLUSION:** In patients with insufficient space for insertion of an implant, pre-prosthetic orthodontic treatment is required for satisfactory results.

## CP 093

Wednesday, June 16 - at 16:45

### CAMOUFLAGE TREATMENT OF A PATIENT WITH A CLASS II DIVISION 1 MALOCCLUSION – A CASE REPORT

**Z Stamenković**, K Brajović, S Milošević, V Dusić, V Filipović,  
Department of Orthodontics, University of Belgrade, Serbia

**AIM:** Presentation of a patient treated with upper and lower fixed appliances and extraction of the upper first premolars.

**SUBJECT AND METHOD:** A 28-year-old female patient. Before treatment the overjet was 10 mm and overbite 4 mm. Lateral cephalometric analysis showed severe protrusion of the upper incisors; angle I/SpP was 38°, while other relevant parameters were SNA 86°, SNB 81.5°, ANB 4.5°, SN/SpP 8°, SN/MP 25°, SpP/MP 17°, sum of angles of Björk's polygon  $\Sigma = 380^\circ$ , i/MP 81°, I/i 100°. The patient was treated for 23 months. During the last stage, elastic intermaxillary traction was used (heavy pull; 184 g; 4.8 mm;).

**RESULTS:** At the end of treatment the overjet was 3 mm and the overbite 4 mm. The following parameters were measured on the lateral cephalographs: SNA 84.5°, SNB 81°, ANB 3.5°, SN/SpP 6.5°, SN/MP 22.5°, SpP/MP 16°, sum of angles of Björk's polygon  $\Sigma = 382^\circ$ , i/MP 76°, I/i 121°. I/SpP was significantly increased  $-62.5^\circ$ . The occlusion was stable at the end of treatment, the overjet was significantly less and the profile and facial aesthetics were improved.

## CP 094

Wednesday, June 16 - at 16:45

### ORTHODONTIC AND SURGICAL TREATMENT OF IMPACTED MANDIBULAR SECOND PREMOLARS

**F Sali<sup>1</sup>**, B Mehmeti<sup>2</sup>, H Mehmeti<sup>1</sup>, J Gjorgova<sup>1</sup>, M Petruseska<sup>1</sup>,  
Departments of Orthodontics, <sup>1</sup>University Dental Clinical Centre, Skopje, Former Yugoslav Republic of Macedonia and <sup>2</sup>University of Prishtina, Kosovo

**AIM:** Impaction of the mandibular second premolars is mostly caused by their idiopathic rotation and position in the mandibular bone during growth and development. The teeth most often impacted, in order of frequency, are the maxillary and mandibular third molars, maxillary canines and mandibular second premolars. The aim of this study is to show the orthodontic surgical of a patient with impacted mandibular second premolars.

**SUBJECT AND METHOD:** A 13-year old boy with bilateral impaction of the mandibular second premolars in the permanent dentition. A clinical case determined by bilateral lack of second mandibular premolars and loss of first permanent molars was used. A panoramic radiograph showed bilateral impaction of the second mandibular premolars, with severe mesial inclination and lingual rotation. During surgical treatment, using the gingivectomy and corticotomy technique, brackets were bonded on the exposed buccal surface of the tooth crowns. Fixed orthodontic appliance treatment was undertaken using the straightwire technique.

**RESULT:** During one year of orthodontic treatment correct positioning of the two premolars in the dental arch was achieved.

**CONCLUSION:** Orthodontic treatment with surgical exposure is a successful procedure for positioning impacted teeth in the dental arch, and has a good long-term prognosis with no adverse pulpal or periodontal risk to the tooth.

**CP 095***Wednesday, June 16 - at 16:45***COMBINED ORTHODONTIC-IMPLANT TREATMENT OF CONGENITALLY MISSING LATERAL INCISORS****M Stojanović<sup>1</sup>**, Z Stajčić<sup>2</sup> <sup>1</sup>Private practice Paraćin and <sup>2</sup>Centre for Maxillofacial Surgery, 'Beograd-centar', Serbia

**AIM:** To undertake orthodontic treatment to create conditions for the construction of crowns on dental implants to replace congenitally missing lateral incisors in order to achieve integrity of the maxillary arch, harmonious dental arches and a pleasant facial appearance.

**SUBJECT AND METHOD:** A 25-year old patient with a traumatic occlusion, hypodontia of the lateral maxillary incisors, a large median diastema, divergent and abraded central incisors and a low fraenum. Treatment included a fraenectomy and orthodontic treatment to close the median diastema, and to create adequate intercoronary and inter-radicular space for future implant-prosthetic restoration of the missing incisors with ongoing dental arch matching, evaluation of the facial profile and correct positioning of the incisors. Two Nobel Biocare dental implants were inserted on each side and ceramic crowns on implants were constructed.

**RESULT:** Treatment resulted in the achievement of good occlusion, good dental and facial aesthetics and a pleasant smile.

**CONCLUSION:** Combined orthodontic-implant treatment proved to be a successful option for the treatment of missing maxillary lateral incisors in adult patient.

**CP 096***Wednesday, June 16 - at 16:45***HERBST APPLIANCE IN THE LATE TREATMENT OF CLASS II DIVISION 1 MALOCCLUSION – A CASE REPORT****N Nedeljković**, Z Stamenković, Department of Orthodontics, University of Belgrade, Serbia

**AIM:** To present the long-term follow-up of a subject treated with the Herbst and fixed appliances.

**SUBJECT AND METHOD:** A 15-year-old female at the beginning of the therapy beyond the ideal treatment period, i.e. before the pubertal peak of growth. A lateral cephalogram showed a skeletal Class II and dentoalveolar Class II division 1 malocclusion. She was treated with a Herbst treatment for 6 months followed by fixed appliances for 9 months. During Herbst treatment a rapid palatal expander was used. The measurements (Pancherz, 1982) were performed on lateral cephalograms before and after treatment: incisor inferior, incisor superior, molar inferior, molar superior, pogonion and subspinale. The distance from these points to Olp were measured and compared on the cephalograms and the temporomandibular joints on tomograms obtained pre- and post-treatment.

**RESULTS:** Correction (maxillary and mandibular) was found in the molar and incisor relationship, overjet and overbite. Sagittal skeletal changes and soft tissue profile improvement were observed. Superimposition of the pre- and post-treatment TMJ tomogram showed remodelling of the condyle and articular fossa.

**CONCLUSION:** The Herbst appliance is effective in the late treatment of Class II malocclusions.

**CP 097***Wednesday, June 16 - at 16:45***THE ORTHODONTIC MANAGEMENT OF CLEFT LIP AND PALATE PATIENTS****E Hussein**, Associate professor, Palestinian Territory

Children born with oral clefts require treatment from a number of health care providers during the first two decades of their life. These children acquire problems ranging from feeding difficulties as infants to frequent ear infections, compromised speech, aberrant dentofacial development, malocclusion, and last but not least esthetic impairment with its unfavorable effects on the psychological adaptation of the patient.

During the deciduous dentition stage of development, orthodontic treatment may be appropriate in some instances. Extensive orthodontic procedures are not usually done at this stage. During the mixed dentition stage, secondary alveolar bone grafting procedures are usually performed. It is generally accepted that the graft placement be preceded by a phase of orthodontic treatment involving the maxillary arch. Alveolar cleft closure affords the patient with a number of advantages, including stabilization (unification) of the maxilla, closure of oronasal communication, provision of an osseous base for the eruption of canines and other teeth, as well as improves esthetics.

This presentation will review the etiology and embryology of oral clefts, the timing of lip and palate closure, the acrylic feeding plate and the cleft palate orthopedics, and will concentrate on the orthodontic management of the cleft patients in all their different stages of development

**CP 098***Wednesday, June 16 - at 16:45***A MODIFIED ELN APPLIANCE IN OPEN BITE TREATMENT OF GROWING PATIENTS – A PRELIMINARY STUDY****F Tomassini**, A Di Lonardo, A Giancotti, University of Rome 'Tor Vergata' - Fatebenefratelli Hospital- Isola Tiberina, Italy

**AIM:** An anterior open bite (AOB) can be related to a vertical skeletal pattern or to deleterious habits (thumb sucking and tongue thrusting) or both. A tongue thrust can be seen as an effect of a skeletal open bite or as the main aetiological factor of the malocclusion. Treatment of functional open bite patients is usually carried out at 6 years of age, after the eruption of the incisors and in simple tongue thrusting habit the treatment is usually based on functional appliances that require high compliance to achieve acceptable results. The envelope linguale nocturne (ELN; Bonnet, 1992) is a functional activator based on an acrylic appliance with a 'tongue tunnel' to correct the tongue thrusting, which requires minimal compliance nighttime wear and 1 hour in the daytime. The aim of this presentation is to illustrate the efficacy of a modified ELN appliance in dental open bite subjects with a tongue thrusting habit.

**SUBJECTS AND METHOD:** Five patients with a Class I normodivergent skeletal pattern and AOB associated with a tongue thrust treated for a mean period of 1 year. Steiner's maxillary and mandibular incisor position, interincisal angle, overbite, overjet, mandibular and palatal planes related to SN were measured before and after treatment.

**RESULTS:** The AOB was successfully treated in all patients and upper and lower incisor proclination was reduced after a mean time of 7 months. The tongue thrust habit was successfully treated and the result was stable.

**CONCLUSION:** The modified ELN appliance seems to be an effective option in dental open bite and tongue thrust treatment.

**CP 099***Wednesday, June 16 - at 16:45***LONG-TERM RETENTION AFTER TREATMENT OF SKELETAL CLASS II GROWING PATIENTS WITH A DEEP BITE\*\*\*****J-H Kim**, Y-J Choi, C Chung, K-H Kim, Department of Orthodontics, Gangnam Severance Dental Hospital, College of Dentistry, Yonsei University, Seoul, South Korea

**AIM:** An anteroposterior (AP) discrepancy in skeletal Class II growing patients can be treated by maxillary growth inhibition and mandibular growth guidance. A deep bite can be resolved by regulating the vertical dimension of the anterior and posterior segments. The aim of this presentation is to show the successful treatment of skeletal Class II growing patients with a deep bite.

**SUBJECTS AND METHOD:** Three growing patients diagnosed as skeletal Class II treated by orthopaedic appliances during the growing phase, followed by fixed appliances with no extractions. Lateral cephalograms taken before, after, and at least 5 years after treatment were superimposed to evaluate skeletal growth and facial development.

**RESULTS:** 1. AP skeletal relationship was improved by regulating maxillary growth with high pull headgear. 2. Relative intrusion of the anterior teeth contributed to the deep bite correction. 3. The mandibular plane was maintained by vertical growth of the mandibular ramus. 4. Treatment results were successfully maintained during retention with fixed retainers.

**CONCLUSION:** Skeletal Class II growing patients with a deep bite can be successfully treated by dentofacial orthopaedic appliances which improve the vertical and horizontal skeletal relationship. Relative intrusion of the anterior teeth with vertical growth of the mandible seems to resolve the deep bite.

## CP 100

Wednesday, June 16 - at 16:45

### EPIDEMIOLOGY OF CLEFT LIP AND PALATE IN THE FORMER YUGOSLAV REPUBLIC OF MACEDONIA

**R Dimovska**<sup>1</sup>, S Naumovski<sup>2</sup>, J Stojanoski<sup>1</sup>, G Curcieva-Cuckova<sup>3</sup>, S Popovski<sup>3</sup>, <sup>1</sup>Private Dental Practice, Krana MS, <sup>2</sup>Clinic of Maxillofacial Surgery and <sup>3</sup>Faculty of Dentistry, Skopje, Former Yugoslav Republic of Macedonia

**AIM:** To retrospectively estimate the prevalence of a cleft lip and palate in the Former Yugoslav Republic of Macedonia, and variability related to form, gender and geographical area.

**MATERIAL AND METHOD:** Data were collected from the Clinic of Maxillofacial surgery in Skopje and the Republic Institute of Health protection. The material comprised cleft children born between 1996-2006. The data were obtained manually because of the lack of a cleft lip and palate team.

**RESULTS:** The estimated incidence of cleft lip and palate was 1:1000, which is quite low.

**CONCLUSION:** Due to the lack of records and the difficulty in manually searching the database, there is a need for the creation of a state centred archive and a national cleft team.

## CP 101

Wednesday, June 16 - at 16:45

### A SEVERE COMPLICATION OF BONDED MANDIBULAR LINGUAL RETAINERS

**P Pazer**, C Katsaros, Department of Orthodontics and Dentofacial Orthopedics, University of Bern, Switzerland

**AIM:** Flexible spiral wire retainers bonded to the lingual surfaces of all six anterior mandibular teeth are a common type of retention and are, in the majority of the patients, very efficient. In 3-5 per cent of subjects, however, they can cause unexpected movements of anterior teeth to such an extent that retreatment is necessary. The aim of this presentation is to show a major complication of a lingual retainer needing retreatment.

**SUBJECT AND METHOD:** A 20-year-old male with a broken retainer between 42 and 43. The retainer had been bonded four years prior to the incident. The root of tooth 43 was approximately 30 degrees buccally torqued. A 6 X 6 cm cone beam computed tomography (CBCT) was performed and analyzed. The root of tooth 43 was almost completely out of the bone on its buccal side. The nerve and vascular bundle had followed the apex to the limit of the bone, so that tooth vitality was preserved. Orthodontic retreatment was performed. Just before the end of retreatment a new CBCT was taken. Although the tooth has been moved back to its original position only the apical area of the root was covered with new bone whereas the rest of the root surface remained uncovered. Clinically slight gingival recession remained. After retreatment a new lingual retainer was bonded.

**RESULTS:** The successful retreatment allowed a satisfying aesthetic result for the patient. Because bone formation and mineralization is a slow process, the lack of bone on the buccal root aspect observed on the second CBCT could have been due to the short time interval after moving the tooth back to its original position.

**CONCLUSION:** Orthodontists, dentists and patients should be aware of possible complications of bonded retainers and instructed on how to detect them, if possible, at an early stage.

## CP 102

Wednesday, June 16 - at 17:30

### AN ALTERNATIVE METHOD FOR OVEREXTRUDED SECOND MOLARS IN ADULT PATIENTS

**J E Vlachojannis**, M Santoro, Columbia University School of Dental and Oral Medicine, New York, USA

Unopposed molar teeth are believed to be at risk for overeruption, impaired masticatory function, and the development of temporomandibular disorders; all problems that are frequently observed. Two approaches are currently used to correct overerupted/extruded posterior teeth: 1) the prosthodontic approach, which reduces the vertical height of the crown of the extruded tooth, and 2) the orthodontic approach. This report presents two cases successfully treated with a novel orthodontic approach for selective molar intrusion the modified palatal arch appliance (TPA). To intrude overerupted maxillary second molars a modified TPA was designed according to a calculated standard vector model. Of course, each clinical application would have variable effects due to the individual environment. The modified TPA represents a force system that is equivalent to a couple with a 30 g net force to move the tooth bodily. A short length of elastomeric chain from the helix to the palatal sheath of the maxillary second molar applied a 50 g force to the overerupted maxillary second molar. The elastomeric chain was replaced every three weeks. Sufficient intrusion of the maxillary second molar was obtained within two months. When intrusion of a single tooth is needed, the modified TPA is an attractive orthodontic alternative to traditional edgewise mechanics, temporary anchorage devices, or removable appliances. The traditional edgewise side-effect of intrusion of the adjacent tooth did not occur. Not all patients are candidates for insertion of temporary anchorage devices, either for psychological, medical, or anatomical reasons. The use of removable appliances also does not seem to be indicated when the terminal tooth requires intrusion. The modified TPA can be used successfully to intrude overerupted maxillary second molars and is a clinically elegant, non-invasive, and cost-effective procedure because it does not require laboratory work. This appliance can be realized in one chairside appointment.

## CP 103

Wednesday, June 16 - at 17:30

### TREATMENT OF A TOOTH WITH A COMPLICATED CROWN-ROOT FRACTURE – A CASE REPORT

**B Bajraktarova**<sup>1</sup>, E Bajraktarova Valjakova<sup>2</sup>, C Misevska<sup>3</sup>, T Miseveski<sup>2</sup>, Departments of <sup>1</sup>Pediatric Dentistry, <sup>2</sup>Prosthetics and <sup>3</sup>Orthodontics, Faculty of Dentistry, Ss. Cyril and Methodius University, Skopje, Former Yugoslav Republic of Macedonia

**AIM:** To show the treatment of a 9-year-old male patient who had injured his maxillary permanent right incisor.

**MATERIALS AND METHOD:** The diagnosis of a crown-root fracture of the incisor (open apex) was based on clinical and radiographic examination. The fracture line began about 2 mm incisal to the marginal gingiva on the labial aspect of the crown, ending a few millimetres below the gingival crevice orally. A partial (calcium hydroxide) pulpectomy was performed before coronal fragment removal. To ensure a good seal, zinc oxide eugenol was placed over the calcium hydroxide, and finally glass-ionomer cement. The tooth was followed at 6, 12 and 21 months, and 6 years after the trauma. During that time, the tooth continued to erupt and completed root development (21 month). At the same time, the apical fragment spontaneously erupted, so that the subgingival margin extruded sufficiently to the position where restoration was possible. After definitive endodontic treatment and root canal filling, the tooth was restored with post-retained acrylic crown. Tooth space was maintained by orthodontic treatment.

**CONCLUSION:** Teeth with incomplete root formation and crown-root fracture may be maintained by endodontic treatment, performed towards apexogenesis. The apical fragment in immature teeth can spontaneously erupt or, if necessary, after root completion, orthodontically extruded, and then restored.

**DISTORTED SHAPE AND SIZE OF TEETH – A CHALLENGE FOR CONTEMPORARY ORTHODONTIC TREATMENT**

**A Sieja**, J Antoszewska, Department of Dentofacial Orthopedics and Orthodontics Medical University in Wrocław, Poland

**AIM:** Facial aesthetics are considered to be socially significant. Contemporary orthodontics is also involved since with correction of dental and skeletal relationships, patients achieve proportional and regular facial features as well as harmonious smiles. However when the natural shape and size of the patient's teeth are distorted, orthodontic treatment alone is insufficient. In such cases aesthetic dentistry may be necessary. Reconstruction of the malformed teeth with composite material is a low-invasive method of regaining normal tooth-anatomy.

**SUBJECTS AND METHOD:** A 30-year-old male patient treated with lingual appliances due to high aesthetic demands. The patient's smile after completion of orthodontic treatment remained unattractive due to the distorted shape and size of the teeth. The upper incisors were restored with composite (GC Gradia Direct, GC Corporation, Japan). This reconstruction resulted in a regular smile-line and diminished diastema. The second patient was a 17-year-old female under treatment with an aesthetic appliance; her main concern was an impacted central left incisor. Presently only 4 mm of the tooth has emerged into the oral cavity. In order to camouflage the anaesthetic appearance, the incisor crown was artificially elongated with composite material (GC Gradia Direct); subsequently an orthodontic bracket was bonded on the new incisal surface to control tooth extrusion.

**CONCLUSION:** Patients may obtain harmonious and attractive smiles even if their teeth are malformed, not only by prosthetic restoration veneers or crowns. Constantly improved composite materials used in cosmetic dentistry enables low-invasive support of orthodontic treatment results.

## CP 105

Wednesday, June 16 - at 17:30

**INTERDISCIPLINARY TREATMENT: ORTHODONTIC EXTRUSION WITH MINI-IMPLANTS**

**A Peniche**<sup>1</sup>, R Medellín<sup>1</sup>, H Ballado<sup>2</sup>, <sup>1</sup>Orthodontic and <sup>2</sup>Dental Prosthesis and Implantology Departments, UNAM, DEPEL FO, México D.F

**AIM:** To show the treatment of a patient with periodontal problems and a low bone level.

**SUBJECT AND METHOD:** A 52-year-old male with periodontal problems and a low bone level, such that it was not possible to insert implants. In order to rehabilitate the patient with multiple implants, keeping the margin at a level compatible with gingival health, the central and lateral incisor teeth were extruded using orthodontic mini-implants. Before extrusion, root canal treatment was carried out. A casted pin and post were then cemented into the root and it was prepared for a head-cured acrylic resin temporary crown. The crown was set with glass ionomer luting agent to prevent debonding during traction.

**RESULTS:** Favourable changes were noted in the anterior bone level with extrusion of the teeth (about 3 mm), so the implants could be inserted in correct place.

**CONCLUSION:** Following the improvement in bone level, the treatment goals were exceeded with the help of orthodontic mini-implants for extrusion.

## CP 106

Thursday, June 17 - at 16:45

**RELATIONAL DEVELOPMENT IN BABIES WITH CLEFT LIP AND PALATE: A STUDY CARRIED OUT IN FRANCE**

**B Grollemund**<sup>1,2</sup>, V Lesnes<sup>3</sup>, W Bacon<sup>1</sup>, <sup>1</sup>Functional Unit of Dentofacial Orthopaedics, University Hospital, <sup>2</sup>Laboratoire d'Ethique et Pratiques Médicales, Université de Strasbourg and <sup>3</sup>Service de chirurgie plastique et maxillo-faciale, Centre Hospitalier Universitaire de Grenoble, France

**AIMS:** The study of patients with a cleft lip(s), with or without a cleft palate (CL/P) during the repetitive appointments required by orthodontic treatment shows the importance of the psychological setting and of the parent-child relationships. The aims of this prospective, multidisciplinary and multicentre research were to assess the psychological perceptions of parents of children affected by a CL/P in the year following the birth, and to analyse the child's degree of psychological suffering as well as the parent-child relationship.

**SUBJECTS AND METHOD:** Four centres that intervene at different times according to the waiting times between the child's birth and the first surgical intervention. The results obtained for each group will be cross-compared according to two sub-groups of parents who found out about the CL/P through embryo diagnosis and those who found out at their child's birth. The underlying hypothesis is that the longer the period prior to the first intervention the more likely the parents' psychological perception will affect the parent-child relationship and influence their child's harmonious development. The mental and psychological dimension related to the malformation and its correction will be analysed in the parental group (importance of the pre-natal diagnosis, relational development with the child, self-esteem, quality of life, etc.) but also in the child (likely distress and withdrawal symptoms).

**RESULTS:** The findings will enable the development of new methods to lessen the psychological impact of the malformation on parents and improve the children's treatment in the long term. Every care centre should consider the mental health of these children and their parents by offering psychological support from an early age.

## CP 107

Thursday, June 17 - at 16:45

**USING CASTILLO MORALES STIMULATION PLATE FOR A PATIENT WITH APERT SYNDROME – A CASE REPORT**

**A Ifert**<sup>1</sup>, R Sader<sup>2</sup>, S Kopp<sup>1</sup>, Department of <sup>1</sup>Orthodontics and <sup>2</sup>Oral, Maxillofacial and Plastic Surgery, Goethe University Frankfurt, Germany

**AIM:** Apert Syndrome is a genetic defect and belongs to the craniofacial anomalies. It occurs in 1 per 160,000 to 200,000 live births. It is characterized by specific malformations of the skull, the midface, the hands, and the feet. Newborns with Apert Syndrome suffer from low oxygen saturation. The patient has a median cleft palatal. The aim of this research was to provide higher oxygen saturation for growth of the baby.

**MATERIALS AND METHOD:** The upper jaw with a median cleft palate is very narrow. The patients tongue has a non-physiological function. The tongue stays at the palate and the oxygenic saturations drop down. To bring the tongue in a physiological position a Castillo Morales stimulation plate was used. A stimulation ring is situated in the area of the palatal folds. The ring shape was changed at every clinic visit.

**RESULTS:** During use of the stimulation plate it was possible to reduce the decreases in oxygen saturation. The patient learned a physiological position and function of the tongue. Home monitoring was still necessary.

**CONCLUSION:** The Castillo Morales stimulation plate is a useful aid for patients with dysfunctions of the tongue. It is important to change the shape of the stimulation ring, and home monitoring is important. Studies with larger groups of patients are necessary.

## CP 108

Thursday, June 17 - at 16:45

**LONG-TERM RETENTION AFTER TREATMENT OF A DEEP BITE IN NON-GROWING PATIENTS**

**J-H Kim**, H-R Yoon, S-S An, K-H Kim, Department of Orthodontics, Gangnam Severance Dental Hospital, College of Dentistry, Yonsei University, Seoul, South Korea

**AIM:** To show successful camouflage treatment by vertically regulating the anterior and posterior segments of non-growing patients with a deep bite.

**SUBJECTS AND METHOD:** Three non-growing patients with a deep bite treated with fixed orthodontic appliances. The anterior teeth were intruded with a three-piece intrusion arch and the results were maintained with a fixed lingual retainer. Lateral cephalograms taken before, after, and at least 5 years after treatment were superimposed in order to evaluate skeletal and dental changes.

**RESULTS:** 1. The deep bite was resolved by absolute intrusion of the maxillary anterior teeth. 2. As there were only very minor changes in the vertical position of the anterior teeth, the results were successfully maintained during the retention period.

**CONCLUSION:** Non-growing patients with a deep bite can be successfully treated with camouflage treatment by improving the horizontal and vertical dental relationships with absolute intrusion of the anterior teeth.

## CP 109

Thursday, June 17 - at 16:45

### TREATMENT OF CLASS II DIVISION 1 MALOCCLUSION, THROUGH COMBINATION OF FIXED AND MOBILE APPLIANCE

**E Çitaku**, A Reshitaj, E Lupçi, E Çitaku, Private Practice 'Magic Dent', Prishtina, Kosova

**AIM:** To show the results of treatment of Class II division 1 anomalies through a combination of fixed and mobile appliances.

**SUBJECT AND METHOD:** A 19 year old, female in the permanent dentition, open frontal bite, an overjet of 10 mm, anterior maxillary protrusion, Class II division 1 malocclusion and interocclusal space in the anterior region of 5 mm. Since the patient had infantile swallowing, a removable appliance was used for six months to eliminate the habit. The maxillary first premolars were then extracted and whilst continuing with treatment of the habit, distalisation of maxillary canines was started since a diastema was present between the maxillary incisors. During the second phase retrusion of the maxilla was commenced followed by fixed appliances to level the teeth.

**RESULTS:** Two year's post-treatment the occlusion was stable.

**CONCLUSION:** Knowledge of removable and fixed devices and their combination gives good results in the treatment of orthodontic anomalies.

## CP 110

Thursday, June 17 - at 16:45

### GERM EXTRACTION – A THERAPEUTIC APPROACH FOR DENTOFACIAL ORTHOPAEDICS USING MINI-IMPLANTS

**H Sarabia**, Universität Hamburg, Germany and Universidad de Chihuahua, Mexico

Germ extraction before the beginning of maximum growth potential is the most favourable therapeutic approach in order to prevent crowding.

Indications A) Class I malocclusion due to bimaxillary protrusion combined with mini-orthoimplants, lack of maxillary basal bone or Pont index 35. B) Class II due to total maxillary protrusion combined with mini-orthoimplants. C) Class III due to total mandibular protrusion (Progenie). D) Open bite due to Pont index 35. E) Deep bite no clinical considerations found.

The alternative to using temporary mini-implants during development and growth gives the opportunity to eliminate total maxillary and bimaxillary protrusions in Class I and II combined with germ-extraction therapy in order to change kids facial profile before the normal eruption time.

## CP 111

Thursday, June 17 - at 16:45

### SIMULTANEOUS FUNCTIONAL MAGNETIC RESONANCE IMAGING – ELECTROMYOGRAPHIC MEASUREMENT DURING TOOTH TAPPING

**J Miyamoto**<sup>1,2,3</sup>, T Hanakawa<sup>3,4</sup>, T Ono<sup>1,2</sup>, T Kurabayashi<sup>1</sup>, K Moriyama<sup>1,2</sup>, <sup>1</sup>Tokyo Medical and Dental University, <sup>2</sup>Global Center of Excellence Program, International Research Center for Molecular Science in Tooth and Bone Diseases, <sup>3</sup>National Center of Neurology and Psychiatry, Precursory Research for Embryonic Science and Technology, <sup>4</sup>Japan Science and Technology Agency, Japan

**AIM:** Recently, several non-invasive techniques for measuring human brain functions have been introduced and applied to examine brain mechanisms of physiological movement. Above all, functional magnetic resonance imaging (fMRI) is a modality with high spatial and temporal resolution, which is an essential prerequisite for precise measurement of brain activity. To evaluate brain functions during movement, it is also important to measure muscle activity quantitatively during fMRI. For example, in some studies that investigated brain activity during finger movement, a simultaneous measurement of fMRI and surface electromyography (EMG) was performed. However, there are no such studies on the brain controlling mechanisms of jaw movement. The aim of this study was to develop a simultaneous fMRI-EMG technique during tooth tapping and to verify the data.

**SUBJECTS AND METHOD:** Healthy volunteers performed a tooth-tapping task with different tapping forces during fMRI scanning and EMG recording from the masseter and temporalis muscles with MR compatible electrodes. Functional imaging data were analyzed to identify the brain areas with task-related signal changes, and EMG data were full-wave rectified and then integrated to quantify muscle activity after eliminating imaging artefacts.

**RESULTS:** Brain activity related to tooth tapping after removing head-motion artefacts associated with jaw movement could be identified. Additionally, differences in muscle activity were quantitatively shown for different force conditions.

**CONCLUSION:** Reliable data were obtained by simultaneous fMRI-EMG measurement during tooth tapping. It is thus suggested that the technique may be useful to investigate the possible correlation between masticatory muscle activity and brain functions controlling tooth tapping.

## CP 112

Thursday, June 17 - at 16:45

### RETENTION: A CASE PRESENTATION FOR THE POSTGRADUATE COURSE AT TOKYO DENTAL COLLEGE

T Nezu, **M Mukai**, M Kobayashi, K Sueishi, Department of Orthodontics, Tokyo Dental College, Japan

**AIM:** In postgraduate education, there is a tendency to emphasize the importance of diagnostics and therapeutics. However, the concept of long-term management also needs to be fully understood, taking into account such factors as relapse and stability of occlusion. The objective of this study was to evaluate patients with an overjet of +5 mm prior to treatment. Pre- and post-operative treatment records were compared.

**SUBJECTS AND METHOD:** Eight Japanese patients (4 males, 4 females with retention time spans ranging from 2 years ~ 4 years 8 months. All presented for initial treatment with an Angle Class II classification. Five patients were treated with, and three without extractions. All wore circumferential type maxillary retainers. With respect to the mandible, three wore circumferential type retainers, one wore both circumferential and fixed retainers, three wore fixed type retainers, while one wore a clip type retainer.

**RESULTS:** Post-treatment relapse was as follows: five presented with maxillary anterior buccal inclination, four with mandibular lingual inclination with slight crowding, and one with relapse to a Class II Angle classification. The overjet was maintained in two patients after bracket removal, five had an increase, while one presented with a decrease. Maxillary anterior buccal inclination relapse can be attributed to tongue thrust, bruxism and mouth-breathing. The correction of deleterious habits is of importance for good long-term retention.

**CONCLUSION:** Long-term stability after orthodontic treatment is an important clinical goal. Observing and comparing the various stages of retention can improve diagnosis and treatment.

**MANDIBULAR SYMPHYSIS ADVANCEMENT WITH TOOTH- AND BONE-BORNE DISTRACTION DEVICES**

T Türk<sup>1</sup>, F Çakmak<sup>2</sup>, M Sumer<sup>3</sup>, Departments of <sup>1</sup>Orthodontics and <sup>3</sup>Oral Surgery, University of Ondokuz Mayıs, Samsun and <sup>2</sup>Department of Orthodontics, University of Zonguldak Karaelmas, Turkey

**AIM:** To evaluate the effects of mandibular symphysis advancement with distraction osteogenesis (DO) on the dentofacial structures.

**SUBJECTS AND METHOD:** Advancement of the mandibular symphysis with DO was carried out in four female patients aged between 12 years 1 month and 18 years 10 months. The osteotomy line, between the canine and the first premolar, was extended from the interdental area to the lower border of the mandibular symphysis on both sides under general anaesthesia. Subsequently the mandibular symphysis was mobilized with the osteotome by making small rotational movements at the osteotomy lines. In all cases, individual tooth- and bone-borne distraction devices were used. After a latency period of 1 week the screws were activated (0.8 mm/day). The distraction device was removed under local anaesthesia after an 8 week consolidation period. Dentofacial changes were evaluated on pre-distraction and post-consolidation cephalograms.

**RESULTS:** Increases of SNB and condylion-gnathion ranged from 2 to 3 degrees and from 0.5 to 5.5 mm, respectively. In three subjects an increase of anterior face height was observed. This increase ranged from 2 to 3 mm. With anterior movement of the mandibular symphysis, ANB angle decreased. This decrease ranged from 1 to 2.5 degrees. A decrease of overjet was noted of between 3 to 6 mm. A 5.5 to 13 mm increase of mandibular arch length was calculated for all patients.

**CONCLUSION:** Mandibular symphysis advancement is an effective method for the treatment of increased overjet, lower anterior crowding, and improvement of the soft tissue profile.

**CONGENITAL ABNORMALITIES OF THE EXTERNAL EAR IN CLINICAL ORTHODONTIC EXAMINATION**

N Vasilakou, P Synodinos, M Papagrigorakis, Department of Orthodontics, School of Dentistry, University of Athens, Greece

**AIM:** The external ear is a complex structure comprising the auricle, the cartilaginous external auditory canal, and the bony external canal. Its physiological function includes the collection, filtering and direction of acoustic signals from the environment to the tympanic membrane. Morphologically normal external ears come in a wide variety of sizes and shapes. This variation is attributed mainly to racial and familial characteristics. Congenital abnormalities of the external ear are not uncommon, affecting 1 in 12,500 newborns. Anomalous morphology of the external ears is most often evident at delivery. The aim of this study was to review the embryology and development of the external ear and to investigate the aetiology of its most commonly observed morphological anomalies.

**SUBJECTS:** Characteristic clinical cases associated with syndromic and non-syndromic abnormalities of the external ear.

**RESULTS:** Dysmorphic signs or abnormalities are diagnosed when the external ears are large/long, small/microtic, malformed, deformed, low-set, prominent/outstanding, abnormally orientated, bilaterally asymmetrical and, in extreme cases, vestigial or totally absent. Though most cases are isolated and sporadic, they may be part of several genetic syndromes, such as trisomy 21, 18 or 22, DiGeorge sequence, Treacher Collins, Robinow, foetal alcohol, and Goldenhar syndromes.

**CONCLUSION:** Given the intricate pattern of external ear development that may be affected by several genetic and/or environmental factors, attention should be given during clinical examination of the external ear.

**PREFORMED DEVICES: TEN-YEAR'S EXPERIENCE**

D Tarantino, G Ierardo, A Guarino, F Petrillo, A Polimeni, Department of Oral Science, Sapienza University of Rome, Italy

**AIM:** During recent years preformed devices have been played an important role in the ortho-paedodontic field. These devices allow management of orthodontic interceptive therapy, combining efficacy with comfort, ease of use and a reduction in treatment time.

**MATERIALS AND METHOD:** Over ten years of experience using these devices in orthodontic interceptive therapy at U.O.C. of Pediatric Dentistry of 'Sapienza', University of Rome will be explained.

**RESULTS:** The process has allowed optimisation of therapeutic interventions regarding efficacy and efficiency. Ease of use is one advantage that allows diversification of timing and goals. Using technology for materials allows treatment special need patients and for those with a relative contraindication to the use traditional device. Last, but not least, are the economic and social aspects of these preformed devices.

**CONCLUSION:** Preformed devices are now completely absorbed in orthodontic interceptive therapy as well as traditional devices.

**ORODENTAL, CRANIOFACIAL AND GENERAL CHARACTERISTICS OF LACRIMO-AURICULO-DENTO-DIGITAL SYNDROME**

D A Papamanou, P Christou, Department of Orthodontics, University of Athens, Greece

**AIM:** Lacrimo-auriculo-dento-digital (LADD) syndrome, also known as Levy-Hollister syndrome, is a rare syndrome (1:2000 live births) characterized by pronounced dysplasia in various organs. It is attributed to a mutation in the gene encoding fibroblast growth factor 10, which is also key factor in the regulation of craniofacial growth. The clinical characteristics are mainly structural deficiencies of lacrimal and salivary tracts, small cup-shaped and/or malformed ears, sensorineuronal or conductive hearing loss, and variable anomalies of the hands and feet. Although the degree to which the different organs are involved varies considerably, there is no clear report of the dental and craniofacial anomalies. The aim of this presentation is to describe the orodental, craniofacial and general characteristics of a patient with LADD syndrome.

**MATERIALS AND METHOD:** Medical, dental history and orthodontic records of a 9-year-old girl, and analysis of two previously reported cases.

**RESULTS:** Clinical examination revealed a full permanent dentition, with no abnormalities in the shape of the teeth, an Angle Class I malocclusion, a hypoplastic chin and zygomatic arches, an anterior open bite and convex profile. Lateral cephalometric analysis showed a retrognathic mandible and hyperdivergence. The patient presented cup-shaped/low-set ears with bilateral hearing loss and dysplasia of the fingers and toes. Nasolacrimal duct obstruction, xerostomia or agenesis of salivary glands is variably reported. The patient, in addition to the typical manifestations of the syndrome, had a palatal cleft at birth that was operated on during the first six months, and agenesis of the thumbs.

**CONCLUSION:** Although LADD syndrome is a developmental disorder it apparently involves pathways that, to a variable degree, affect craniofacial growth.

**THE EFFECTS OF THE PENDEX APPLIANCE IN A CLASS II DIVISION 2 MALOCCLUSION SUBJECT**

T Moreira<sup>1</sup>, A Ângelo<sup>2</sup>, A L Maló<sup>3</sup>, M Gomes<sup>3</sup>, <sup>1</sup>Private Practice, Oporto, <sup>2</sup>Private Practice, Coimbra, and <sup>3</sup>Department of Orthodontics, Faculty of Medicine, University of Coimbra, Portugal

**AIM:** To describe the treatment effects of the Pendex of Hilgers appliance in a patient with a Class II division 2 malocclusion. This

type of mechanotherapy that requires no patient cooperation is a common treatment modality for subjects with maxillary skeletal and/or dentoalveolar protrusion.

**SUBJECT AND METHOD:** An adult female with a Class II division 2 malocclusion treated with the Pendex appliance to distalize the upper first molars. Cephalometric evaluation was undertaken and the sagittal relationship was determined.

**CONCLUSION:** The Pendex is an effective method of treatment for Class II malocclusions.

#### CP 118

Thursday, June 17 - at 17:30

##### A DILACERATED UPPER INCISOR IN A YOUNG PATIENT – ORTHODONTIC TREATMENT

**S Tronca**, L Giannin, G Galbiati, F Iazzetti, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** Dilacerated teeth are dental elements presenting an abnormal angle between the crown and root, usually following childhood trauma. This pathology may alter normal tooth development and eruption and result in poor aesthetics. The aim of this presentation is to show the treatment of a young patient with an upper dilacerated incisor.

**SUBJECT AND METHOD:** A 5-year-old male in good health. The parents reported trauma to the upper primary incisors during childhood that had resulted in an alteration in the shape of the permanent upper central incisor.

**RESULTS:** Orthodontic-surgical treatment of the permanent upper central dilacerated incisor resulted in its correct position in the dental arch.

**CONCLUSION:** Tooth and periodontal conditions at the end of treatment were good and there was no tooth mobility. Aesthetic and functional results were optimal and they have remained stable.

#### CP 119

Thursday, June 17 - at 17:30

##### RELATIONSHIP BETWEEN RAPID PALATAL EXPANSION AND SAGITTAL DISCREPANCIES

**F Iazzetti**, L Giannini, G Galbiati, S Tronca, C Maspero, Department of Orthodontics, University of Milan, Italy

**AIM:** Rapid palatal expansion (RPE) in growing patients often has effects on skeletal growth and the sagittal position of the maxillary and mandibular bones. The aim of this presentation is to show the cephalometric results after RPE.

**SUBJECTS AND METHOD:** Seventy-five patients of both genders, aged 5 to 13 years with transverse maxillary hypoplasia. Twenty-five had a skeletal Class I, 24 a skeletal Class II and 26 a skeletal Class III malocclusion. Cephalometric analysis was undertaken on lateral telerradiographs obtained before and after expansion. SNA, SNB, ANB, SN-SNPSNA and SN-GoMe angles were evaluated.

**RESULTS:** An improvement in ANB angle was observed in all patients which improved in the Class III subjects (+2.15) and diminished in the Class II patients (-1.53). Minimal change was observed in the Class I group (-0.18]. In the Class II patients the maxilla advanced inconsistently and mandible advanced constantly. The average improvement in SNA and SNB angles was 0.19 and 1.68 degrees, respectively.

**CONCLUSION:** With RPE it is possible to obtain an improvement in the maxillary transverse diameter and consequent advantages for the sagittal plane.

#### CP 120

Thursday, June 17 - at 17:30

##### ORTHODONTIC TREATMENT OF INCOMPLETE TRANSPOSITION IN THE LATE MIXED DENTITION

**Ş Karaçay**<sup>1</sup>, F Sabuncuoğlu<sup>2</sup>, M Erkan<sup>1</sup>, Departments of Orthodontics, <sup>1</sup>GMMA Haydarpasha Education Hospital İstanbul and <sup>2</sup>GMMA Center of Dental Sciences, Ankara, Turkey

**AIM:** Transposition is a relatively rare dental anomaly, characterized by an interchange in the position of two adjacent permanent

teeth on the same side of the dental arch. Incomplete transposition is a condition describing an interchange in the positions of the crowns of two permanent teeth within the same quadrant of the dental arch, while the root apices remain in their relative positions. Complete transposition is a similar situation in which both the crowns and the entire root structure are transposed. The aim of this presentation is to show the orthodontic treatment of incomplete unilateral transposition of the mandibular canine and lateral incisor during the late mixed dentition.

**SUBJECT AND METHOD:** A 12 year-old female with incomplete transposition of the mandibular left canine and lateral incisor. The panoramic radiograph revealed that the apex of mandibular left lateral was placed on the mesial corner of the crown of the mandibular left canine while the crown was erupted ectopically near the mandibular left second molar. Following uprighting of the lateral incisor with a removable appliance, a utility arch was used up to completion of the permanent dentition. After eruption of the permanent teeth, the treatment continued with full fixed appliances.

**RESULTS:** At the end of the orthodontic treatment lasting 26 months, the transposition of the canine and lateral incisor were corrected, and a Class I occlusion with a favourable smile was obtained.

**CONCLUSION:** Malocclusions with transposition or severely ectopic eruption of the teeth may be treated orthodontically. In these cases early diagnosis affects the prognosis and treatment procedures.

#### CP 121

Thursday, June 17 - at 17:30

##### SEGMENTAL MANDIBULAR DENTOALVEOLAR DISTRACTION: AN OPTION TO EXTRACTION THERAPY

**G Sgritta**, **D Esercizio**<sup>1</sup>, R De Santis, S Nucci, L Perillo, <sup>1</sup>Department of Orthodontics, <sup>2</sup>University of Naples, <sup>3</sup>maxillo-facial surgeon, <sup>3</sup>specialist in orthodontics and odonstomatology, Italy

**AIM:** In the craniofacial region, osteodistraction can be applied to patients with facial anomalies to lengthen either one or both jaws, or a single piece of bone such as a segmental distraction of the anterior dentoalveolar process. This procedure includes four steps: pre-surgical orthodontics, banding of the distractor, osteotomy and movement of the alveolar process, and tooth movement into the created space. The aim of this presentation is to show that osteodistraction is a reliable therapy for patients with skeletal or dentoalveolar discrepancies.

**MATERIALS AND METHOD:** A lingual arch with a two screw-type expander mechanism was bonded on the first molars and lower incisors. Activation started one week after the osteotomy and ended when the required space was gained. After two months, the teeth were moved into the new bone.

**RESULTS:** Four adult patients with dento-skeletal Class II malocclusions were successfully treated with osteodistraction. The required space corrected lower crowding of 6 mm in one patient; Class II molar and incisor crowding associated with a biretrusive profile in another; allowed treatment of a Class II division 2 in a patient with a retroalveolar process and previous mandibular trauma; and improved severe crowding and overjet in the last patient.

**CONCLUSION:** In appropriate cases, segmental osteodistraction can offer a quick biological advantage and good aesthetic results.



# INDEX OF AUTHORS

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Aartman, I	SP 058	72					Ateş, M	OP 24	25				
Ábalos Labruzzi, C	SP 012	60	SP 029	64			Athanasidou, A	SP 217	114	SP 254	123		
Abou Jamra, B	OP 57	42					Attin, T	SP 195	108				
Abou Jamra, R	OP 57	42					Austin, D F	SP 060	72				
Abrahamsson, C	OP 50	38					Avventi, T	OP 37	31	SP 130	91	SP 131	91
Acar, A	OP 24	25					Aydemir, H	SP 142	94				
Ackermans, M	OP 14	19					Aydin Mustafa, F	SP 280	129				
Adeyemi, A	OP 32	29					Aydogdu, E	OP 33	29				
Adlam, D	OP 36	31					Baccetti, T	HAO 04	5	OP 35	30	OP 39	32
Agache, F	CP 013	149						OP 42	34	SP 048	69	SP 050	70
Aghaeipour, N	SP 139	93					Bachleda-Księdzularz, M	SP 035	66				
Aguilar, F	CP 082	166					Baciut, G	SP 112	86				
Ahmeti, B	CP 017	150					Bacon, W	CP 106	171	CP 064	161	OP 60	43
Ahn, H W	SP 064	73						SP 270	127				
Ahn, S H	SP 116	87	SP 117	88			Baek, S H	SP 064	73				
Akcam, M O	SP 173	102					Baharoğlu, E	CP 041	156				
Aksoy Doğan, A	CP 85	166					Bahat Yalvac, A	SP 280	129				
Albreht Zore, M	SP 300	135					Bajraktarova Valjakova, E	CP 103	170	SP 250	122		
Alcan, T	OP 40	33	SP 059	72			Bajraktarova, B	CP 103	170				
Algera, T	SP 188	106					Balcioglu, A H	SP 176	103				
Algera, T J	SP 225	115					Ballado, H	CP 105	171				
Alikhani, M	SP 037	66					Ballanti, F	SP 214	117	SP 045	68		
Alkiş, H	CP 085	184	SP 176	103	SP 095	81	Balon-Perin, A	CP 045	153	CP 049	158	CP 055	159
Almasan, O	SP 112	86						SP 078	77				
Almeida, M	SP 076	77					Balos Tuncer, B	SP 289	132				
Al-Moammar, K	SP 257	124					Bantleon, H P	SP 103	83	SP 202	110	SP 275	128
Almoqubel, A	SP 021	62						SP 298	134	SP 089	80		
Al-Nawas, B	CP 011	149					Bantleon, J	SP 298	134				
Alqerban, A	OP 18	22					Barbato, E	CP 083	166	SP 180	104	SP 077	77
Altug, A T	OP 30	28	SP 271	127				SP 094	81				
Alzahawi, K	OP 51	39					Barbosa, H	SP 076	77				
Amasyali, M	SP 320	140					Barbosa, H A	SP 274	128				
Amfiloxiou, A	CP 087	167					Barekzai, J	CP 038	155				
An, S S	CP 108	171	SP 226	116			Barreto, L	CP 028	153				
Anderson, I M	SP 109	86					Barrilà, S	SP 174	102				
Andjelić, J	CP 078	165	SP 168	101			Bartley, N	SP 277	103				
Andrić, M	CP 020	151					Bartzela, T	SP 054	71				
Aneiros Ardao, A	CP 088	167					Battelino, T	SP 282	130				
Ângelo, A	CP 117	173					Bavbek Karakaya, I	SP 279	129				
Anikiyenko, A	SP 159	99					Baykul, T	SP 095	81				
Annousaki, O	SP 302	135					Baytorun, E	CP 084	166				
Annousaki-Thormann, O	SP 209	112					Bazargani, F	SP 002	57				
Antonarakis, G	HAP 01	6					Bebnowski, D	OP 13	19				
Antonini, G	CP 025	152	CP 026	152			Beerens, M W	SP 186	106	SP 187	106		
Antonini, M	HAO 03	7					Behets, C	SP 093	81				
Antoszewska, J	CP 104	171	CP 015	150	CP 077	165	Behnaz, M	SP 124	90				
	SP 040	67					Behr, M	SP 227	116	SP 228	116		
Apicella, D	SP 148	95					Bekir, S	CP 065	162				
Apruzzese, I	SP 090	80					Bencze, A	CP 018	151	SP 129	91		
Araghbidi Kashani, M	CP 090	167					Bencze, A M	CP 030	153				
Aranda, L	SP 169	101					Bencze, M A	SP 123	89				
Arhouni, K	SP 078	77					Bengi, O	CP 048	158				
Arko Kampus, K	CP 074	164					Bergé, S	OP 27	26				
Arponen, H	CP 080	165					Bertl, M	SP 275	128				
Arrigoni, P	SP 174	102					Berto, C	OP 37	31	SP 130	91	SP 131	91
Arsova, E	CP 065	162						SP 132	91	SP 133	92		
Assumma, V	SP 148	95					Beuran, I	SP 084	79				
Atac, M S	OP 30	28					Bigliazzi, R	SP 233	118	SP 239	119	SP 076	77
Atamozlu, E K	CP 044	157					Bille, M-L B	SP 223	115				
Ataoglu, T	SP 181	104					Bister, D	SP 177	103	SP 178	103		
Atay, A	CP 092	168					Bjerklin, K	SP 081	78				

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Blaga, J	SP 146	95					Cheremisova, V	SP 258	124	SP 259	124		
Bocchino, T	SP 184	105					Cheveresan, A	CP 007	148				
Bock, N	SP 055	71					Chew, M T	SP 157	98				
Bodini, A	SP 261	125					Chin, A	OP 08	16	SP 192	107		
Bøe, O E	OP 51	39	SP 152	97	SP 235	118	Cho, J-H	SP 322	140				
Bøe, O E B	SP 236	118					Choi, I R	CP 054	159				
Bolamperti, L	SP 209	112	SP 026	63			Choi, J Y	SP 064	73				
Bolender, Y	SP 169	101					Choi, S H	SP 319	140				
Bonadeo, S	CP 022	151	CP 004	147	SP 306	136	Choi, Y J	CP 099	169	SP 226	116		
	SP 307	137					Chourova, I	SP 219	114				
Bondemark, L	OP 50	38	SP 190	107	SP 281	130	Christou, P	CP 116	173	CP 069	163	SP 096	82
	SP 081	78					Chung, C	CP 099	169	SP 226	116		
Bongaarts, C	OP 54	40					Cilenšek, M	SP 282	130	SP 283	130		
Bonhof, L S	SP 187	106					Cioffi, I	HAO 01	4				
Booij, J W	SP 253	122					Circieva-Cuckova, G	SP 119	88				
Bordanzi, A	SP 306	136	SP 307	137			Çitaku, E	CP 109	172	CP 109	172		
Borsos, G	OP 07	16					Colak, C	OP 16	20	SP 277	129	SP 301	135
Bortun, C	CP 007	148	CP 008	148			Colić, S	CP 020	151				
Borzabadi-Farahani, A	HAP 02	7	SP 001	57			Consoli, G	SP 285	131				
Borzabadi-Farahani, A	HAP 02	7	SP 001	57			Contardo, L	SP 241	120	SP 243	120		
Boskovic-Brkanovic, T	SP 170	101					Cooke, M	OP 36	31				
Boucher, C	CP 045	157					Cordasco, G	HAO 01	4				
Bourauel, C	OP 52	39	SP 221	114	SP 292	133	Corega, C	SP 260	124				
	SP 294	133	SP 296	134	SP 297	134	Cornelis, M	OP 41	33	SP 093	81		
	SP 134	92					Coskun, M	OP 33	29				
Božič, M	SP 289	132					Cotič, J	SP 199	109				
Bozkurt, S	CP 068	162	SP 099	82			Cozza, P	SP 047	69				
Božnik, V	SP 137	93					Crisan, M	CP 060	161				
Bracco, A A	CP 033	154	SP 137	93	SP 138	93	Cristache, C	SP 165	100				
Bracco, P	CP 093	168					Csiki, I	SP 120	88				
Brajović, K	CP 093	168					Cubas, P	OP 02	13				
Brandt, T	SP 003	57					Cubrilo, D	SP 249	121				
Bratu, D C	CP 013	149	SP 071	75			Culea, L	SP 071	75				
Bratu, E	SP 071	75	SP 083	78	SP 092	81	Curcieva-Cuckova, G	CP 100	170	SP 242	120	SP 252	122
Breuning, H	OP 19	22						SP 269	127	SP 087	79		
Bronkhorst, E	SP 054	71					Cutrera, A	SP 077	77				
Brudvik, P	OP 51	39	SP 236	118			Cuvelier, C	OP 01	13				
Brusveen, E M G	SP 236	118					D'Alatri, L	CP 025	152				
Buck, I	SP 003	57	SP 065	74			D'Antò, V	OP 09	17				
Bucur, A	SP 088	80					Dagistanli, S	SP 105	85				
Bugaighis, I	SP 007	58					Dagsuyu, I M	SP 105	85				
Buhl, J	OP 15	20					Dalessandri, D	CP 022	151	CP 033	154	CP 004	147
Bumann, A	SP 074	76						SP 306	136	SP 307	137		
Büntemeyer, B	SP 194	108					Dalessandri, M	SP 306	136				
Burllibasa, M	SP 164	100	SP 165	100			Dalstra, M	OP 15	20				
Burnik, S	SP 104	84					Darendeliler, M A	OP 16	20	OP 35	30	SP 277	129
Buschang, P H	OP 48	37						SP 301	135				
Busscher, H J	SP 032	65					Darendeliler, N	SP 279	129				
Cacucci, L	CP 006	148					Darija, M	SP 141	94				
Çakır, E	SP 113	87					Davari, M	CP 091	168				
Çakırer, B	SP 041	67					David, S N	SP 274	128				
Çakmak, F	CP 113	173					De Baets, E	SP 256	123				
Cangialosi, T	OP 46	36					De Clerck, H	OP 42	34				
Cannavale, R	SP 318	139					De Gregorio, F	SP 291	132				
Cappabianca, S	SP 314	138					de Mezer, M	SP 267	101				
Caprioglio, A	SP 135	92	SP 026	63			De Pauw, G	OP 01	13	OP 53	40	SP 238	119
Cara, P	SP 033	65						SP 070	75				
Castaldo, A	SP 243	120					De Santis, R	SP 310	137	CP 121	174		
Castaldo, M I	SP 318	139					De Somere, S	SP 070	75				
Cattaneo, P	SP 302	135					Debernardi, C	SP 137	93				
Çehreli, S B	SP 208	111					DeClerck, H	OP 41	33				
Celis, J P	SP 183	105					Del Giglio, R	SP 184	105				
Çerekja, E	CP 034	65	SP 041	67			Del Torre, M	OP 23	24				
Cernescu, A	SP 030	64					Deleurant, Y	OP 20	23				
Cerović, R	CP 076	77					Deli, R	CP 025	152	CP 026	152	CP 006	148
Ceschi, M	SP 243	120						SP 237	119				
Cesta, A	OP 37	31	SP 130	91	SP 131	91	Deljo, E	SP 245	120				
	SP 132	91	SP 133	92			Demir, P	SP 136	92				
Cevidanis, L	OP 41	33	OP 42	34			Demirer, S	SP 262	125				
Cha, J Y	SP 319	140	SP 039	67			Demirkaya, A A	CP 034	154				
Chai, L	HAP 04	8	SP 192	107			Derelli, E	SP 271	127				
Chan, Y H	SP 167	101					Dermaut, L	OP 01	13				
Charezinski, M	CP 049	158	CP 055	159	SP 213	113	Deschepper, E	SP 238	119				
Chartres, I	SP 037	66					Desoi, D	SP 009	149				
Chen, C C	CP 081	166					Dhopatkar, A	HAO 02	4				
Chenin, D	OP 03	14											

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Di Carlo, G	SP 201	109					Favero, L	CP 071	163	SP 293	133		
Di Gioia, E	SP 237	119					Fayed, M	SP 185	105				
Di Lenarda, R	SP 241	120					Fekonja, A	CP 042	156	CP 052	159		
Di Lonardo, A	CP 098	169					Ferrari, M	SP 174	102				
Di Lorenzo, M	CP 010	149	SP 234	118			Ferro, F	SP 148	95				
Di Maria, E	SP 243	120					Fieuws, S	OP 18	22				
Di Raffaele, S	SP 210	88					Filipović, G	CP 056	160				
Didilescu, A C	CP 030	153	CP 031	153	SP 305	136	Filipović, V	CP 093	168				
Dimberg, L	SP 190	107					Filleul, M P	SP 169	101				
Dimovska, R	CP 100	170	CP 016	150	SP 242	120	Finotti, M	OP 23	24				
	SP 087	79					Flahiardekani, F	SP 139	93				
Dinca, O	SP 088	80					Fleser Szuhaneck, C, T	CP 008	148				
Diya, L	SP 256	123					Fleser, C	CP 007	148				
Djordjević, J	OP 21	23					Fontana, P	CP 022	151				
Dominiak, M	SP 082	78					Foong, K W C	SP 011	59				
Dorobat, V	SP 027	64					Foong, W C K	SP 167	101				
Dostalova, T	SP 031	65					Franchi, L	OP 39	32	SP 162	99	SP 047	69
Dragan, M	SP 110	86						SP 048	69	SP 050	70		
Dragomir, B	SP 073	76					Freudenthaler, J	SP 298	134				
Drerup, I	CP 023	62	CP 046	69			Freyss, C	CP 067	162				
Drevenšek, A	OP 38	32					Frezzato, I	SP 293	133				
Drevenšek, G	SP 268	131					Frongia, G	SP 137	93	SP 138	93		
Drevenšek, M	CP 019	151	CP 037	155	CP 073	163	Frumusa, E	SP 135	92				
	SP 104	84	SP 268	123	SP 282	130	Fudalej, P	OP 54	40	SP 110	86		
	SP 283	130					Funaki, J	SP 043	68	SP 044	68		
Drobyshev, A	SP 066	74					Funaki, S	SP 043	68	SP 044	68		
Drobysheva, N	SP 122	89	SP 066	74			Fusco, R	SP 310	137				
Droschl, H	SP 171	102					Gabrieli, P	SP 241	120				
Dudkiewicz, Z	OP 54	40					Galan, E	CP 018	151				
Dumu Dundu Jr., E	CP 045	157					Galantucci, L M	SP 237	119				
Durka-Zajac, M	SP 155	98	SP 072	76			Galbiati, G	CP 118	174	CP 119	174	SP 308	137
Dushenkova, M	SP 159	99						SP 309	137	SP 311	138	SP 312	138
Dusić, V	CP 093	168						SP 313	138	SP 315	139		
Efstratiadis, S	OP 46	36					Galluccio, G	CP 083	166	SP 180	104		
Eftene, O	SP 084	79					Ganea, AD	CP 031	153				
Eghiazaryan, A	SP 166	100					Gao, H	SP 052	70				
Ehmer, U	CP 039	156					Gašpirc, B	CP 029	153				
Eigon, A	CP 068	162					Gedrange, T	OP 55	41	OP 07	16	SP 246	121
Ekberg, E C	OP 50	38						OP 247	121	SP 304	136	SP 082	78
Ekestubbe, A	SP 061	73					Ghafari, J	OP 12	18	OP 49	37		
El, H	OP 03	14					Ghorbani, A	SP 124	90				
El-Bialy, T	OP 52	39	SP 294	133			Giancotti, A	CP 021	151	CP 066	162	CP 079	165
Eleftheriadi, I	CP 086	167	CP 087	167				CP 098	169	SP 261	125	SP 067	74
Elekdag-Turk, S	OP 16	20	SP 277	129	SP 301	135		SP 090	80				
Eliades, T	SP 221	114	SP 254	123	SP 292	133	Giannin, L	CP 118	174				
Elmi, P	SP 068	75					Giannini, L	CP 119	174	SP 308	137	SP 309	137
Emani, Z	SP 118	88						SP 311	138	SP 312	138	SP 313	138
Emeljanova, O	SP 085	79	SP 086	79				SP 315	139				
Emral, M E	SP 173	102					Gil, J	SP 014	60				
Enache, A M	CP 032	154	SP 305	136			Gilkens, C	SP 078	77				
Engel, U	SP 063	73					Gioeva, Y	SP 085	79	SP 086	79		
Erbe, C	SP 264	125	SP 284	131			Giordano, C I	SP 318	139				
Erkan, M	CP 047	157	CP 048	158	CP 092	168	Giordano, D	OP 11	18				
	CP 120	174					Giot-Wirgot, S	CP 055	159				
Erken, S	SP 288	132					Giuntini, V	SP 162	99	SP 048	69	SP 049	70
Ersoy, U	OP 46	36					Gjethammer, M R	SP 152	97				
Ertas, H	SP 317	139					Gjorgova, J	CP 094	168	SP 206	111	SP 250	122
Esenlik, E	SP 038	66					Gkantidis, N	SP 096	82				
Esercizio, D	CP 121	174					Glavan Jianu, R, F	CP 005	147				
Eslamipour, F	SP 001	57					Glavan, F	CP 007	148	CP 008	148	SP 030	64
Esposito, V	SP 314	138						SP 083	78	SP 092	81		
Evcil, E H	SP 038	66					Glineur, R	CP 045	157	CP 049	158	CP 055	159
Fabbrizi, M	SP 285	131						SP 213	113	SP 078	77		
Fabré, M	CP 069	163					Glišić, B	OP 21	23	SP 249	121		
Færovig, E	OP 51	39					Gokalp, H	SP 115	87	SP 136	92	SP 280	104
Falkensammer, F	SP 202	110						SP 057	72				
Faltermeier, A	SP 227	115	SP 228	116	SP 299	135	Golland, D	SP 158	98				
	SP 005	58	SP 006	58			Golland, L	SP 195	108				
Faltin Jr., K	SP 233	118	SP 239	119	SP 274	128	Golovinova, N	SP 085	79	SP 086	79		
Faltin, K	SP 162	99					Gomes, M	CP 117	173				
Faltina, K	SP 076	77					González Nieto, B	SP 004	58				
Farčnik, F	CP 062	161	CP 063	161	OP 38	32	González Nieto, E	SP 004	58				
	SP 147	95	SP 300	135			González Sanz, A M	SP 004	58				
Fattahi, H R	CP 091	168	SP 106	85	SP 160	99	Görgülü, G	SP 136	92				
Faur, N	SP 030	64					Götz, W	OP 57	42	SP 297	134		

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Goyenc, YB	CP 044	157					Ionescu, E	CP 030	153	CP 031	153	CP 060	160
Grammatopoulos, E	HAO 02	4						SP 101	83	SP 123	89	SP 128	90
Grassia, V	SP 310	137						SP 129	91				
Grbic, J	OP 46	36					Ionescu, I	SP 164	100	SP 165	100		
Greco, F	CP 006	148					Isaza Penco, S P	CP 010	149				
Greco, M	CP 021	151	CP 066	162	CP 079	165	İşcan, D	SP 113	87				
	OP 11	18	SP 261	125	SP 067	74	Iscan, D	SP 289	132				
	SP 090	80					İşcan, H N	SP 113	87				
Gredes, T	OP 55	41	SP 246	121	SP 247	121	Iscan, H N	SP 289	132				
	SP 304	136					Ismani, A	CP 017	150	CP 057	160	CP 058	160
Grigore, R	SP 128	90	SP 129	91			Jacobs, R	OP 18	22				
Grollemund, B	CP 106	171	CP 064	161	OP 60	43	Jacquet, W	SP 089	80				
	SP 270	127					Jäger, A	OP 52	39	OP 57	42	SP 292	133
	OP 56	41						SP 294	133	SP 296	134	SP 297	134
Gruenheid, T	SP 182	105					Jakupi, L	CP 065	162				
Grünheid, T	CP 115	173					Jamilian, A	SP 118	88	SP 124	90	SP 139	93
Guarino, A	SP 287	131	SP 095	81				SP 203	110				
Gungor, A Y	SP 105	85					Jämsä, M	SP 251	122				
Gurbuz, T	OP 40	33					Janega, M	SP 031	65				
Guzel, M Z	SP 068	75					Janev, R	CP 065	162				
Habets, L	OP 49	37					Janošević, M	CP 056	160				
Haddad, R	SP 257	124					Janošević, P	SP 126	90				
Haers, P	SP 019	61					Jaroni, A	OP 29	27				
Hägg, U	SP 181	104					Jeremiah, H	OP 36	31				
Hakki, S	SP 253	122					Jianu, R	SP 030	64	SP 092	81		
Halazonetis, D	SP 120	88					Jiménez Planas, A	SP 029	64				
Hamada, M	SP 056	71					Jo, YM	SP 039	67				
Hamza, E	CP 111	172					Jobbágy-Óvári, G	SP 025	63				
Hanakawa, T	OP 13	19					Jolevski, L	SP 269	101				
Hänggi, M	OP 22	24					Joss, C	HAO 03	5				
Hansen, K	CP 027	152					Jost-Brinkmann, P G	SP 036	66				
Haralabakis, O N	OP 24	25					Juloski, J	SP 249	121				
Haralambidis, A	SP 150	96					Jung, Y H	SP 191	107				
Hary, D	OP 25	25					Jurca, A M	SP 193	107				
Harzer, W	SP 276	129	SP 278	129			Jurišin, S	CP 072	163				
Hasegawa, Y	CP 023	152					Kahl-Nieke, B	CP 038	155	OP 43	34	SP 286	131
Heckmann, K	SP 140	93	SP 145	95			Kamínek, M	SP 248	121	SP 075	76		
Hedayati, Z	SP 013	60					Kan, H	CP 043	157				
Hemmatpour, S	OP 59	43					Kania, A	SP 230	117				
Henneman, S	OP 50	38					Karaçay, Ş	CP 047	157	CP 048	158	CP 092	168
Henrikson, T	OP 58	42						CP 120	174				
Herlin, T	SP 179	104					Karagöz, H	CP 047	157				
Hewson, A R	OP 32	29	SP 008	59			Karamifar, A	CP 091	168	SP 196	108	SP 197	108
Higham, S M	SP 042	67						SP 203	110	SP 244	120		
Hirvonen, O	SP 177	103					Karamifar, A A	SP 160	99				
Hj Abd Wahab, S	SP 007	58					Karasu, H A	OP 30	28	SP 142	94		
Hobson, R	OP 26	26					Kärkkäinen, S	SP 251	122				
Hocevar-Boltezar, I	SP 031	65					Karton, E	SP 166	100				
Hofmanová, P	SP 053	71					Katsaros, C	CP 101	170	OP 10	17	OP 54	40
Hong, J S	SP 284	131						SP 185	105	SP 253	122	SP 054	71
Hornikel, S	SP 183	105					Kau, C H	SP 134	92				
Hoste, S	OP 34	30					Kawala, B	CP 077	165	SP 040	67	SP 082	78
Huggare, J	OP 04	14					Kecsed, S	SP 193	107				
Hunt, N	CP 097	169					Kędzia, A	SP 211	112				
Hussein, E	SP 319	140					Keilig, L	SP 292	133	SP 296	134		
Hwang, C J	SP 322	140					Kern, W	SP 171	102				
Hwang, H-S	SP 027	64					Khaje, F	SP 127	90	SP 244	120		
Iacob, M	SP 067	74					Khoo, E	SP 037	66				
Iacopini, L	SP 112	86					Kiliaridis, S	HAO 03	5	HAP 01	6	CP 069	163
Iancu, M	SP 314	138						SP 316	139				
Iaselli, F	CP 118	174	CP 119	174	SP 308	137	Kim, C S	SP 111	86				
Iazzetti, F	SP 309	137	SP 311	138	SP 312	138	Kim, I G	CP 003	147				
	SP 313	138	SP 315	139			Kim, J H	CP 108	171	CP 099	169		
Ibel, G	SP 033	65	SP 089	80			Kim, K H	CP 108	171	CP 099	169	SP 226	116
Ierardo, G	CP 115	173	SP 201	109			Kim, Y	SP 111	86				
Ifert, A	CP 107	171	SP 266	126			Kim, Y J	SP 053	71				
Ifert, F	SP 266	126					Kirovska, T	SP 242	120				
Ihan Hren, N	CP 072	163	SP 134	92			Kirovski, I	CP 016	150	SP 242	120	SP 087	79
Ilhan, I	OP 40	33	SP 059	72			Kjær, I	OP 17	21	SP 198	109	SP 223	115
Imfeld, T	SP 195	108						SP 063	73				
Inglezos, E	CP 086	167					Kjær, KW	SP 063	73				
Innami, H	SP 020	62					Kjellberg, H	HAP 01	6	OP 28	27	SP 163	100
Inoue-Arai, M S	CP 014	150					Kleverlaan, C J	SP 188	106	SP 225	115		
Iodice, G	SP 184	105					Klimova, T	SP 023	62				
							Klinke, T	SP 246	121				

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Klukowska, M	SP 264	125					Lennartsson, B	SP 190	107				
Klytynskiy, Y A	SP 015	60					Leonardi, R	OP 11	18	SP 077	77	SP 094	81
Kobayashi, M	CP 112	172					Lesnes, V	CP 106	171				
Kocadereli, İ	CP 041	156	CP 084	166			Levrini, L	SP 026	63				
Kocoğlu Altan, A B	SP 262	125					Lewis, M	OP 04	14				
Kocova, M	SP 250	122					Lim, I K F	SP 011	59				
Koizumi, N	SP 222	115					Lindskog Stokland, B	SP 061	73				
Koklu, A	CP 012	149					Linke, M	SP 143	94				
Kolesov, M	SP 022	62					Lione, R	SP 121	89	SP 045	68	SP 047	69
Kolokithas, G	SP 217	114					Lisson, J	CP 023	152	CP 046	157		
Komarov, G	OP 32	29					Liu, Y	SP 037	66				
Kook, Y A	SP 111	86					Livas, C	SP 253	122				
Kopač, I	SP 097	82	SP 098	82			Löblein, S	CP 011	149				
Kopp, S	CP 107	171	SP 010	59	SP 194	108	Lomakina, V	SP 258	124	SP 259	124		
	SP 266	126	SP 003	57	SP 065	74	Lombardi, C C	SP 224	115				
	SP 009	59					Löppönen, H	SP 091	80				
Korbmacher, H	SP 286	131					Lossdörfer, S	OP 57	42				
Korfage, J A M	SP 182	105					Loster, B W	SP 035	66				
Kornblit, R	SP 201	109					Loster, B W	SP 125	90				
Korunoska-Stevkovska, V	SP 119	88					Louryan, S	SP 213	113				
Koselj, M	SP 282	130	SP 283	130			Lubinska, K	SP 155	98	SP 072	76		
Kosem, R	SP 104	84					Lucchi, P	SP 135	92	SP 144	94		
Kotova, M	SP 207	111					Lucke, S	SP 247	121				
Kotthaus, M	CP 038	155					Luigi, A	OP 09	17				
Kovacevska, G	SP 252	122					Luk, D	SP 272	128				
Kovalenko, A	SP 122	89	SP 066	74			Lund, H	OP 22	24				
Kowalska, E	SP 265	126					Lupçi, E	CP 109	86				
Koyuturk, A E	SP 320	140					Luther, F	SP 290	132				
Kozanecka, A	CP 015	150					Luzzi, V	SP 285	131				
Kramberger, D	CP 052	159	CP 070	163			Lyakhov, D	SP 022	62	SP 023	62		
Kristensen, K D	OP 58	42					Maal, T	OP 19	22				
Krneta, B	SP 100	83	SP 141	94			Macari, A	OP 12	18				
Krysiński, Z	SP 267	126					Macchi, A	SP 026	63				
Küçükkeleş, N	OP 24	25					Machata, W	CP 059	160				
Kuijpers-Jagtman, A M	HAO 03	5	OP 19	22	OP 27	26	Maddalone, M	SP 174	102				
	OP 54	40	SP 054	71			Mady Maričić, B	CP 076	164	SP 149	95		
	SP 058	72					Mahmoudi, F	SP 200	109				
Kuitert, R	OP 45	35	SP 231	117	SP 232	117	Mahony, D	CP 040	156				
Kuitert, RB	OP 55	41	SP 246	121	SP 247	121	Maiorana, F	OP 11	18				
Kunert-Keil, C	SP 304	136					Makou, M	SP 221	114				
	CP 111	172					Malas, M A	SP 038	66				
Kurabayashi, T	SP 066	74					Malinar, D	SP 100	83				
Kurakin, K	SP 295	133					Malinger, M	CP 063	161				
Kurojedova, V	OP 28	27	SP 163	100	SP 002	57	Malkiewicz, E	SP 212	112				
Kuroł, J	OP 30	28					Malkiewicz, K	SP 212	112	SP 216	113		
Kurt, G	OP 58	42					Malkoc, S	SP 062	73				
Küseler, A	SP 056	71					Maló, A L	CP 117	173				
KWong, R W	SP 274	128					Maltha, J C	OP 59	43				
Ladislau, A S	CP 022	151	CP 033	154	CP 004	147	Maneva, M	SP 252	122	SP 269	127		
Laffranchi, L	SP 307	137					Manzo, P	HAO 01	4				
	SP 215	113	SP 224	115			Marc, J	SP 268	126				
Laganà, G	CP 036	155	CP 037	155	CP 062	161	Marek, I	SP 248	121	SP 075	76		
Lah Kravanja, S	SP 205	110					Marino, A	SP 224	115				
Lahti, S	SP 184	105					Marion, A	CP 075	164				
Laino, A	CP 051	158	CP 070	163			Marion, L	SP 097	82				
Lajh, M	SP 256	123					Markic, G	OP 13	19				
Lambrechts, H	OP 56	41					Markova, M	SP 079	77				
Langenbach, G	SP 182	105					Marolt Mušič, M	OP 61	44				
Langenbach, G E J	SP 248	121	SP 075	76			Marrocco, S	CP 066	162				
Langova, K	CP 089	167					Marshall, D	OP 06	15				
Lara, P	OP 01	13					Martina, R	HAO 01	4	OP 09	17	SP 184	105
Laureys, W	SP 209	112					Martínez de Aragón Lancha, R	SP 004	58				
Laursen, M G	SP 108	85					Maru, N	SP 305	136				
Lee, A	SP 117	88					Maselli, A	CP 021	151				
Lee, C S	SP 107	85	SP 191	107			Masotto, G	SP 229	116				
Lee, J H	SP 053	71					Masoudi, A	SP 203	110				
Lee, K H	SP 039	67					Maspero, C	CP 118	174	CP 119	174	SP 308	137
Lee, K J	SP 322	140						SP 309	137	SP 311	138	SP 312	138
Lee, K-M	SP 191	107						SP 313	138	SP 315	139		
Lee, S M	SP 116	87						OP 39	32	SP 049	70		
Lee, Y S	SP 054	71					Masucci, C	CP 064	161	CP 067	162	OP 60	43
Leenarts, C	SP 149	95					Matern, O	SP 270	127				
Legović, M	SP 002	57						CP 067	162				
Leissner, L	SP 256	123					Mathis, R	CP 078	165	SP 321	140		
Lemiere, J	SP 166	100					Matijević, S	SP 024	63				
Lendengolts, Z							Matthews-Brzozowska, T						

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Mattick, C	SP 007	58					Neitzert, S V	OP 16	20				
Mavili, E	CP 041	156					Nemes, B	SP 025	63				
Mavragani, M	OP 51	39	SP 236	118			Nespriadko, T	SP 015	60				
Mazurkiewicz-Janik, M	SP 035	66					Nespryadko, V	SP 015	60				
M-Bille, LB	SP 223	115					Nevéus, T	SP 002	57				
McDonald, F	SP 257	124					Newton, T	SP 257	124				
McNamara, C M	SP 179	104					Nezu, T	CP 112	172				
Medellín, R	CP 105	171					Ng, H P	SP 167	101				
Meh, A	SP 268	126	SP 282	130	SP 283	130	Ng, I W	CP 014	150				
Mehmeti, B	CP 057	160	CP 094	168			Nguyen, E	OP 02	13				
Mehmeti, H	CP 094	168					Nguyen, T	OP 41	33	OP 42	34		
Mei, L	SP 032	65					Nicolescu, I	CP 032	154				
Melink, S	CP 062	161	OP 26	26	OP 61	44	Nieminen, P	SP 091	80				
Melsen, B	OP 15	20	SP 302	135			Nieto Sanchez, I	SP 161	99				
Merzhvinskaya, E	SP 066	74					Nijkamp, P	SP 068	75				
Meyer, P	SP 036	66					Nikolić, P	CP 020	151				
Midtbø, M	SP 152	97	SP 235	118			Nikolin, D	OP 03	14				
Migliore, M	SP 077	77					Nimigean, V R	SP 305	136				
Miguez Contreras, M	SP 161	99					Nitka, M	SP 074	76				
Mikac Cankar, M	SP 151	97					Nizankowska-Jędrzejczyk, A	SP 230	117				
Milačić, M	CP 072	163					Noens, V	OP 53	40				
Milicescu Jr., S	CP 018	151	CP 060	160			Nolting, D	SP 063	73				
Milicescu, V M	CP 018	151	CP 060	160	SP 123	89	Nordås, K	SP 235	118				
Million, A	SP 009	59					Nori, M	SP 139	93				
Milosavljević, Z	CP 020	151					Notaristefano, C	CP 026	152	CP 006	148		
Milošević, S	CP 093	168					Novakov, B	CP 050	158				
Milutinovic, J	SP 218	114					Novakov, S	CP 050	158				
Mina, K	SP 051	70					Noverraz, R	OP 27	26				
Miotti, F A	OP 23	24	OP 37	31	SP 130	91	Novruzov, Z	CP 012	149	SP 189	106		
	SP 131	91	SP 132	91	SP 133	92	Nucci, S	SP 148	95	CP 121	174		
	SP 229	116					Nucera, R	HAO 01	4				
Mirra, C	SP 285	131					Numata, M	SP 020	62				
Mischak, I	SP 202	110					O'Brien, K D	SP 172	102				
Miseveski, T	CP 103	170					O'Connor, D J	SP 016	61				
Misevska, C	CP 103	170	SP 206	111	SP 250	122	Odinokova, T	SP 240	119				
Miyamoto, J	CP 111	172					Oduncuoglu, B F	OP 33	29				
Mlee, S	SP 107	85					Ogodescu, A	CP 013	149	SP 073	76		
Mocuta, D	SP 164	100					Ohlendorf, D	SP 010	59	SP 194	108	SP 009	59
Mohammadi, A	SP 200	109					Okçu, K M	CP 048	158				
Moise, M	SP 071	75					Oliverio, A	CP 026	152				
Mojsoska, A	SP 269	127					Ong, S H	SP 167	101				
Momeni Danaei, S	SP 127	90	SP 196	108	SP 244	120	Ongkosuwito, E M	OP 48	37				
Mommaerts, N	SP 070	75					Ono, T	CP 111	172	CP 014	150		
Montasser, M	OP 52	39	SP 294	133			Orajärvi, M	SP 042	67				
Moreira, T	CP 117	173					Orellana, M	OP 02	13	SP 150	96		
Moriyama, K	CP 111	172					Ortolani, C F	SP 274	128				
Moriyama, K	CP 014	150					Oshagh, M	SP 196	108	SP 197	108		
Mossaz, C	CP 069	163					Ovsenik, M	CP 029	153	CP 036	155	CP 062	161
Mostafa, Y	OP 27	26						CP 068	162	CP 070	163	CP 072	163
Moullas, A	SP 217	114						CP 075	164	OP 26	26	OP 61	44
Mozzicato, P	CP 079	165						SP 100	83	SP 102	83	SP 134	92
Mucedero, M	SP 121	89	SP 215	87				SP 141	94	SP 147	95	SP 151	97
Muchitsch, A P	SP 273	128						SP 199	109	SP 300	135	SP 097	82
Muchitsch, P	SP 171	102						SP 098	82	SP 099	82		
Mukai, M	CP 112	172					Oyake, R	SP 222	115				
Müller, L	OP 31	28					Özcan, S	SP 320	140				
Müller-Hartwich, R	SP 036	66					Ozdemir, B	SP 289	132				
Muntean, A	CP 005	147	CP 008	148	SP 083	78	Ozdiler, E	SP 271	127				
	SP 092	81					Ozel, B	CP 012	149				
Mušič, M	CP 029	153	CP 070	163	OP 26	26	Ozer, M	CP 009	148				
	SP 099	82					Ozkaya, S	OP 24	25				
Mustafa, L	CP 017	150	CP 058	160			Özsoy, S	SP 208	111	SP 288	132		
Myrda, A	SP 302	135					Öztürk, F	SP 262	125				
Nabiev, N	SP 022	62	SP 023	62			Ozturk, O	SP 176	103				
Nada, R	OP 27	26					Pacurar, M	SP 193	107	SP 260	124	SP 073	76
Nagai, M	CP 002	147					Padricelli, G	SP 318	139				
Nakhleh, A	CP 010	149	SP 234	118			Paganelli, C	CP 033	154				
Nalcaci, R	SP 262	125					Paik, I K	SP 116	87	SP 117	88		
Naoumova, J	OP 28	27	SP 163	100			Pajevic, T	SP 249	121				
Narimani, M	CP 001	147					Pakshir, H	SP 244	120				
Naseri, M M	SP 196	108	SP 197	108			Pakshir, H R	CP 091	168				
Nasiri, F	SP 106	85					Palazón Martínez, C	SP 012	60	SP 029	64		
Nastalek, P	SP 230	117					Palm, R	OP 28	27				
Naumovski, S	CP 100	170					Palomo, J M	OP 03	14				
Nedeljkovic, N	CP 050	158	CP 096	169	SP 218	114	Pandis, N	SP 221	114				

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Pankratova, N	SP 022	62	SP 023	62	SP 240	119	Preoteasa, CT	CP 030	153	CP 031	153	SP 123	89
Panuszka-Landowska, J	SP 034	65					Preoteasa, E	CP 030	153	CP 031	153		
Papadopoulos, M	SP 254	123					Primožič, J	CP 036	155	CP 063	161	OP 38	32
Papadopoulou, K	SP 292	133						SP 098	82				
Papagrigorakis, M	CP 114	173	CP 086	167	CP 087	167	Prisco, D	SP 310	137				
Papamanou, D A	CP 116	173	CP 027	152			Proczek, K	OP 04	14				
Park, JU	SP 111	86					Proff, P	OP 55	41	SP 227	116	SP 228	116
Park, YC	SP 039	67						SP 299	135	SP 005	58	SP 006	58
Park, YH	SP 053	71					Pseiner, B	SP 298	134				
Patcas, R	OP 31	28					Pusch, K	SP 010	59				
Paúl Escolano, A	SP 012	60	SP 029	64			Quaggio, A	SP 239	119				
Paulsson-Björnsson, L	SP 281	130					Quaggio, A M	SP 233	118				
Pavlovic, D	SP 304	136					Quaglia, P S G	SP 233	118				
Pavlović, R	CP 029	153					Rabie, A B M	HAP 04	8	HAP 05	8	OP 08	16
Pavoni, C	SP 214	113	SP 045	68				SP 109	86	SP 016	61	SP 017	61
Payer, M	SP 273	128						SP 192	107	SP 021	62	SP 056	71
Pazera, P	CP 101	170	SP 185	105			Rabie, B	SP 153	97	SP 154	97		
Pedersen, T K	OP 15	20	OP 58	42			Racek, J	SP 219	114				
Pedroso, L	SP 076	77					Radlanski, R J	SP 036	66				
Peltonmäki, T	OP 13	19	OP 20	23	OP 31	28	Raftowicz-Wójcik, K	CP 077	165				
Penco, SI	OP 234	118					Ramoglu, I	SP 062	73				
Pender, N	OP 32	29	SP 008	59			Ramoglu, S I	SP 317	139				
Pender, S M	OP 32	29					Randjelović, N	CP 050	158				
Pendov, H	SP 269	127					Rangel, F	OP 19	22				
Peniche, A	CP 105	171	CP 089	167			Rapin, C	SP 169	101				
Pennings, B	OP 47	36					Raucci, M G	OP 09	17				
Percoco, G	SP 237	119					Raustia, A	SP 018	61	SP 042	67		
Perillo, L	SP 148	95	SP 291	132	SP 310	137	Raymond, J L	CP 064	161	CP 067	162		
	SP 314	138	SP 318	139	CP 121	174	Rehacek, A	SP 031	65				
Perinetti, G	HAO 04	5	SP 241	120	SP 243	120	Reicheneder, C	SP 227	116	SP 228	116	SP 299	135
Perrin-Schmitt, F	OP 60	43	SP 270	127				SP 005	58	SP 006	58		
Persin, L	SP 122	89	SP 022	62	SP 023	62	Reichert, C	SP 297	134				
Persin, LS	SP 204	110	SP 240	119	SP 069	75	Reijers, R	OP 59	43				
	SP 079	77					Reimann, S	OP 52	39	SP 297	134		
Petcu, A	SP 073	76					Reinhardt, R	SP 295	133				
Petrén, S	SP 081	78					Reitsma, J H	OP 48	37				
Petri, N	SP 198	109					Rejc Novak, M	CP 019	151				
Petrick, S	OP 25	25					Remisz, B	SP 035	66				
Petrillo, F	CP 115	173					Ren, Y	SP 032	65	SP 052	70		
Petrovska, J	SP 220	114					Repina, T	SP 204	110				
Petrovski, D	SP 220	114					Reshitaj, A	CP 109	172				
Petruseska, M	CP 094	168					Rexhepi, A	CP 017	150	CP 057	160	CP 058	160
Piana, G	SP 303	136					Rexhepi, S	CP 017	150	CP 057	160	CP 058	160
Piaino, M G	CP 033	154	SP 137	93	SP 138	93	Rheden, R	OP 47	36				
Pichelmayer, M	SP 171	102	SP 273	128			Ribackova, M	SP 204	110				
Pietrzak, P	SP 175	103	SP 265	123			Ricchiuti, M R	SP 214	113	SP 215	113		
Pietrzak-Bilinska, B	SP 216	113					Richmond, S	OP 21	23	OP 38	32	OP 06	15
Pietrzak-Bilińska, B	SP 080	78						SP 100	83	SP 102	83	SP 134	92
Pilotto, A	CP 083	166	SP 180	104				SP 141	94	SP 156	98		
Pirilä-Parkkinen, K	SP 091	80					Richter, V	SP 074	76				
Pirte, A	SP 260	124					Rimes, R	OP 36	31				
Pirttiniemi, P	SP 018	61	SP 205	110	SP 251	122	Roberto, M	OP 23	24				
	SP 042	67	SP 091	80			Rocca, P	SP 201	109				
Pisek, M	OP 38	32					Rodriguez Arnal, F	SP 161	99				
Playle, R	OP 06	15					Römer, P	SP 227	116	SP 228	116	SP 299	135
Poikela, A	SP 251	122						SP 005	58	SP 006	58		
Polat-Özsoy, Ö	OP 33	29	SP 208	111	SP 288	132	Romero Maroto, M	SP 161	99	SP 004	58		
Polimeni, A	CP 115	173	SP 201	109			Roos, M	OP 13	19	SP 158	98		
Polma, L	SP 258	124	SP 259	124	SP 079	77	Rosa, M	SP 135	92	SP 144	94		
Polychronopoulou, A	SP 254	123					Rossini, G	SP 094	81				
Pontarolo, E	SP 229	116					Rozzi, M	SP 121	89				
Pop Stefanova-Trposka, M	SP 119	88	SP 087	79			Ruano Gomez, R	SP 161	99				
Popa, M	SP 128	90					Rubleva, I	SP 069	75				
Popescu, M	SP 101	83	SP 084	79	SP 088	80	Rudblad, S	SP 002	57				
Popova, I	SP 204	110					Ruf, S	SP 055	71				
Popoviciu, O	SP 101	83	SP 129	91	SP 084	79	Ruinpeikar, M	SP 127	90				
	SP 088	80					Rusanen, J	SP 205	110				
Popovski, S	CP 100	170	SP 242	120	SP 087	79	Russel, F	OP 47	36				
Porumb, A	SP 260	124					Ružman Knežević, R	SP 149	95				
Pospieszńska, M	SP 114	87					Saadat, S	SP 197	108	SP 244	120		
Postnikov, M	SP 240	119					Sabuncuoglu, F	CP 120	174				
Pourdanish, F	SP 124	90					Sacomanno, S	CP 025	152	CP 026	152	CP 006	148
Pozzani, E	SP 210	112	SP 303	136			Sader, R	CP 107	171	SP 266	126		
Präger, T M	SP 036	66					Sağdıç, D	SP 320	140				
Prahl-Andersen, B	OP 48	37					Salehi, P	SP 196	108	SP 197	108		

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Sali, F	CP 057	160	CP 094	168			Stanciu, D	SP 088	80				
Salvador, C F A	SP 233	118	SP 239	119			Stanizzi, A	SP 174	102				
Sancak, Ö	SP 271	127					Staudt, CB	SP 316	139				
Sandy, J R	SP 179	104					Stawarczyk, B	SP 158	98				
Sano, N	SP 276	129	SP 278	129			Stec-Slonicz, M	CP 046	157				
Santigli, E	SP 209	112					Stefanović, G	SP 263	125				
Santorio, M	CP 102	170					Stefanović, N	OP 03	14	SP 263	125		
Sapač, A	CP 019	151					Stellini, E	CP 071	163				
Sar, Ç	SP 208	111					Stepanov, G	SP 240	119				
Sarabia, H	CP 110	172					Stevens, M	SP 186	106				
Sari, Z M	SP 181	104					Stojanowski, J	CP 100	170	CP 016	150		
Sarioğlu, M	OP 05	15					Stojanović, D	SP 126	90				
Sarkarat, F	SP 203	110					Stojanović, M	CP 035	155	CP 095	169		
Sarul, M	SP 040	67					Stojković, I	CP 053	159				
Sato, K	CP 059	160					Stojković, M	CP 053	159				
Sauleau, E A	SP 270	127					Stós, W	SP 034	65	SP 035	66		
Scarpa, D	CP 025	152					Stošić, M	SP 126	90				
Scepan, I	SP 249	121					Stoustrup, P	OP 58	42				
Scharstuhl, A	OP 47	36					Suárez, C	SP 014	60				
Schätzle, M	SP 158	98	SP 195	108			Sueishi, K	CP 112	172				
Schmidlin, P R	SP 195	108					Sumbullu, M A	SP 105	85				
Schmidtman, I	SP 284	131					Sumer, M	CP 113	173				
Schulze, M	OP 43	34					Sun, M-K	SP 322	140				
Schuster, G	SP 143	94					Svabova, M	SP 219	114				
Schwartz, C	CP 064	161	CP 067	162			Svanholt, P	SP 198	109				
Scopelliti, D	CP 066	162					Svejda, M	SP 103	83				
Seeborg, J	OP 15	20					Synodinos, F	CP 086	167	CP 087	167		
Selan, D	CP 075	164	SP 102	83			Synodinos, P	CP 114	173	CP 024	152		
Seoane, J	SP 255	123					Syrinska, M	SP 155	98	SP 072	76		
Serbanescu, A	SP 112	86					Szczurowski, P	SP 125	90				
Sever, E	SP 097	82	SP 098	82			Szuhaneck, C	CP 005	147	SP 030	64	SP 083	78
Sevilla, P	SP 014	60						SP 092	81				
Sfeatcu, R	SP 165	100					Szuhaneck, T	CP 007	148				
Sgritta, G	SP 291	132	CP 121	174			Szumielewicz, M	CP 015	150	SP 024	63		
Shah, R	OP 04	14					Szyper-Szczurowska, J	SP 125	90				
Shahidi, S	SP 051	70					Ślaj, M	CP 076	164	SP 046	69		
Sham, P Y E	SP 167	101					Špalj, S	SP 046	69	SP 149	95		
Sherriff, M	SP 177	103	SP 178	103			Štaban, T	SP 118	88	SP 203	110		
Shinohara, Y	SP 222	115					Tagliabue, A	SP 026	63				
Showkatbakhsh, A	SP 124	90	SP 139	93			Takada, J I	CP 014	150				
Showkatbakhsh, R	SP 118	88					Talalaeva, E	SP 079	77				
Sidiropoulou-Chatzigianni, S	SP 217	114					Talpone, F	SP 138	93				
Sieja, A	CP 104	171					Tamer, T	SP 301	135				
Sifakakis, I	SP 221	114					Tan, Y	OP 08	16				
Signorelli, L	OP 31	28					Tan, Y Y	HAP 04	8				
Silvola, A S	SP 205	110					Tanase, G	SP 164	100	SP 165	100		
Sipilä, K	SP 205	110					Taner, T	CP 043	157				
Sisman, C	SP 317	139					Tang, A T H	SP 019	61				
Sjögren, A	OP 34	30					Taoufik, K	SP 254	123				
Skrypnyk, I	OP 015	60					Tarantino, D	CP 115	173				
Slabkovskaya, A	SP 122	89	SP 069	75			Tarmure, V	SP 112	172				
Slette, G	SP 235	118					Tarraf, N E	OP 35	30				
Smiech-Slomkowska, G	SP 175	103					Tastan, H	SP 271	127				
Smileva-Nacevska, M	SP 252	122					Tausche, E	OP 25	25				
Söderfeldt, B	SP 190	107					Tavira, S	CP 082	166				
Sökücü, O	SP 262	125					Tavventi	SP 132	91	SP 133	92		
Solano Reina, E	SP 012	60	SP 029	64			Taylor, P	OP 35	30				
Sönmez, G	CP 047	157					Temmerman, L	SP 238	119				
Sonnesen, L	SP 198	109	SP 063	73			Teodorescu, E	CP 018	151	CP 060	160	SP 123	89
Sorrentino, M	SP 291	132						SP 128	90	SP 129	91		
Soumas, V	SP 316	139					Terada, K	SP 276	129	SP 278	129		
Spaić, G	CP 072	163					Terashima, Y	SP 278	129				
Spary, D	CP 028	153					Tezza, F	SP 210	112				
Spasov, Z	SP 206	111					Thomas, E R	SP 008	89				
Spasov, A	OP 55	41	SP 246	121	SP 247	121	Thongdee, P	SP 017	61				
	SP 304	136					Tian, C	CP 004	147				
Speel, R J	SP 225	115					Tiddeman, B	SP 007	58				
Spidlen, M	SP 248	121					Tiilikainen, P	SP 018	61	SP 042	67		
Spinella, P	SP 291	132					Timm, H	SP 264	125				
Springer, I	OP 43	34					Tindlund, R S	OP 44	35				
Sprogar, Š	SP 268	126					Tokalak, F	CP 009	148				
Stajčić, Z	CP 095	169					Toliat, M	SP 118	88				
Stamenković, Z	CP 050	158	CP 093	168	CP 096	169	Tolonen, U	SP 091	80				
	SP 218	114					Toma, A	OP 21	23	OP 06	15	SP 156	98
Stamm, T	CP 039	156					Tomassini, F	CP 098	169				

Author	Abs	Pg	Abs	Pg	Abs	Pg	Author	Abs	Pg	Abs	Pg	Abs	Pg
Tomažević, T	SP 104	84					Von den Hoff, J W	OP 59	43				
Topouzelis, N	SP 096	82					von Kodolitsch, Y	SP 286	131				
Torensma, R	OP 10	17					Vral, A	SP 238	119				
Torkan, S	CP 091	168	SP 127	90			Vreeke, M	SP 182	105				
Toseska, N	SP 250	122					Vukosavljević, L	SP 263	125				
Toseska-Spasova, N	SP 206	111					Vuličević, T	CP 020	151				
Toth Pirnovar, A	SP 147	95					Wagener, F	OP 47	36				
Toxopeus, A H	SP 186	106					Wagner, W	CP 011	149				
Toygar Memikoğlu, U	SP 142	94					Waldron, J M	SP 179	104				
Tran, J	SP 037	66					Waltimo-Sirén, J	CP 080	165				
Treccani, M	SP 302	135					Walverde, E A	SP 239	119				
Triaca, A	HAO 03	5					Wangwarunyoo, K	SP 154	97				
Tronca, S	CP 118	174	CP 119	174	SP 308	137	Waring, D	SP 172	102				
	SP 309	137	SP 311	138	SP 312	138	Wedrychowska-Szulc, B	SP 110	86				
	SP 313	138	SP 315	139			Wee, E C K	SP 011	59				
Tsaknaki, I	SP 264	125					Wehrbein, H	CP 011	149	OP 29	27	SP 264	125
Tschill, P	OP 60	43						SP 284	131	SP 295	133		
Tsetsilas, M	SP 296	134					Wendl, B	SP 171	102	SP 273	128		
Tsourakis, A K	CP 027	152					Wennström, J	SP 061	73				
Tufano, G	SP 314	138					Whang, C Z Y	SP 178	103				
Tulloch, J F C	OP 41	33	OP 42	34			White, A	HAO 02	4				
Tuna, S H	SP 176	103					Willems, G	OP 18	22	SP 183	105	SP 256	123
Tunçbilek, G	CP 041	156					Winkler, A	CP 071	163	SP 293	133		
Türk, T	CP 113	173					Winsauer, H	SP 273	128				
Turk, T	OP 16	20	SP 277	129			Wojtaszek-Słomińska, A	SP 211	112				
Turkkahraman, H	OP 05	15	SP 176	103	SP 287	131	Wong, R	SP 153	97	SP 154	97		
	SP 095	81					Wong, R W K	HAP 04	8	HAP 05	8	OP 08	16
	CP 034	154						SP 109	86	SP 016	61	SP 017	61
Uğurlu, K	CP 047	157						SP 192	107	SP 021	62	SP 028	64
Ulker, E	SP 272	128					Wozniak, K	SP 155	98	SP 072	76		
Ulku, Z	OP 31	28					Wriedt, S	CP 011	149	OP 29	27		
Ullrich, O	SP 230	117					Wu, A T	SP 301	135				
Undas, A	SP 208	111					Xiao, D	SP 052	70				
Ünver, B	SP 039	67					Yajima, E	SP 278	129				
Urban, A	SP 207	111					Yamaaki, F	SP 020	62				
Urbanova, W	SP 283	130					Yanagisawa, M	CP 059	160	CP 059	160		
Uršič-Bratina, N	SP 115	87	SP 189	106			Yang, B	SP 107	85				
Uslu Akçam, O	SP 181	104	SP 062	73			Yang, B H	SP 191	107				
Usumez, S	SP 317	139	SP 320	140	SP 062	73	Yang, Y	OP 08	16				
Uysal, T	SP 166	100					Yang, Y Q	HAP 04	8	SP 192	107	SP 028	64
Vagapov, Z	SP 286	131					Yavorovkaya, T	SP 085	79				
Vahle-Hinz, E	SP 260	124					Yavorovskaya, T	SP 086	79				
Vaida, L	SP 083	78					Yoon, H R	CP 108	171				
Valceanu, A	SP 138	93					Yoon, J W	SP 319	140				
Vallelonga, T	OP 09	17					Yoon, M S	SP 107	85				
Valletta, R	SP 187	106					Yosefipoor, E	SP 160	99				
van der Kaaij, N C W	SP 032	65					Yu, H S	SP 116	87	SP 117	88		
van der Mei, H C	SP 186	106	SP 187	106			Yugaraj, P	HAP 05	8				
van der Veen, M H	SP 232	117					Yun, M S	SP 191	107				
van der Vlag, J	SP 058	72					Yurov, V	SP 069	75				
Van Parys, K	SP 213	113					Zadurska, M	SP 080	78				
Van Sint Jan, S	OP 19	22					Zaletel, E	CP 061	161				
van Vlijmen, O	SP 188	106					Zamiri, B	SP 051	70				
van Westing, K	SP 033	65	SP 089	80			Zandi, S	SP 160	99				
Vande Vannet, B	SP 049	70	SP 050	70			Zapala, J	SP 125	90				
Vangelisti, A	SP 207	111					Zavareian, S	SP 106	85				
Vankova, Z	SP 213	113					Zecevic, B	CP 053	159				
Vanmuylder, N	SP 025	63					Zentner, A	OP 45	35	OP 56	41	SP 182	105
Varga, G	CP 114	173	CP 024	152				SP 231	117	SP 232	117	SP 058	72
Vasilakou, N	SP 153	97					Zetu, I N	SP 193	107	SP 027	64	SP 073	76
Veerawattanatigul, A	OP 07	16	SP 120	88	SP 025	63	Zhu, S J	SP 028	64				
Végh, A	SP 097	82	SP 098	82			Zhurov, A	OP 21	23	OP 06	15	SP 100	83
Vehovec, E	CP 068	162	CP 075	164	SP 102	83		SP 102	83	SP 141	94	SP 156	98
Verdenik, M	SP 164	100					Ziebur, T	CP 039	156				
Vereanu, A	CP 073	163					Ziętek, M	SP 082	78				
Verhovec, K	CP 049	158					Zimmermann, A	OP 20	23				
Verly, N	SP 169	101					Zinchenko, A	SP 204	110				
Vernière, A	OP 10	17					Zotti, F	CP 022	151	CP 004	147	SP 306	136
Verstappen, J	CP 076	164						SP 307	137				
Vidaković, R	SP 014	60					Zuo, P	HAP 05	8	SP 021	62		
Vilar, T	SP 172	102					Zupančič, S	SP 199	109				
Visram, S	CP 102	170	OP 46	36			Zuzelova, M	SP 252	122				
Vlachojannis, J E	CP 029	153	CP 068	162	CP 070	163							
Volk, J	OP 26	26	OP 61	44	SP 099	82							
Von den Hoff, H	OP 10	17	OP 14	20									

